

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07D 265/32, 279/12, 413/06, 413/04, 413/14	A1	(11) International Publication Number: WO 95/16679 (43) International Publication Date: 22 June 1995 (22.06.95)
(21) International Application Number: PCT/US94/14497 (22) International Filing Date: 13 December 1994 (13.12.94) (30) Priority Data: 169,889 17 December 1993 (17.12.93) US (60) Parent Application or Grant (63) Related by Continuation US 169,889 (CIP) Filed on 17 December 1993 (17.12.93) (71) Applicant (for all designated States except US): MERCK & CO., INC. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): DORN, Conrad, P. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). FINKE, Paul, E. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). HALE, Jeffrey, J. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). MacCOSS, Malcolm [GB/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). MILLS, Sander, G. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). SHAH, Shrenik,	K. [IN/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). CHAMBERS, Mark, Stuart [GB/GB]; Terlings Park, Eastwick Road, Harlow, Essex CM20 2QR (GB). HARRISON, Timothy [GB/GB]; Terlings Park, Eastwick Road, Harlow, Essex CM20 2QR (GB). LADDUWA-HETTY, Tamara [GB/GB]; Terlings Park, Eastwick Road, Harlow, Essex CM20 2QR (GB). WILLIAMS, Brian, John [GB/GB]; Terlings Park, Eastwick Road, Harlow, Essex CM20 2QR (GB). (74) Common Representative: MERCK & CO., INC.; Patent Dept., 126 East Lincoln Avenue, Rahway, NJ 07065 (US). (81) Designated States: AM, AU, BB, BG, BR, BY, CA, CN, CZ, EE, FI, GE, HU, JP, KG, KR, KZ, LK, LR, LT, LV, MD, MG, MN, NO, NZ, PL, RO, RU, SI, SK, TJ, TT, UA, US, UZ, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, MW, SD, SZ). Published <i>With international search report.</i>	
(54) Title: MORPHOLINE AND THIOMORPHOLINE TACHYKININ RECEPTOR ANTAGONISTS		
<div style="text-align: center;"> <p>(I)</p> </div>		
(57) Abstract <p>Substituted heterocycles of general structural formula (I) or a pharmaceutically acceptable salt thereof, wherein: R¹ is selected from the group consisting of: hydrogen; C₁₋₆ alkyl, unsubstituted or substituted; C₂₋₆ alkenyl, unsubstituted or substituted; C₂₋₆ alkynyl; phenyl, unsubstituted or substituted; X is selected from the group consisting of: -O-, -S-, -SO-, -SO₂-; Y is selected from the group consisting of: a single bond, -O-, -S-, -CO-, -CH₂-, -CHR₅- and -CR¹⁵R¹⁶-, Z = C₁₋₆ alkyl, are tachykinin receptor antagonists useful in the treatment of inflammatory diseases, pain or migraine, asthma and emesis, and calcium channel blockers useful in the treatment of cardiovascular conditions such as angina, hypertension or ischemia.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

- 1 -

TITLE OF THE INVENTIONMORPHOLINE AND THIOMORPHOLINE TACHYKININ
RECEPTOR ANTAGONISTS5 BACKGROUND OF THE INVENTION

Analgesia has historically been achieved in the central nervous system by opiates and analogs which are addictive, and peripherally by cyclooxygenase inhibitors that have gastric side effects. Substance P antagonists may induce analgesia both centrally and
10 peripherally. In addition, substance P antagonists are inhibitory of neurogenic inflammation.

The neuropeptide receptors for substance P (neurokinin-1; NK-1) are widely distributed throughout the mammalian nervous system (especially brain and spinal ganglia), the circulatory system and
15 peripheral tissues (especially the duodenum and jejunum) and are involved in regulating a number of diverse biological processes. This includes sensory perception of olfaction, vision, audition and pain, movement control, gastric motility, vasodilation, salivation, and micturition (B. Pernow, Pharmacol. Rev., 1983, 35, 85-141). The
20 NK-1 and NK-2 receptor subtypes are implicated in synaptic transmission (Laneuville et al., Life Sci., 42: 1295-1305 (1988)).

The receptor for substance P is a member of the superfamily of G protein-coupled receptors. This superfamily is an extremely diverse group of receptors in terms of activating ligands and
25 biological functions. In addition to the tachykinin receptors, this receptor superfamily includes the opsins, the adrenergic receptors, the muscarinic receptors, the dopamine receptors, the serotonin receptors, a thyroid-stimulating hormone receptor, a luteinizing hormone-choriogonadotropic hormone receptor, the product of the oncogene ras,
30 the yeast mating factor receptors, a Dictyostelium cAMP receptor, and receptors for other hormones and neurotransmitters (see A.D. Hershey, et al., J. Biol. Chem., 1991, 226, 4366-4373).

Substance P (also called "SP" herein) is a naturally occurring undecapeptide belonging to the tachykinin family of peptides,

- 2 -

the latter being so-named because of their prompt contractile action on extravascular smooth muscle tissue. The tachykinins are distinguished by a conserved carboxyl-terminal sequence Phe-X-Gly-Leu-Met-NH₂. In addition to SP the known mammalian tachykinins include neurokinin A and neurokinin B. The current nomenclature designates the receptors for SP, neurokinin A, and neurokinin B as NK-1, NK-2, and NK-3, respectively.

More specifically, substance P is a pharmacologically-active neuropeptide that is produced in mammals and possesses a characteristic amino acid sequence (Chang *et al.*, Nature New Biol. **232**, 86 (1971); D.F. Veber *et al.*, U.S. Patent No. 4,680,283).

Substance P is a pharmacologically-active neuropeptide that is produced in mammals and acts as a vasodilator, a depressant, stimulates salivation and produces increased capillary permeability. It is also capable of producing both analgesia and hyperalgesia in animals, depending on dose and pain responsiveness of the animal (see R.C.A. Frederickson *et al.*, Science, **199**, 1359 (1978); P. Oehme *et al.*, Science, **208**, 305 (1980)) and plays a role in sensory transmission and pain perception (T.M. Jessell, Advan. Biochem. Psychopharmacol. **28**, 189 (1981)). For example, substance P is believed to be involved in the neurotransmission of pain sensations [Otsuka *et al.*, "Role of Substance P as a Sensory Transmitter in Spinal Cord and Sympathetic Ganglia" in 1982 Substance P in the Nervous System, Ciba Foundation Symposium 91, 13-34 (published by Pitman) and Otsuka and Yanagisawa, "Does Substance P Act as a Pain Transmitter?" TIPS, **8** 506-510 (Dec. 1987)], specifically in the transmission of pain in migraine (see B.E.B. Sandberg *et al.*, Journal of Medicinal Chemistry, **25**, 1009 (1982); M. A. Moskowitz, Trends Pharmacol. Sci., **13**, 307-311 (1992)), and in arthritis (Levine, *et al.* Science, **226** 547-549 (1984); M. Lotz, *et al.*, Science, **235**, 893-895 (1987)). Tachykinins have also been implicated in gastrointestinal (GI) disorders and diseases of the GI tract, such as inflammatory bowel disease [see Mantyh *et al.*, Neuroscience, **25** (3), 817-37 (1988) and D. Regoli in "Trends in Cluster Headache" Ed. F. Sicuteri *et al.*, Elsevier Scientific Publishers, Amsterdam, pp. 85-95

- 3 -

(1987)], and emesis [Trends Pharmacol. Sci., 9, 334-341 (1988), F.D. Tattersall, et al., Eur. J. Pharmacol., 250, R5-R6 (1993)].

It is also hypothesized that there is a neurogenic mechanism for arthritis in which substance P may play a role [Kidd et al., "A
5 Neurogenic Mechanism for Symmetric Arthritis" in The Lancet, 11 November 1989 and Gronblad et al., "Neuropeptides in Synovium of Patients with Rheumatoid Arthritis and Osteoarthritis" in J. Rheumatol.
15 15(12) 1807-10 (1988)]. Therefore, substance P is believed to be involved in the inflammatory response in diseases such as rheumatoid
10 arthritis and osteoarthritis [O'Byrne et al., Arthritis and Rheumatism, 33 1023-8 (1990)].

Evidence for the usefulness of tachykinin receptor antagonists in pain, headache, especially migraine, Alzheimer's disease, multiple sclerosis, attenuation of morphine withdrawal, cardiovascular
15 changes, oedema, such as oedema caused by thermal injury, chronic inflammatory diseases such as rheumatoid arthritis, asthma/bronchial hyperreactivity and other respiratory diseases including allergic rhinitis, inflammatory diseases of the gut including ulcerative colitis and
20 Crohn's disease, ocular injury and ocular inflammatory diseases, proliferative vitreoretinopathy, irritable bowel syndrome and disorders of bladder function including cystitis and bladder detrusor hyperreflexia is reviewed in "Tachykinin Receptors and Tachykinin Receptor
Antagonists," C.A. Maggi, R. Patacchini, P. Rovero and A. Giachetti, J. Auton. Pharmacol., 13, 23-93 (1993); see also R. M. Snider, et al.,
25 Chem. Ind., 11, 792-794 (1991). Neurokinin-1 receptor antagonists alone or in combination with bradykinin receptor antagonists may also be useful in the prevention and treatment of inflammatory conditions in the lower urinary tract, especially cystitis [Giuliani, et al., J. Urology,
30 150, 1014-1017 (1993)]. Other disease areas where tachykinin antagonists are believed to be useful are allergic conditions [Hamelet et al., Can. J. Pharmacol. Physiol., 66, 1361-7 (1988)], immunoregulation [Lotz, et al., Science, 241 1218-21 (1988), Kimball, et al., J. Immunol., 141 (10) 3564-9 (1988); A. Perianin, et al., Biochem. Biophys. Res Commun. 161, 520 (1989)], post-operative pain and nausea [C. Bountra,

- 4 -

et al., Eur. J. Pharmacol., 249, R3-R4 (1993), F. D. Tattersall, et al., Neuropharmacology, 33, 259-260 (1994)], vasodilation, bronchospasm, reflex or neuronal control of the viscera [Mantyh et al., PNAS, 85, 3235-9 (1988)] and, possibly by arresting or slowing β -amyloid-mediated neurodegenerative changes [Yankner et al., Science, 250, 279-82 (1990)] in senile dementia of the Alzheimer type, Alzheimer's disease and Downs Syndrome. Substance P may also play a role in demyelinating diseases such as multiple sclerosis and amyotrophic lateral sclerosis [J. Lubner-Narod, et al., poster C.I.N.P. XVIIIth Congress, 28th June-2nd July, 1992], and in disorders of bladder function such as bladder detrusor hyper-reflexia [Lancet, 16th May 1992, 1239]. Antagonists selective for the neurokinin-1 (NK-1) and/or the neurokinin-2 (NK-2) receptor may be useful in the treatment of asthmatic disease (Frossard et al., Life Sci., 49, 1941-1953 (1991); Advenier, et al., Biochem. Biophys. Res. Comm., 184(3), 1418-1424 (1992); P. Barnes, et al., Trends Pharmacol. Sci., 11, 185-189 (1993)). Tachykinin antagonists may also be useful in the treatment of small cell carcinomas, in particular small cell lung cancer (SCLC) [Langdon et al., Cancer Research, 52, 4554-7 (1992)].

It has furthermore been suggested that tachykinin receptor antagonists have utility in the following disorders: depression, dysthymic disorders, chronic obstructive airways disease, hypersensitivity disorders such as poison ivy, vasospastic diseases such as angina and Reynauld's disease, fibrosing and collagen diseases such as scleroderma and eosinophilic fasciitis, reflex sympathetic dystrophy such as shoulder/hand syndrome, addiction disorders such as alcoholism, stress related somatic disorders, neuropathy, neuralgia, disorder related to immune enhancement or suppression such as systemic lupus erythematous (EPO Publication No. 0,436,334), ophthalmic diseases such as conjunctivitis, vernal conjunctivitis, and the like, and cutaneous diseases such as contact dermatitis, atopic dermatitis, urticaria, and other eczematoid dermatitis (EPO Publication No. 0,394,989).

Substance P antagonists may be useful in mediating neurogenic mucus secretion in mammalian airways and hence provide

- 5 -

treatment and symptomatic relief in diseases characterized by mucus secretion, in particular, cystic fibrosis [S. Ramnarine, et al., abstract presented at 1993 ALA/ATS Int'l Conference, 16-19 May, 1993, published in Am. Rev. of Respiratory Dis., May 1993].

5 In the recent past, some attempts have been made to provide peptide-like substances that are antagonists for the receptors of substance P and other tachykinin peptides in order to more effectively treat the various disorders and diseases mentioned above. For example, Lowe, Drugs of the Future, 17 (12) 1115-1121 (1992) and EPO
10 Publication Nos. 0,347,802, 0,401,177 and 0,412,452 disclose various peptides as neurokinin A antagonists. Also, PCT Patent Publication WO 93/14113 discloses certain peptides as tachykinin antagonists. In addition, EPO Publication No. 0,336,230 discloses heptapeptides which are substance P antagonists useful in the treatment of asthma. Merck
15 U.S. Patent No. 4,680,283 also discloses peptidal analogs of substance P. Certain inhibitors of tachykinins have been described in U.S. Patent No. 4,501,733, by replacing residues in substance P sequence by Trp residues. A further class of tachykinin receptor antagonists, comprising a monomeric or dimeric hexa- or heptapeptide unit in linear or cyclic
20 form, is described in GB-A-2216529.

The peptide-like nature of such substances make them too labile from a metabolic point of view to serve as practical therapeutic agents in the treatment of disease. The non-peptidic antagonists of the present invention, on the other hand, do not possess this drawback, as
25 they are expected to be more stable from a metabolic point of view than the previously-discussed agents.

It is known that in the central nervous system baclofen [β -(aminoethyl)-4-chlorobenzenepropanoic acid] effectively blocks the excitatory activity of substance P, but because in many areas the
30 excitatory responses to other compounds such as acetylcholine and glutamate are inhibited as well, baclofen is not considered a specific substance P antagonist. Pfizer WIPO patent applications (PCT Publication Nos. WO 90/05525, WO 90/05729, WO 91/18899, WO 92/12151 and WO 92/12152) and publications (Science, 251, 435-437

- 6 -

(1991); Science, 251, 437-439 (1991); J. Med. Chem., 35, 2591-2600 (1992)) disclose 2-arylmethyl-3-substituted amino-quinuclidine derivatives which are disclosed as being useful as substance P antagonists for treating gastrointestinal disorders, central nervous system disorders, inflammatory diseases and pain or migraine.

5 A Glaxo European patent application (EPO Publication No. 0.360.390) discloses various spirolactam-substituted amino acids and peptides which are antagonists or agonists of substance P. A Pfizer WIPO patent application (PCT Publication No. WO 92/06079) discloses fused-ring

10 analogs of nitrogen-containing nonaromatic heterocycles as useful for the treatment of diseases mediated by an excess of substance P. A Pfizer WIPO patent application (PCT Publication No. WO 92/15585) discloses 1-azabicyclo[3.2.2]nonan-3-amine derivatives as substance P antagonists. A Pfizer WIPO patent application (PCT Publication No. WO 93/10073)

15 discloses ethylenediamine derivatives as substance P antagonists. PCT Publication No. WO 93/01169 discloses certain aromatic compounds as tachykinin receptor antagonists. A Sanofi publication (Life Sci., 50, PL101-PL106 (1992)) discloses a 4-phenyl piperidine derivative as an antagonist of the neurokinin A (NK2) receptor.

20 Howson et al. (Biorg. & Med. Chem. Lett., 2 (6), 559-564 (1992)) disclose certain 3-amino and 3-oxy quinuclidine compounds and their binding to substance P receptors. EPO Publication 0.499.313 discloses certain 3-oxy and 3-thio azabicyclic compounds as tachykinin antagonists. U.S. Patent No. 3,506,673 discloses certain 3-hydroxy

25 quinuclidine compounds as central nervous system stimulants. A Pfizer EPO Patent application (EPO Publication 0.436.334) discloses certain 3-aminopiperidine compounds as substance P antagonists. U.S. Patent No. 5,064,838 discloses certain 1,4-disubstituted piperidinyl compounds as analgesics. PCT Publication No. WO 92/12128 discloses certain


30 piperidine and pyrrolidine compounds as analgesics. Peyronel, et al. (Biorg & Med. Chem. Lett., 2 (1), 37-40 (1992)) disclose a fused ring pyrrolidine compound as a substance P antagonist. EPO Publication No. 0.360.390 discloses certain spirolactam derivatives as substance P antagonists. U.S. Patent No. 4,804,661 discloses certain piperazine

- 7 -

compounds as analgesics. U.S. Patent No. 4,943,578 discloses certain piperazine compounds useful in the treatment of pain. PCT Publication No. WO 92/01679 discloses certain 1,4-disubstituted piperazines useful in the treatment of mental disorders in which a dopaminergic deficit is implicated. PCT Publication No. WO 94/00440 discloses certain morpholine compounds as substance P antagonists.

SUMMARY OF THE INVENTION

This invention is concerned with novel compounds represented by structural formula I:



wherein R¹, R², R³, R⁴, R⁵, and X are hereinafter defined.

The invention is also concerned with pharmaceutical formulations comprising these novel compounds as active ingredients and the use of the novel compounds and their formulations in the treatment of certain disorders.


The compounds of this invention are tachykinin receptor antagonists and are useful in the treatment of inflammatory diseases, pain or migraine, asthma and emesis.

Also, some of these compounds are calcium channel blockers and are useful in the treatment of cardiovascular disorders such as angina, hypertension or ischemia.

- 8 -

DETAILED DESCRIPTION OF THE INVENTION

The novel compounds of this invention are represented by structural formula I:



or a pharmaceutically acceptable salt thereof, wherein:

R¹ is selected from the group consisting of:

- (1) hydrogen;
- (2) C₁-6 alkyl, unsubstituted or substituted with one or more of the substituents selected from:
 - (a) hydroxy,
 - (b) oxo,
 - (c) C₁-6 alkoxy,
 - (d) phenyl-C₁-3 alkoxy,
 - (e) phenyl,
 - (f) -CN,
 - (g) halo, wherein halo is fluoro, chloro, bromo or iodo,
 - (h) -NR⁹R¹⁰, wherein R⁹ and R¹⁰ are independently selected from:
 - (i) hydrogen,
 - (ii) C₁-6 alkyl,
 - (iii) hydroxy-C₁-6 alkyl, and
 - (iv) phenyl,
 - (i) -NR⁹COR¹⁰, wherein R⁹ and R¹⁰ are as defined above,
 - (j) -NR⁹CO₂R¹⁰, wherein R⁹ and R¹⁰ are as defined above,
 - (k) -CONR⁹R¹⁰, wherein R⁹ and R¹⁰ are as defined above,
 - (l) -COR⁹, wherein R⁹ is as defined above,

- 9 -

- (m) $-\text{CO}_2\text{R}^9$, wherein R^9 is as defined above;
- (n) heterocycle, wherein the heterocycle is selected from the group consisting of:
- (A) benzimidazolyl,
 - (B) benzofuranyl,
 - (C) benzothiophenyl,
 - (D) benzoxazolyl,
 - (E) furanyl,
 - (F) imidazolyl,
 - (G) indolyl,
 - (H) isooxazolyl,
 - (I) isothiazolyl,
 - (J) oxadiazolyl,
 - (K) oxazolyl,
 - (L) pyrazinyl,
 - (M) pyrazolyl,
 - (N) pyridyl,
 - (O) pyrimidyl,
 - (P) pyrrolyl,
 - (Q) quinolyl,
 - (R) tetrazolyl,
 - (S) thiadiazolyl,
 - (T) thiazolyl,
 - (U) thienyl,
 - (V) triazolyl,
 - (W) azetidiny,
 - (X) 1,4-dioxanyl,
 - (Y) hexahydroazepinyl,
 - (Z) piperazinyl,
 - (AA) piperidiny,
 - (AB) pyrrolidinyl,
 - (AC) tetrahydrofuranyl, and
 - (AD) tetrahydrothienyl,

and wherein the heterocycle is unsubstituted or substituted with

- 10 -

one or more substituent(s) selected from:

- 5
- (i) C₁-6 alkyl, unsubstituted or substituted with halo, -CF₃, -OCH₃, or phenyl,
 - (ii) C₁-6 alkoxy,
 - (iii) oxo,
 - (iv) hydroxy,
 - (v) thioxo,
 - (vi) -SR⁹, wherein R⁹ is as defined above,
 - 10 (vii) halo,
 - (viii) cyano,
 - (ix) phenyl,
 - (x) trifluoromethyl,
 - (xi) -(CH₂)_m-NR⁹R¹⁰, wherein m is 0, 1 or 2, and R⁹ and R¹⁰ are as defined above,
 - 15 (xii) -NR⁹COR¹⁰, wherein R⁹ and R¹⁰ are as defined above,
 - (xiii) -CONR⁹R¹⁰, wherein R⁹ and R¹⁰ are as defined above,
 - 20 (xiv) -CO₂R⁹, wherein R⁹ is as defined above, and
 - (xv) -(CH₂)_m-OR⁹, wherein m and R⁹ are as defined above;

25 (3) C₂-6 alkenyl, unsubstituted or substituted with one or more of the substituent(s) selected from:

- (a) hydroxy,
- (b) oxo,
- (c) C₁-6 alkoxy,
- 30 (d) phenyl-C₁-3 alkoxy,
- (e) phenyl,
- (f) -CN,
- (g) halo,
- (h) -CONR⁹R¹⁰ wherein R⁹ and R¹⁰ are as defined above,

- 11 -

- (i) -COR⁹ wherein R⁹ is as defined above,
 - (j) -CO₂R⁹, wherein R⁹ is as defined above,
 - (k) heterocycle, wherein the heterocycle is as defined above;
- 5 (4) C₂-6 alkynyl;
- (5) phenyl, unsubstituted or substituted with one or more of the substituent(s) selected from:
- (a) hydroxy,
 - (b) C₁-6 alkoxy,
 - 10 (c) C₁-6 alkyl,
 - (d) C₂-5 alkenyl,
 - (e) halo,
 - (f) -CN,
 - (g) -NO₂,
 - 15 (h) -CF₃,
 - (i) -(CH₂)_m-NR⁹R¹⁰, wherein m, R⁹ and R¹⁰ are as defined above,
 - (j) -NR⁹COR¹⁰, wherein R⁹ and R¹⁰ are as defined above,
 - 20 (k) -NR⁹CO₂R¹⁰, wherein R⁹ and R¹⁰ are as defined above,
 - (l) -CONR⁹R¹⁰, wherein R⁹ and R¹⁰ are as defined above,
 - 25 (m) -CO₂NR⁹R¹⁰, wherein R⁹ and R¹⁰ are as defined above,
 - (n) -COR⁹, wherein R⁹ is as defined above;
 - (o) -CO₂R⁹, wherein R⁹ is as defined above;
- 30 R² and R³ are independently selected from the group consisting of:
- (1) hydrogen,
 - (2) C₁-6 alkyl, unsubstituted or substituted with one or more of the substituents selected from:
 - (a) hydroxy,
 - (b) oxo,

- 12 -

- 5
- (c) C₁₋₆ alkoxy,
 - (d) phenyl-C₁₋₃ alkoxy,
 - (e) phenyl,
 - (f) -CN,
 - (g) halo,
 - (h) -NR⁹R¹⁰, wherein R⁹ and R¹⁰ are as defined above,
 - (i) -NR⁹COR¹⁰, wherein R⁹ and R¹⁰ are as defined above,
 - 10 (j) -NR⁹CO₂R¹⁰, wherein R⁹ and R¹⁰ are as defined above,
 - (k) -CONR⁹R¹⁰, wherein R⁹ and R¹⁰ are as defined above,
 - 15 (l) -COR⁹, wherein R⁹ is as defined above, and
 - (m) -CO₂R⁹, wherein R⁹ is as defined above;
- (3) C₂₋₆ alkenyl, unsubstituted or substituted with one or more of the substituent(s) selected from:
- 20
- (a) hydroxy,
 - (b) oxo,
 - (c) C₁₋₆ alkoxy,
 - (d) phenyl-C₁₋₃ alkoxy,
 - (e) phenyl,
 - (f) -CN,
 - 25 (g) halo,
 - (h) -CONR⁹R¹⁰ wherein R⁹ and R¹⁰ are as defined above,
 - (i) -COR⁹ wherein R⁹ is as defined above,
 - (j) -CO₂R⁹, wherein R⁹ is as defined above;
- 30
- (4) C₂₋₆ alkynyl;
- (5) phenyl, unsubstituted or substituted with one or more of the substituent(s) selected from:
- (a) hydroxy,
 - (b) C₁₋₆ alkoxy,

- 13 -

- 5
- (c) C₁₋₆ alkyl,
 - (d) C₂₋₅ alkenyl,
 - (e) halo,
 - (f) -CN,
 - (g) -NO₂,
 - (h) -CF₃,
 - (i) -(CH₂)_m-NR⁹R¹⁰, wherein m, R⁹ and R¹⁰ are as defined above,
 - 10 (j) -NR⁹COR¹⁰, wherein R⁹ and R¹⁰ are as defined above,
 - (k) -NR⁹CO₂R¹⁰, wherein R⁹ and R¹⁰ are as defined above,
 - (l) -CONR⁹R¹⁰, wherein R⁹ and R¹⁰ are as defined above,
 - 15 (m) -CO₂NR⁹R¹⁰, wherein R⁹ and R¹⁰ are as defined above,
 - (n) -COR⁹, wherein R⁹ is as defined above;
 - (o) -CO₂R⁹, wherein R⁹ is as defined above;

20 and the groups R¹ and R² may be joined together to form a heterocyclic ring selected from the group consisting of:

- 25
- (a) pyrrolidinyl,
 - (b) piperidinyl,
 - (c) pyrrolyl,
 - (d) pyridinyl,
 - (e) imidazolyl,
 - (f) oxazolyl, and
 - (g) thiazolyl,

30 and wherein the heterocyclic ring is unsubstituted or substituted with one or more substituent(s) selected from:

- (i) C₁₋₆alkyl,
- (ii) oxo,
- (iii) C₁₋₆alkoxy,
- (iv) -NR⁹R¹⁰, wherein R⁹ and R¹⁰ are as defined

- 14 -

above,

(v) halo, and

(vi) trifluoromethyl;

5 and the groups R² and R³ may be joined together to form a carbocyclic ring selected from the group consisting of:

(a) cyclopentyl,

(b) cyclohexyl,

(c) phenyl,

10 and wherein the carbocyclic ring is unsubstituted or substituted with one or more substituents selected from:

(i) C₁₋₆alkyl,

(ii) C₁₋₆alkoxy,

15 (iii) -NR⁹R¹⁰, wherein R⁹ and R¹⁰ are as defined above,

(iv) halo, and

v) trifluoromethyl;

20 and the groups R² and R³ may be joined together to form a heterocyclic ring selected from the group consisting of:

(a) pyrrolidinyl,

(b) piperidinyl,

(c) pyrrolyl,

25 (d) pyridinyl,

(e) imidazolyl,

(f) furanyl,

(g) oxazolyl,

(h) thienyl, and

30 (i) thiazolyl,

and wherein the heterocyclic ring is unsubstituted or substituted with one or more substituent(s) selected from:

(i) C₁₋₆alkyl,

(ii) oxo,

(iii) C₁₋₆alkoxy,

- 15 -


- (iv) $-NR^9R^{10}$, wherein R^9 and R^{10} are as defined above,
- (v) halo, and
- (vi) trifluoromethyl;

5

R^4 is selected from the group consisting of:

(1)

10



15

(2) $-Y-C_{1-8}$ alkyl, wherein the alkyl is unsubstituted or substituted with one or more of the substituents selected from:

20

- (a) hydroxy,
- (b) oxo,
- (c) C_{1-6} alkoxy,
- (d) phenyl- C_{1-3} alkoxy,
- (e) phenyl,
- (f) $-CN$,
- (g) halo,
- (h) $-NR^9R^{10}$, wherein R^9 and R^{10} are as defined above,
- (i) $-NR^9COR^{10}$, wherein R^9 and R^{10} are as defined above,
- (j) $-NR^9CO_2R^{10}$, wherein R^9 and R^{10} are as defined above,
- (k) $-CONR^9R^{10}$, wherein R^9 and R^{10} are as defined above,
- (l) $-COR^9$, wherein R^9 is as defined above, and
- (m) $-CO_2R^9$, wherein R^9 is as defined above;

25

30

- 16 -

(3) -Y-C₂₋₆ alkenyl, wherein the alkenyl is unsubstituted or substituted with one or more of the substituent(s) selected from:

- 5
- (a) hydroxy,
 - (b) oxo,
 - (c) C₁₋₆ alkoxy,
 - (d) phenyl-C₁₋₃ alkoxy,
 - (e) phenyl,
 - 10 (f) -CN,
 - (g) halo,
 - (h) -CONR⁹R¹⁰ wherein R⁹ and R¹⁰ are as defined above,
 - (i) -COR⁹ wherein R⁹ is as defined above,
 - 15 (j) -CO₂R⁹, wherein R⁹ is as defined above;

(4) -O(CO)-phenyl, wherein the phenyl is unsubstituted or substituted with one or more of R⁶, R⁷ and R⁸;

20 R⁵ is selected from the group consisting of:

- (1) phenyl, unsubstituted or substituted with one or more of R¹¹, R¹² and R¹³;
- (2) C₁₋₈ alkyl, unsubstituted or substituted with one or more of the substituents selected from:

25

 - (a) hydroxy,
 - (b) oxo,
 - (c) C₁₋₆ alkoxy,
 - (d) phenyl-C₁₋₃ alkoxy,
 - (e) phenyl,
 - 30 (f) -CN,
 - (g) halo,
 - (h) -NR⁹R¹⁰, wherein R⁹ and R¹⁰ are as defined above,
 - (i) -NR⁹COR¹⁰, wherein R⁹ and R¹⁰ are as defined above,

- 17 -

- (j) $-NR^9CO_2R^{10}$, wherein R^9 and R^{10} are as defined above,
(k) $-CONR^9R^{10}$, wherein R^9 and R^{10} are as defined above,
5 (l) $-COR^9$, wherein R^9 is as defined above, and
(m) $-CO_2R^9$, wherein R^9 is as defined above;
- (3) C2-6 alkenyl, unsubstituted or substituted with one or more of the substituent(s) selected from:
10 (a) hydroxy,
(b) oxo,
(c) C1-6 alkoxy,
(d) phenyl-C1-3 alkoxy,
(e) phenyl,
15 (f) $-CN$,
(g) halo,
(h) $-CONR^9R^{10}$ wherein R^9 and R^{10} are as defined above,
(i) $-COR^9$ wherein R^9 is as defined above,
20 (j) $-CO_2R^9$, wherein R^9 is as defined above;
- (4) heterocycle, wherein the heterocycle is as defined above;
- 25 R^6 , R^7 and R^8 are independently selected from the group consisting of:
(1) hydrogen;
(2) C1-6 alkyl, unsubstituted or substituted with one or more of the substituents selected from:
30 (a) hydroxy,
(b) oxo,
(c) C1-6 alkoxy,
(d) phenyl-C1-3 alkoxy,
(e) phenyl,
(f) $-CN$,

- 18 -

- (g) halo,
- (h) $-NR^9R^{10}$, wherein R^9 and R^{10} are as defined above,
- 5 (i) $-NR^9COR^{10}$, wherein R^9 and R^{10} are as defined above,
- (j) $-NR^9CO_2R^{10}$, wherein R^9 and R^{10} are as defined above,
- (k) $-CONR^9R^{10}$, wherein R^9 and R^{10} are as defined above,
- 10 (l) $-COR^9$, wherein R^9 is as defined above, and
- (m) $-CO_2R^9$, wherein R^9 is as defined above;
- (3) C₂-6 alkenyl, unsubstituted or substituted with one or more of the substituent(s) selected from:
- 15 (a) hydroxy,
- (b) oxo,
- (c) C₁-6 alkoxy,
- (d) phenyl-C₁-3 alkoxy,
- (e) phenyl,
- 20 (f) -CN,
- (g) halo,
- (h) $-CONR^9R^{10}$ wherein R^9 and R^{10} are as defined above,
- (i) $-COR^9$ wherein R^9 is as defined above,
- (j) $-CO_2R^9$, wherein R^9 is as defined above;
- 25 (4) C₂-6 alkynyl;
- (5) phenyl, unsubstituted or substituted with one or more of the substituent(s) selected from:
- 30 (a) hydroxy,
- (b) C₁-6 alkoxy,
- (c) C₁-6 alkyl,
- (d) C₂-5 alkenyl,
- (e) halo,
- (f) -CN,
- (g) -NO₂,

- 19 -

- 5 (h) $-\text{CF}_3$,
(i) $-(\text{CH}_2)_m-\text{NR}^9\text{R}^{10}$, wherein m , R^9 and R^{10}
are as defined above,
(j) $-\text{NR}^9\text{COR}^{10}$, wherein R^9 and R^{10} are as
defined above,
(k) $-\text{NR}^9\text{CO}_2\text{R}^{10}$, wherein R^9 and R^{10} are as
defined above,
10 (l) $-\text{CONR}^9\text{R}^{10}$, wherein R^9 and R^{10} are as
defined above,
(m) $-\text{CO}_2\text{NR}^9\text{R}^{10}$, wherein R^9 and R^{10} are as
defined above,
(n) $-\text{COR}^9$, wherein R^9 is as defined above;
(o) $-\text{CO}_2\text{R}^9$, wherein R^9 is as defined above;
15 (6) halo,
(7) $-\text{CN}$,
(8) $-\text{CF}_3$,
(9) $-\text{NO}_2$,
(10) $-\text{SR}^{14}$, wherein R^{14} is hydrogen or C_1 -5alkyl,
(11) $-\text{SOR}^{14}$, wherein R^{14} is as defined above,
20 (12) $-\text{SO}_2\text{R}^{14}$, wherein R^{14} is as defined above,
(13) $\text{NR}^9\text{COR}^{10}$, wherein R^9 and R^{10} are as defined above,
(14) $\text{CONR}^9\text{COR}^{10}$, wherein R^9 and R^{10} are as defined above,
(15) NR^9R^{10} , wherein R^9 and R^{10} are as defined above,
(16) $\text{NR}^9\text{CO}_2\text{R}^{10}$, wherein R^9 and R^{10} are as defined above,
25 (17) hydroxy,
(18) C_1 -6alkoxy,
(19) COR^9 , wherein R^9 is as defined above,
(20) CO_2R^9 , wherein R^9 is as defined above,
30 (21) 2-pyridyl,
(22) 3-pyridyl,
(23) 4-pyridyl,
(24) 5-tetrazolyl,
(25) 2-oxazolyl, and
(26) 2-thiazolyl;

- 20 -

R11, R12 and R13 are independently selected from the definitions of R6, R7 and R8;

5 X is selected from the group consisting of:

- (1) -O-,
- (2) -S-,
- (3) -SO-, and
- (4) -SO₂-;

10

Y is selected from the group consisting of:

- (1) a single bond,
- (2) -O-,
- (3) -S-,
- (4) -CO-,
- (5) -CH₂-,
- (6) -CHR₁₅-, and
- (7) -CR₁₅R₁₆-, wherein R₁₅ and R₁₆ are independently selected from the group consisting of:

20

(a) C₁₋₆ alkyl, unsubstituted or substituted with one or more of the substituents selected from:

- (i) hydroxy,
- (ii) oxo,
- (iii) C₁₋₆ alkoxy,
- (iv) phenyl-C₁₋₃ alkoxy,
- (v) phenyl,
- (vi) -CN,
- (vii) halo,
- (viii) -NR⁹R₁₀, wherein R⁹ and R₁₀ are as defined above,
- (ix) -NR⁹COR₁₀, wherein R⁹ and R₁₀ are as defined above,
- (x) -NR⁹CO₂R₁₀, wherein R⁹ and R₁₀ are as defined above,

25

30

- 21 -

- (xi) $-\text{CONR}^9\text{R}^{10}$, wherein R^9 and R^{10} are as defined above,
(xii) $-\text{COR}^9$, wherein R^9 is as defined above, and
(xiii) $-\text{CO}_2\text{R}^9$, wherein R^9 is as defined above;
5 (b) phenyl, unsubstituted or substituted with one or more of the substituent(s) selected from:
(i) hydroxy,
(ii) C1-6 alkoxy,
(iii) C1-6 alkyl,
10 (iv) C2-5 alkenyl,
(v) halo,
(vi) $-\text{CN}$,
(vii) $-\text{NO}_2$,
(viii) $-\text{CF}_3$,
15 (ix) $-(\text{CH}_2)_m-\text{NR}^9\text{R}^{10}$, wherein m , R^9 and R^{10} are as defined above,
(x) $-\text{NR}^9\text{COR}^{10}$, wherein R^9 and R^{10} are as defined above,
(xi) $-\text{NR}^9\text{CO}_2\text{R}^{10}$, wherein R^9 and R^{10} are as defined above,
20 (xii) $-\text{CONR}^9\text{R}^{10}$, wherein R^9 and R^{10} are as defined above,
(xiii) $-\text{CO}_2\text{NR}^9\text{R}^{10}$, wherein R^9 and R^{10} are as defined above,
25 (xiv) $-\text{COR}^9$, wherein R^9 is as defined above, and
(xv) $-\text{CO}_2\text{R}^9$, wherein R^9 is as defined above;

Z is selected from:

- 30 (1) hydrogen,
(2) C1-6 alkyl, and
(3) hydroxy, with the proviso that if Y is $-\text{O}-$, Z is other than hydroxy, or if Y is $-\text{CHR}^{15}-$, then Z and R^{15} may be joined together to form a double bond.

- 22 -

The compounds of the present invention have asymmetric centers and this invention includes all of the optical isomers and mixtures thereof.

5 In addition compounds with carbon-carbon double bonds may occur in Z- and E- forms with all isomeric forms of the compounds being included in the present invention.

When any variable (e.g., alkyl, aryl, R⁶, R⁷, R⁸, R⁹, R¹⁰, R¹¹, R¹², R¹³, etc.) occurs more than one time in any variable or in Formula I, its definition on each occurrence is independent of its
10 definition at every other occurrence.

As used herein, the term "alkyl" includes those alkyl groups of a designated number of carbon atoms of either a straight, branched, or cyclic configuration. Examples of "alkyl" include methyl, ethyl, propyl, isopropyl, butyl, iso- sec- and tert-butyl, pentyl, hexyl, heptyl,
15 3-ethylbutyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, norbornyl, and the like. "Alkoxy" represents an alkyl group of indicated number of carbon atoms attached through an oxygen bridge, such as methoxy, ethoxy, propoxy, butoxy and pentoxy. "Alkenyl" is intended to include hydrocarbon chains of a specified
20 number of carbon atoms of either a straight- or branched- configuration and at least one unsaturation, which may occur at any point along the chain, such as ethenyl, propenyl, butenyl, pentenyl, dimethylpentenyl, and the like, and includes E and Z forms, where applicable. "Halogen" or "halo", as used herein, means fluoro, chloro, bromo and iodo.

25 The term "aryl" means phenyl or naphthyl either unsubstituted or substituted with one, two or three substituents selected from the group consisting of halo, C₁₋₄-alkyl, C₁₋₄-alkoxy, NO₂, CF₃, C₁₋₄-alkylthio, OH, -N(R⁶)₂, -CO₂R⁶, C₁₋₄-perfluoroalkyl, C₃₋₆-perfluorocycloalkyl, and tetrazol-5-yl.


30 The term "heteroaryl" means an unsubstituted, monosubstituted or disubstituted five or six membered aromatic heterocycle comprising from 1 to 3 heteroatoms selected from the group consisting of O, N and S and wherein the substituents are members selected from the group consisting of -OH, -SH, -C₁₋₄-alkyl,

- 23 -

-C₁-4-alkoxy, -CF₃, halo, -NO₂, -CO₂R⁹, -N(R⁹R¹⁰) and a fused benzo group.

As will be understood by those skilled in the art, pharmaceutically acceptable salts include, but are not limited to salts with inorganic acids such as hydrochloride, sulfate, phosphate, diphosphate, hydrobromide, and nitrate or salts with an organic acid such as malate, maleate, fumarate, tartrate, succinate, citrate, acetate, lactate, methanesulfonate, p-toluenesulfonate, 2-hydroxyethylsulfonate, pamoate, salicylate and stearate. Similarly, pharmaceutically acceptable cations include, but are not limited to sodium, potassium, calcium, aluminum, lithium and ammonium.

A preferred embodiment of the present invention is directed to compounds of the formula:



or a pharmaceutically acceptable salt thereof, wherein:

R¹ is selected from the group consisting of:

- (1) hydrogen;
- (2) C₁-6 alkyl, unsubstituted or substituted with one or more of the substituents selected from:
 - (a) hydroxy,
 - (b) oxo,
 - (c) C₁-6 alkoxy,
 - (d) phenyl-C₁-3 alkoxy,
 - (e) phenyl,
 - (f) -CN,
 - (g) halo, wherein halo is fluoro, chloro, bromo or iodo,

- 24 -

- 5
- (h) -NR⁹R¹⁰, wherein R⁹ and R¹⁰ are independently selected from:
- (i) hydrogen,
 - (ii) C₁₋₆ alkyl,
 - (iii) hydroxy-C₁₋₆ alkyl, and
 - (iv) phenyl,
- 10
- (i) -NR⁹COR¹⁰, wherein R⁹ and R¹⁰ are as defined above,
- (j) -NR⁹CO₂R¹⁰, wherein R⁹ and R¹⁰ are as defined above,
- (k) -CONR⁹R¹⁰, wherein R⁹ and R¹⁰ are as defined above,
- (l) -COR⁹, wherein R⁹ is as defined above,
- 15
- (m) -CO₂R⁹, wherein R⁹ is as defined above;
- (n) heterocycle, wherein the heterocycle is selected from the group consisting of:
- (A) benzimidazolyl,
 - (B) benzofuranyl,
 - (C) benzothiophenyl,
 - 20
 - (D) benzoxazolyl,
 - (E) furanyl,
 - (F) imidazolyl,
 - (G) indolyl,
 - (H) isooxazolyl,
 - 25
 - (I) isothiazolyl,
 - (J) oxadiazolyl,
 - (K) oxazolyl,
 - (L) pyrazinyl,
 - (M) pyrazolyl,
 - 30
 - (N) pyridyl,
 - (O) pyrimidyl,
 - (P) pyrrolyl,
 - (Q) quinolyl,
 - (R) tetrazolyl,

- 25 -

- (S) thiadiazolyl,
(T) thiazolyl,
(U) thienyl,
(V) triazolyl,
(W) azetidinyI,
(X) 1,4-dioxanyl,
(Y) hexahydroazepinyl,
(Z) piperazinyl,
(AA) piperidinyl,
(AB) pyrrolidinyl,
(AC) tetrahydrofuranyl, and
(AD) tetrahydrothienyl,
and wherein the heterocycle is unsubstituted or substituted with one or more substituent(s) selected from:
- (i) C₁₋₆ alkyl, unsubstituted or substituted with halo, -CF₃, -OCH₃, or phenyl,
(ii) C₁₋₆ alkoxy,
(iii) oxo,
(iv) hydroxy,
(v) thioxo,
(vi) -SR⁹, wherein R⁹ is as defined above,
(vii) halo,
(viii) cyano,
(ix) phenyl,
(x) trifluoromethyl,
(xi) -(CH₂)_m-NR⁹R¹⁰, wherein m is 0, 1 or 2, and R⁹ and R¹⁰ are as defined above,
(xii) -NR⁹COR¹⁰, wherein R⁹ and R¹⁰ are as defined above,
(xiii) -CONR⁹R¹⁰, wherein R⁹ and R¹⁰ are as defined above,
(xiv) -CO₂R⁹, wherein R⁹ is as defined above, and

- 26 -

(xv) $-(\text{CH}_2)_m\text{-OR}^9$, wherein m and R^9 are as defined above;

- 5 (3) C₂-6 alkenyl, unsubstituted or substituted with one or more of the substituent(s) selected from:
- (a) hydroxy,
 - (b) oxo,
 - (c) C₁-6 alkoxy,
 - 10 (d) phenyl-C₁-3 alkoxy,
 - (e) phenyl,
 - (f) -CN,
 - (g) halo,
 - (h) $-\text{CONR}^9\text{R}^{10}$ wherein R^9 and R^{10} are as defined above,
 - 15 (i) $-\text{COR}^9$ wherein R^9 is as defined above,
 - (j) $-\text{CO}_2\text{R}^9$, wherein R^9 is as defined above,
 - (k) heterocycle, wherein the heterocycle is as defined above;
- 20 (4) C₂-6 alkynyl;
- (5) phenyl, unsubstituted or substituted with one or more of the substituent(s) selected from:
- (a) hydroxy,
 - (b) C₁-6 alkoxy,
 - (c) C₁-6 alkyl,
 - 25 (d) C₂-5 alkenyl,
 - (e) halo,
 - (f) -CN,
 - (g) -NO₂,
 - (h) -CF₃,
 - 30 (i) $-(\text{CH}_2)_m\text{-NR}^9\text{R}^{10}$, wherein m , R^9 and R^{10} are as defined above,
 - (j) $-\text{NR}^9\text{COR}^{10}$, wherein R^9 and R^{10} are as defined above,

- 27 -

- (k) $-\text{NR}^9\text{CO}_2\text{R}^{10}$, wherein R^9 and R^{10} are as defined above,
- (l) $-\text{CONR}^9\text{R}^{10}$, wherein R^9 and R^{10} are as defined above,
- 5 (m) $-\text{CO}_2\text{NR}^9\text{R}^{10}$, wherein R^9 and R^{10} are as defined above,
- (n) $-\text{COR}^9$, wherein R^9 is as defined above;
- (o) $-\text{CO}_2\text{R}^9$, wherein R^9 is as defined above;
- 10 R^2 and R^3 are independently selected from the group consisting of:
- (1) hydrogen,
- (2) C1-6 alkyl, unsubstituted or substituted with one or more of the substituents selected from:
- 15 (a) hydroxy,
- (b) oxo,
- (c) C1-6 alkoxy,
- (d) phenyl-C1-3 alkoxy,
- (e) phenyl,
- 20 (f) $-\text{CN}$,
- (g) halo,
- (h) $-\text{NR}^9\text{R}^{10}$, wherein R^9 and R^{10} are as defined above,
- (i) $-\text{NR}^9\text{COR}^{10}$, wherein R^9 and R^{10} are as defined above,
- 25 (j) $-\text{NR}^9\text{CO}_2\text{R}^{10}$, wherein R^9 and R^{10} are as defined above,
- (k) $-\text{CONR}^9\text{R}^{10}$, wherein R^9 and R^{10} are as defined above,
- (l) $-\text{COR}^9$, wherein R^9 is as defined above, and
- 30 (m) $-\text{CO}_2\text{R}^9$, wherein R^9 is as defined above;
- (3) C2-6 alkenyl, unsubstituted or substituted with one or more of the substituent(s) selected from:
- (a) hydroxy,
- (b) oxo,

- 28 -

- 5
- (c) C₁₋₆ alkoxy,
 - (d) phenyl-C₁₋₃ alkoxy,
 - (e) phenyl,
 - (f) -CN,
 - (g) halo,
 - (h) -CONR⁹R¹⁰ wherein R⁹ and R¹⁰ are as defined above,
 - (i) -COR⁹ wherein R⁹ is as defined above,
 - (j) -CO₂R⁹, wherein R⁹ is as defined above;
- 10
- (4) C₂₋₆ alkynyl;
 - (5) phenyl, unsubstituted or substituted with one or more of the substituent(s) selected from:
- 15
- (a) hydroxy,
 - (b) C₁₋₆ alkoxy,
 - (c) C₁₋₆ alkyl,
 - (d) C₂₋₅ alkenyl,
 - (e) halo,
 - (f) -CN,
 - (g) -NO₂,
 - (h) -CF₃,
 - (i) -(CH₂)_m-NR⁹R¹⁰, wherein m, R⁹ and R¹⁰ are as defined above,
 - (j) -NR⁹COR¹⁰, wherein R⁹ and R¹⁰ are as defined above,
 - (k) -NR⁹CO₂R¹⁰, wherein R⁹ and R¹⁰ are as defined above,
 - (l) -CONR⁹R¹⁰, wherein R⁹ and R¹⁰ are as defined above,
 - (m) -CO₂NR⁹R¹⁰, wherein R⁹ and R¹⁰ are as defined above,
 - (n) -COR⁹, wherein R⁹ is as defined above;
 - (o) -CO₂R⁹, wherein R⁹ is as defined above;
- 20
- 25
- 30

- 29 -

and the groups R¹ and R² may be joined together to form a heterocyclic ring selected from the group consisting of:

- (a) pyrrolidinyl,
- (b) piperidinyl,
- (c) pyrrolyl,
- (d) pyridinyl,
- (e) imidazolyl,
- (f) oxazolyl, and
- (g) thiazolyl,

and wherein the heterocyclic ring is unsubstituted or substituted with one or more substituent(s) selected from:

- (i) C₁-6alkyl,
- (ii) oxo,
- (iii) C₁-6alkoxy,
- (iv) -NR⁹R¹⁰, wherein R⁹ and R¹⁰ are as defined above,
- (v) halo, and
- (vi) trifluoromethyl;

and the groups R² and R³ may be joined together to form a carbocyclic ring selected from the group consisting of:

- (a) cyclopentyl,
- (b) cyclohexyl,
- (c) phenyl,

and wherein the carbocyclic ring is unsubstituted or substituted with one or more substituents selected from:

- (i) C₁-6alkyl,
- (ii) C₁-6alkoxy,
- (iii) -NR⁹R¹⁰, wherein R⁹ and R¹⁰ are as defined above,
- (iv) halo, and
- (v) trifluoromethyl;

- 30 -

and the groups R² and R³ may be joined together to form a heterocyclic ring selected from the group consisting of:

- (a) pyrrolidinyl,
- (b) piperidinyl,
- (c) pyrrolyl,
- (d) pyridinyl,
- (e) imidazolyl,
- (f) furanyl,
- (g) oxazolyl,
- (h) thienyl, and
- (i) thiazolyl,

and wherein the heterocyclic ring is unsubstituted or substituted with one or more substituent(s) selected from:

- (i) C₁₋₆alkyl,
- (ii) oxo,
- (iii) C₁₋₆alkoxy,
- (iv) -NR⁹R¹⁰, wherein R⁹ and R¹⁰ are as defined above,
- (v) halo, and
- (vi) trifluoromethyl;

R⁶, R⁷ and R⁸ are independently selected from the group consisting of:

- (1) hydrogen;
- (2) C₁₋₆ alkyl, unsubstituted or substituted with one or more of the substituents selected from:
 - (a) hydroxy,
 - (b) oxo,
 - (c) C₁₋₆ alkoxy,
 - (d) phenyl-C₁₋₃ alkoxy,
 - (e) phenyl,
 - (f) -CN,
 - (g) halo,
 - (h) -NR⁹R¹⁰, wherein R⁹ and R¹⁰ are as defined above,

- 31 -

- 5
- (i) $-\text{NR}^9\text{COR}^{10}$, wherein R^9 and R^{10} are as defined above,
- (j) $-\text{NR}^9\text{CO}_2\text{R}^{10}$, wherein R^9 and R^{10} are as defined above,
- (k) $-\text{CONR}^9\text{R}^{10}$, wherein R^9 and R^{10} are as defined above,
- (l) $-\text{COR}^9$, wherein R^9 is as defined above, and
- (m) $-\text{CO}_2\text{R}^9$, wherein R^9 is as defined above;
- 10 (3) C₂-6 alkenyl, unsubstituted or substituted with one or more of the substituent(s) selected from:
- (a) hydroxy,
- (b) oxo,
- (c) C₁-6 alkoxy,
- 15 (d) phenyl-C₁-3 alkoxy,
- (e) phenyl,
- (f) -CN,
- (g) halo,
- (h) $-\text{CONR}^9\text{R}^{10}$ wherein R^9 and R^{10} are as defined above,
- 20 (i) $-\text{COR}^9$ wherein R^9 is as defined above,
- (j) $-\text{CO}_2\text{R}^9$, wherein R^9 is as defined above;
- (4) C₂-6 alkynyl;
- (5) phenyl, unsubstituted or substituted with one or more of the substituent(s) selected from:
- 25 (a) hydroxy,
- (b) C₁-6 alkoxy,
- (c) C₁-6 alkyl,
- (d) C₂-5 alkenyl,
- 30 (e) halo,
- (f) -CN,
- (g) -NO₂,
- (h) -CF₃,
- (i) $-(\text{CH}_2)_m-\text{NR}^9\text{R}^{10}$, wherein m , R^9 and R^{10} are as defined above,

- 32 -

- (j) $-\text{NR}^9\text{COR}^{10}$, wherein R^9 and R^{10} are as defined above,
- (k) $-\text{NR}^9\text{CO}_2\text{R}^{10}$, wherein R^9 and R^{10} are as defined above,
- 5 (l) $-\text{CONR}^9\text{R}^{10}$, wherein R^9 and R^{10} are as defined above,
- (m) $-\text{CO}_2\text{NR}^9\text{R}^{10}$, wherein R^9 and R^{10} are as defined above,
- 10 (n) $-\text{COR}^9$, wherein R^9 is as defined above;
- (o) $-\text{CO}_2\text{R}^9$, wherein R^9 is as defined above;
- (6) halo,
- (7) $-\text{CN}$,
- (8) $-\text{CF}_3$,
- (9) $-\text{NO}_2$,
- 15 (10) $-\text{SR}^{14}$, wherein R^{14} is hydrogen or C_{1-5} alkyl,
- (11) $-\text{SOR}^{14}$, wherein R^{14} is as defined above,
- (12) $-\text{SO}_2\text{R}^{14}$, wherein R^{14} is as defined above,
- (13) $\text{NR}^9\text{COR}^{10}$, wherein R^9 and R^{10} are as defined above,
- 20 (14) $\text{CONR}^9\text{COR}^{10}$, wherein R^9 and R^{10} are as defined above,
- (15) NR^9R^{10} , wherein R^9 and R^{10} are as defined above,
- (16) $\text{NR}^9\text{CO}_2\text{R}^{10}$, wherein R^9 and R^{10} are as defined above,
- (17) hydroxy,
- (18) C_{1-6} alkoxy,
- 25 (19) COR^9 , wherein R^9 is as defined above,
- (20) CO_2R^9 , wherein R^9 is as defined above,
- (21) 2-pyridyl,
- (22) 3-pyridyl,
- (23) 4-pyridyl,
- 30 (24) 5-tetrazolyl,
- (25) 2-oxazolyl, and
- (26) 2-thiazolyl;

R^{11} , R^{12} and R^{13} are independently selected from the definitions of R^6 , R^7 and R^8 ;

- 33 -

X is selected from the group consisting of:

- 5
- (1) -O-,
 - (2) -S-,
 - (3) -SO-, and
 - (4) -SO₂-;

Y is selected from the group consisting of:

- 10
- (1) a single bond,
 - (2) -O-,
 - (3) -S-,
 - (4) -CO-,
 - (5) -CH₂-,
 - (6) -CHR¹⁵-, and
 - 15 (7) -CR¹⁵R¹⁶-, wherein R¹⁵ and R¹⁶ are independently selected from the group consisting of:
 - (a) C₁₋₆ alkyl, unsubstituted or substituted with one or more of the substituents selected from:

20

 - (i) hydroxy,
 - (ii) oxo,
 - (iii) C₁₋₆ alkoxy,
 - (iv) phenyl-C₁₋₃ alkoxy,
 - (v) phenyl,
 - 25 (vi) -CN,
 - (vii) halo,
 - (viii) -NR⁹R¹⁰, wherein R⁹ and R¹⁰ are as defined above,
 - (ix) -NR⁹COR¹⁰, wherein R⁹ and R¹⁰ are as defined above,
 - 30 (x) -NR⁹CO₂R¹⁰, wherein R⁹ and R¹⁰ are as defined above,
 - (xi) -CONR⁹R¹⁰, wherein R⁹ and R¹⁰ are as defined above,
 - (xii) -COR⁹, wherein R⁹ is as defined above, and

- 34 -

(xiii) $-\text{CO}_2\text{R}^9$, wherein R^9 is as defined above;

(b) phenyl, unsubstituted or substituted with one or more of the substituent(s) selected from:

5

(i) hydroxy,

(ii) C₁₋₆ alkoxy,

(iii) C₁₋₆ alkyl,

(iv) C₂₋₅ alkenyl,

10

(v) halo,

(vi) $-\text{CN}$,

(vii) $-\text{NO}_2$,

(viii) $-\text{CF}_3$,

15

(ix) $-(\text{CH}_2)_m-\text{NR}^9\text{R}^{10}$, wherein m, R^9 and R^{10} are as defined above,

(x) $-\text{NR}^9\text{COR}^{10}$, wherein R^9 and R^{10} are as defined above,

(xi) $-\text{NR}^9\text{CO}_2\text{R}^{10}$, wherein R^9 and R^{10} are as defined above,

20

(xii) $-\text{CONR}^9\text{R}^{10}$, wherein R^9 and R^{10} are as defined above,

(xiii) $-\text{CO}_2\text{NR}^9\text{R}^{10}$, wherein R^9 and R^{10} are as defined above,

(xiv) $-\text{COR}^9$, wherein R^9 is as defined above, and

25

(xv) $-\text{CO}_2\text{R}^9$, wherein R^9 is as defined above;

Z is C₁₋₆ alkyl.

In the preferred embodiment of the present compounds it is more preferred that:

30

R^1 is C₁₋₆ alkyl, substituted with one or more of the substituents selected from:

heterocycle, wherein the heterocycle is selected from the group consisting of:

(A) benzimidazolyl,

- 35 -

- 5 (B) imidazolyl,
(C) isooxazolyl,
(D) isothiazolyl,
(E) oxadiazolyl,
(F) pyrazinyl,
(G) pyrazolyl,
(H) pyridyl,
(I) pyrrolyl,
10 (J) tetrazolyl,
(K) thiadiazolyl,
(L) triazolyl, and
(M) piperidinyl,
and wherein the heterocycle is unsubstituted or
substituted with one or more substituent(s) selected
15 from:

- (i) C₁₋₆ alkyl, unsubstituted or
substituted with halo, -CF₃,
-OCH₃, or phenyl,
20 (ii) C₁₋₆ alkoxy,
(iii) oxo,
(iv) thioxo,
(v) cyano,
(vi) -SCH₃,
25 (vii) phenyl,
(viii) hydroxy,
(ix) trifluoromethyl,
(x) -(CH₂)_m-NR⁹R¹⁰, wherein m is
0, 1 or 2, and wherein R⁹ and
30 R¹⁰ are independently selected
from:
(I) hydrogen,
(II) C₁₋₆ alkyl,
(III) hydroxy-C₁₋₆ alkyl, and
(IV) phenyl,

- 36 -

(xi) $-NR^9COR^{10}$, wherein R^9 and R^{10}
are as defined above, and

(xii) $-CONR^9R^{10}$, wherein R^9 and R^{10}
are as defined above.

5

In the preferred embodiment of present compounds it is
also more preferred that:

R^2 and R^3 are independently selected from the group consisting of:

- 10
- (1) hydrogen,
 - (2) C1-6 alkyl,
 - (3) C2-6 alkenyl, and
 - (4) phenyl;

R^6 , R^7 and R^8 are independently selected from the group consisting of:

- 15
- (1) hydrogen,
 - (2) C1-6 alkyl,
 - (3) fluoro,
 - (4) chloro,
 - (5) bromo,
 - (6) iodo, and
 - 20 (7) $-CF_3$;

R^{11} , R^{12} and R^{13} are independently selected from the group consisting
of:

- 25
- (1) hydrogen,
 - (2) C1-6 alkyl,
 - (3) fluoro,
 - (4) chloro,
 - (5) bromo,
 - (6) iodo, and
 - 30 (7) $-CF_3$;

X is $-O-$;


Y is $-O-$; and

Z is C1-4 alkyl.

- 37 -


In the compounds of the present invention a preferred embodiment includes those compounds wherein Z is C₁₋₄ alkyl. An especially preferred embodiment of the present compounds includes those compounds wherein Z is -CH₃. These compounds bearing a substituent on the alpha-carbon atom exhibit advantageous pharmacological properties, in particular, enhanced duration of action in models of extravasation, presumably due to biological stability and resistance to enzymatic degradation.

An embodiment of the novel compounds of this invention is that wherein X is O, R⁴ is -YCHZ-phenyl, and R⁵ is phenyl of structural formula:



or a pharmaceutically acceptable salt thereof, wherein R¹, R², R³, R⁶, R⁷, R⁸, R¹¹, R¹², R¹³, Y and Z are as defined above.


Within this embodiment, a preferred class of compounds includes those compounds of the structural formula:



or a pharmaceutically acceptable salt thereof, wherein R¹, R², R³, R⁶, R⁷, R⁸, R¹¹, R¹², R¹³, Y and Z are as defined in Claim 1.


- 38 -

An even more preferred class of compounds within this embodiment includes those compounds of the structural formula III:



or a pharmaceutically acceptable salt thereof, wherein R¹, R², R³, R⁶, R⁷, R⁸, R¹¹, R¹², R¹³ and Z are as defined in Claim 1.

15 An alternate class of compounds within this embodiment includes those compounds of the structural formula II:




or a pharmaceutically acceptable salt thereof, wherein R¹, R², R³, R⁶, R⁷, R⁸, R¹¹, R¹², R¹³ and Z are as defined in Claim 1.

30


- 39 -

Another embodiment of the novel compounds of this invention is that wherein X is S, R⁴ is -Y-CHZ-phenyl, and R⁵ is phenyl of structural formula:



or a pharmaceutically acceptable salt thereof, wherein R¹, R², R³, R⁶, R⁷, R⁸, R¹¹, R¹², R¹³, Y and Z are as defined above.

15 Another embodiment of the novel compounds of this invention is that wherein X is SO, R⁴ is -Y-CHZ-phenyl, and R⁵ is phenyl of structural formula:



or a pharmaceutically acceptable salt thereof, wherein R¹, R², R³, R⁶, R⁷, R⁸, R¹¹, R¹², R¹³, Y and Z are as defined above.


30

- 40 -

Another embodiment of the novel compounds of this invention is that wherein X is SO₂, R⁴ is -Y-CHZ-phenyl, and R⁵ is phenyl of structural formula:

5

10



or a pharmaceutically acceptable salt thereof, wherein R¹, R², R³, R⁶, R⁷, R⁸, R¹¹, R¹², R¹³, Y and Z are as defined above.


15

In the compounds of the present invention a preferred embodiment is that in which R¹ is selected from the following group of substituents:

20




25

30





- 41 -



5





10





15








20




25



30



- 42 -



A particularly preferred embodiment of the compounds of the present invention includes those compounds wherein R^1 is (1,2,4-triazolo)methyl or (5-oxo-1H,4H-1,2,4-triazolo)methyl.

Another particularly preferred embodiment of the compounds of the present invention includes those compounds wherein R^1 is (1,3-imidazolo)methyl or (2-oxo-1,3-imidazolo)methyl.

Specific preferred compounds within the present invention include:

- 1) (+/-)-2-(3,5-bis(trifluoromethyl)benzyloxy)-3-phenyl-morpholine;
- 2) (2R,S)-(3,5-bis(trifluoromethyl)benzyloxy)-(3R)-phenyl-(6R)-methyl-morpholine;

- 43 -

- 3) (2R,S)-(3,5-bis(trifluoromethyl)benzyloxy)-(3S)-phenyl-(6R)-methyl-morpholine;
- 5 4) (+/-)-2-(3,5-bis(trifluoromethyl)benzyloxy)-3-phenyl-4-methylcarboxamido-morpholine;
- 5) (+/-)-2-(3,5-bis(trifluoromethyl)benzyloxy)-3-phenyl-4-methoxy-carbonylmethyl-morpholine;
- 10 6) 2-(2-(3,5-bis(trifluoromethyl)phenyl)ethenyl)-3-phenyl-5-oxo-morpholine;
- 7) 3-phenyl-2-(2-(3,5-bis(trifluoromethyl)phenyl)-ethyl)-morpholine;
- 15 8) 2-(R)-(3,5-bis(trifluoromethyl)benzyloxy)-3-(R)-phenyl-6-(S)-methyl-morpholine;
- 9) 2-(R)-(3,5-bis(trifluoromethyl)benzyloxy)-3-(S)-phenyl-6-(S)-methyl-morpholine;
- 20 10) 2-(S)-(3,5-bis(trifluoromethyl)benzyloxy)-3-(R)-phenyl-6-(S)-methyl-morpholine;
- 25 11) 2-(S)-(3,5-bis(trifluoromethyl)benzyloxy)-3-(S)-phenyl-6-(S)-methyl-morpholine;
- 12) 2-(R)-(3,5-bis(trifluoromethyl)benzyloxy)-3-(R)-phenyl-5-(R)-methyl-morpholine;
- 30 13) 2-(R)-(3,5-bis(trifluoromethyl)benzyloxy)-3-(S)-phenyl-5-(R)-methyl-morpholine;
- 14) 2-(S)-(3,5-bis(trifluoromethyl)benzyloxy)-3-(R)-phenyl-5-(R)-

- 44 -

methyl-morpholine;

- 5 15) 2-(S)-(3,5-bis(trifluoromethyl)benzyloxy)-3-(S)-phenyl-5-(R)-methyl-morpholine;
- 16) 2-(S)-(3,5-bis(trifluoromethyl)benzyloxy)-3-(S)-phenyl-morpholine;
- 10 17) 4-(3-(1,2,4-triazolo)methyl)-2-(S)-(3,5-bis(trifluoromethyl)benzyloxy)-3-(S)-phenyl-morpholine;
- 18) 4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-2-(S)-(3,5-bis(trifluoromethyl)benzyloxy)-3-(S)-phenyl-morpholine;
- 15 19) 2-(R)-(3,5-bis(trifluoromethyl)benzyloxy)-3-(R)-phenyl-6-(R)-methyl-morpholine;
- 20 20) 2-(S)-(3,5-bis(trifluoromethyl)benzyloxy)-3-(R)-phenyl-6-(R)-methyl-morpholine;
- 21) 2-(R)-(3,5-bis(trifluoromethyl)benzyloxy)-3-(S)-phenyl-6-(R)-methyl-morpholine;
- 25 22) 2-(S)-(3,5-bis(trifluoromethyl)benzyloxy)-3-(S)-phenyl-6-(R)-methyl-morpholine;
- 23) 2-(R)-(3,5-bis(trifluoromethyl)-benzyloxy)-3-(S)-phenyl-5-(S)-methyl-morpholine;
- 30 24) 2-(S)-(3,5-bis(trifluoromethyl)benzyloxy)-3-(S)-phenyl-5-(S)-methyl-morpholine;
- 25) 2-(R)-(3,5-bis(trifluoromethyl)benzyloxy)-3-(R)-phenyl-5-(S)-methyl-morpholine;

- 45 -

- 26) 2-(R)-(3,5-bis(trifluoromethyl)benzyloxy)-3-(S)-phenyl-5-(R)-phenyl-morpholine;
- 5 27) 2-(S)-(3,5-bis(trifluoromethyl)benzyloxy)-3-(S)-phenyl-5-(R)-phenyl-morpholine;
- 28) 2-(S)-(3,5-bis(trifluoromethyl)benzyloxy)-3-(R)-phenyl-5-(S)-phenyl-morpholine;
- 10 29) 2-(R)-(3,5-bis(trifluoromethyl)benzyloxy)-3-(R)-phenyl-5-(S)-phenyl-morpholine;
- 30) 2-(S)-(3,5-bis(trifluoromethyl)benzyloxy)-6-(R)-methyl-3-(S)-phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 15 31) 2-(S)-(3,5-bis(trifluoromethyl)benzyloxy)-6-(R)-methyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-3-(S)-phenyl-morpholine;
- 20 32) 2-(S)-(3,5-bis(trifluoromethyl)benzyloxy)-3-(R)-phenyl-morpholine;
- 33) 4-(3-(1,2,4-triazolo)methyl)-2-(S)-(3,5-bis(trifluoromethyl)benzyloxy)-3-(R)-phenyl-morpholine;
- 25 34) 4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-2-(S)-(3,5-bis(trifluoromethyl)benzyloxy)-3-(R)-phenyl-morpholine;
- 35) 4-(2-(imidazolo)methyl)-2-(S)-(3,5-bis(trifluoromethyl)benzyloxy)-3-(R)-phenyl-morpholine;
- 30 36) 4-(4-(imidazolo)methyl)-2-(S)-(3,5-bis(trifluoromethyl)benzyloxy)-3-(R)-phenyl-morpholine;

- 46 -

- 37) 4-(aminocarbonylmethyl)-2-(S)-(3,5-bis(trifluoromethyl)-benzyloxy)-3-(R)-phenyl-morpholine;
- 5 38) 4-(2-(imidazolo)methyl)-2-(S)-(3,5-bis(trifluoromethyl)-benzyloxy)-3-(S)-phenyl-morpholine;
- 39) 4-(4-(imidazolo)methyl)-2-(S)-(3,5-bis(trifluoromethyl)-benzyloxy)-3-(S)-phenyl-morpholine;
- 10 40) 4-(2-(imidazolo)methyl)-2-(S)-(3,5-bis(trifluoromethyl)-benzyloxy)-3-(S)-phenyl-6-(R)-methyl-morpholine;
- 41) 4-(4-(imidazolo)methyl)-2-(S)-(3,5-bis(trifluoromethyl)-benzyloxy)-3-(S)-phenyl-6(R)-methyl-morpholine;
- 15 42) 2-(S)-(3,5-bis(trifluoromethyl)benzyloxy)-4-((6-hydroxy)-hexyl)-3-(R)-phenyl-morpholine;
- 20 43) 2-(S)-(3,5-bis(trifluoromethyl)benzyloxy)-4-(5-(methylamino-carbonyl)pentyl)-3-(R)-phenyl-morpholine;
- 44) 4-(3-(1,2,4-triazolo)methyl)-2-(3,5-dimethylbenzyloxy)-3-phenylmorpholine;
- 25 45) 4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-2-(3,5-dimethyl)-benzyloxy)-3-phenyl-morpholine;
- 46) 4-(3-(1,2,4-triazolo)methyl)-2-(3,5-di(tert-butyl)-benzyloxy)-3-phenylmorpholine;
- 30 47) 4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-2-(3,5-di(tert-butyl)-benzyloxy)-3-phenyl-morpholine;
- 48) 4-(3-(1,2,4-triazolo)methyl)-2-(3-(tert-butyl)-5-methyl-

- 47 -

benzyloxy)-3-phenyl-morpholine;

- 5 49) 4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-2-(3-(tert-butyl)-5-methylbenzyloxy)-3-phenyl-morpholine;
- 50) 4-(3-(1,2,4-triazolo)methyl)-2-(3-(trifluoro-methyl)-5-methyl-benzyloxy)-3-phenyl-morpholine;
- 10 51) 4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-2-(3-(trifluoro-methyl)-5-methylbenzyloxy)-3-phenyl-morpholine;
- 52) 4-(3-(1,2,4-triazolo)methyl)-2-(3-(tert-butyl)-5-(trifluoro-methyl)benzyloxy)-3-phenyl-morpholine;
- 15 53) 4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-2-(3-(tert-butyl)-5-(trifluoromethyl)benzyloxy)-3-phenyl-morpholine;
- 54) 4-(2-(imidazolo)methyl)-2-(3,5-dimethyl-benzyloxy)-3-phenyl-morpholine;
- 20 55) 4-(4-(imidazolo)methyl)-2-(3,5-dimethyl-benzyloxy)-3-phenyl-morpholine;
- 56) 4-(2-(imidazolo)methyl)-2-(3,5-di(tert-butyl)-benzyloxy)-3-phenyl-morpholine;
- 25 57) 4-(4-(imidazolo)methyl)-2-(3,5-di(tert-butyl)-benzyloxy)-3-phenyl-morpholine;
- 30 58) 4-(2-(imidazolo)methyl)-2-(3-(tert-butyl)-5-methylbenzyloxy)-3-phenyl-morpholine;
- 59) 4-(4-(imidazolo)methyl)-2-(3-(tert-butyl)-5-methylbenzyloxy)-3-phenyl-morpholine;

- 48 -

- 60) 4-(2-(imidazolo)methyl)-2-(3-(trifluoro-methyl)-5-methyl-benzyloxy)-3-phenyl-morpholine;
- 5 61) 4-(4-(imidazolo)methyl)-2-(3-(trifluoro-methyl)-5-methyl-benzyloxy)-3-phenyl-morpholine;
- 62) 4-(2-(imidazolo)methyl)-2-(3-(tert-butyl)-5-(trifluoromethyl)-benzyloxy)-3-phenyl-morpholine;
- 10 63) 2-(S)-(3,5-dichlorobenzyloxy)-3-(S)-phenyl-morpholine;
- 64) 2-(S)-(3,5-dichlorobenzyloxy)-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-3-(S)-phenylmorpholine;
- 15 65) 2-(S)-(3,5-bis(trifluoromethyl)benzyloxy)-4-(methoxycarbonylmethyl)-3-(S)-phenyl-morpholine;
- 66) 2-(S)-(3,5-bis(trifluoromethyl)benzyloxy)-4-(carboxymethyl)-3-(S)-phenyl-morpholine;
- 20 67) 2-(S)-(3,5-bis(trifluoromethyl)benzyloxy)-4-((2-aminoethyl)-aminocarbonylmethyl)-3-(S)-phenyl-morpholine;
- 25 68) 2-(S)-(3,5-bis(trifluoromethyl)benzyloxy)-4-((3-aminopropyl)-amino carbonylmethyl)-3-(S)-phenylmorpholine;
- 69) 4-benzyl-5-(S),6-(R)-dimethyl-3-(S)-phenylmorpholinone and 4-benzyl-5-(R),6-(S)-dimethyl-3-(S)-phenyl-morpholinone;
- 30 70) 2-(R)-(3,5-bis(trifluoromethyl)benzyloxy)-[5-(S),6-(R) or 5-(R),6-(S)-dimethyl]-3-(S)-phenyl-morpholinone;
- 71) 2-(S)-(3,5-bis(trifluoromethyl)benzyloxy)-[5-(R),6-(S) or 5-

- 49 -

(S),6-(R)-dimethyl]-3-(S)-phenyl-morpholinone;

- 5 72) 2-(R)-(3,5-bis(trifluoromethyl)benzyloxy)-4-(3-(1,2,4-triazolo)methyl)-[5-(S),6-(R) or 5-(R),6-(S)-dimethyl]-3-(S)-phenyl-morpholinone;
- 10 73) 2-(R)-(3,5-bis(trifluoromethyl)benzyloxy)-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-[5-(S),6-(R) or 5-(R),6-(S)-dimethyl]-3-(S)-phenyl-morpholinone;
- 15 74) 2-(S)-(3,5-bis(trifluoromethyl)benzyloxy)-4-(3-(1,2,4-triazolo)methyl)-[5-(R),6-(S) or 5-(S),6-(R)-dimethyl]-3-(S)-phenyl-morpholinone;
- 20 75) 2-(S)-(3,5-bis(trifluoromethyl)benzyloxy)-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-[5-(R),6-(S) or 5-(S),6-(R)-dimethyl]-3-(S)-phenyl-morpholinone;
- 25 76) 2-(S)-(3,5-bis(trifluoromethyl)benzyloxy)-4-(2-(1-(4-benzyl)-piperidino)ethyl)-3-(S)-phenyl-morpholine;
- 77) 3-(S)-(4-fluorophenyl)-4-benzyl-2-morpholinone;
- 28 78) 2-(S)-(3,5-bis(trifluoromethyl)benzyloxy)-3-(S)-(4-fluorophenyl)-4-benzyl-morpholine;
- 30 79) 2-(S)-(3,5-Bis(trifluoromethyl)benzyloxy)-3-(S)-(4-fluorophenyl)morpholine;
- 80) 2-(S)-(3,5-bis(trifluoromethyl)benzyloxy)-3-(S)-(4-fluorophenyl)-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 81) 2-(S)-(3,5-bis(trifluoromethyl)benzyloxy)-4-((3-pyridyl)methyl carbonyl)-3-(R)-phenyl-morpholine;

- 50 -

- 82) 2-(S)-(3,5-bis(trifluoromethyl)benzyloxy)-4-(methoxycarbonyl-pentyl)-3-(R)-phenyl-morpholine;
- 5 83) 2-(S)-(3,5-bis(trifluoromethyl)benzyloxy)-4-(carboxypentyl)-3-(R)-phenyl-morpholine;
- 84) 2-(S)-(3,5-bis(trifluoromethyl)benzyloxy)-4-(methylamino-carbonylpentyl)-6-oxo-hexyl)-3-(R)-phenyl-morpholine;
- 10 85) 2-(R)-(3,5-bis(trifluoromethyl)benzoyloxy)-3-(S)-phenyl-4-benzyl-morpholine;
- 86) 2-(R)-(1-(3,5-bis(trifluoromethyl)phenyl)ethenyloxy)-3-(S)-phenyl-4-benzyl-morpholine;
- 15 87) 2-(R)-(1-(S)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-phenyl-morpholine;
- 88) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-phenyl-morpholine;
- 20 89) 2-(R)-(1-(S)-(3,5-Bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl-morpholine;
- 25 90) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl-morpholine;
- 91) 2-(R)-(3,5-bis(trifluoromethyl)benzoyloxy)-3-(S)-(4-fluoro)-phenyl-4-benzyl-morpholine;
- 30 92) 2-(R)-(1-(3,5-bis(trifluoromethyl)phenyl)ethenyloxy)-3-(S)-(4-fluoro)phenyl-4-benzyl-morpholine;

- 51 -

- 93) 2-(R)-(1-(S)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-morpholine;
- 5 94) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-morpholine;
- 95) 2-(R)-(1-(S)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 10 96) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 15 97) 2-(R)-(1-(R)-(1-(3-(methyl)naphthyl))ethoxy)-3-(S)-phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 98) 2-(R)-(1-(R)-(1-(3-(methyl)naphthyl))ethoxy)-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 20 99) 2-(R)-(1-(R)-(3-(fluoro)-5-(trifluoromethyl)phenyl)ethoxy)-3-(S)-phenyl-morpholine;
- 100) 2-(R)-(1-(R)-(3-(fluoro)-5-(trifluoromethyl)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 25 101) 2-(R)-(1-(R)-(3-(chloro)-5-(trifluoromethyl)phenyl)ethoxy)-3-(S)-phenyl-morpholine;
- 30 102) 2-(R)-(1-(R)-(3-(chloro)-5-(trifluoromethyl)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;

- 52 -

- 103) 2-(R)-(1-(R)-(3,5-(dimethyl)phenyl)ethoxy)-3-(S)-phenyl-morpholine;
- 5 104) 2-(R)-(1-(R)-(3,5-(dimethyl)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 105) 2-(R)-(1-(R)-(3-(fluoro)-5-(methyl)phenyl)ethoxy)-3-(S)-phenyl-morpholine;
- 10 106) 2-(R)-(1-(R)-(3-(fluoro)-5-(methyl)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 107) 2-(R)-(1-(R)-(3-(chloro)-5-(methyl)phenyl)ethoxy)-3-(S)-phenyl-morpholine;
- 15 108) 2-(R)-(1-(R)-(3-(chloro)-5-(methyl)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 109) 2-(R)-(1-(R)-(3-(bromo)-5-(methyl)phenyl)ethoxy)-3-(S)-phenyl-morpholine;
- 20 110) 2-(R)-(1-(R)-(3-(bromo)-5-(methyl)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 111) 2-(R)-(1-(R)-(3-(isopropoxy)phenyl)ethoxy)-3-(S)-phenyl-morpholine;
- 25 112) 2-(R)-(1-(R)-(3-(isopropoxy)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 30 113) 2-(R)-(1-(R)-(3-(isopropoxy)-5-trifluoromethyl)phenyl)ethoxy)-3-(S)-phenyl-morpholine;

- 53 -

- 114) 2-(R)-(1-(R)-(3-(isopropoxy)-5-trifluoromethyl)phenyl)ethoxy)-
3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-
morpholine;
- 5 115) 2-(R)-(1-(R)-(3-(chloro)-5-(isopropoxy)phenyl)ethoxy)-3-(S)-
phenyl-morpholine;
- 116) 2-(R)-(1-(R)-(3-(chloro)-5-(isopropoxy)phenyl)ethoxy)-3-(S)-
10 phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 117) 2-(R)-(1-(R)-(3-(fluoro)-5-(isopropoxy)phenyl)ethoxy)-3-(S)-
phenyl-morpholine;
- 118) 2-(R)-(1-(R)-(3-(fluoro)-5-(isopropoxy)phenyl)ethoxy)-3-(S)-
15 phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 119) 2-(R)-(1-(R)-(1-(3-(trifluoromethyl)naphthyl))ethoxy)-3-(S)-
phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 20 120) 2-(R)-(1-(R)-(1-(3-(trifluoromethyl)naphthyl))ethoxy)-3-(S)-
phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 121) 2-(R)-(1-(R)-(3-(t-butyl)-5-(chloro)phenyl)ethoxy)-3-(S)-phenyl-
25 morpholine;
- 122) 2-(R)-(1-(R)-(3-(t-butyl)-5-(chloro)phenyl)ethoxy)-3-(S)-phenyl-
4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 123) 2-(R)-(1-(R)-(3-(t-butyl)-5-(trifluoromethyl)phenyl)ethoxy)-3-
30 (S)-phenyl-morpholine;
- 124) 2-(R)-(1-(R)-(3-(t-butyl)-5-(trifluoromethyl)phenyl)ethoxy)-3-
(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-
morpholine;

- 54 -

- 125) 2-(R)-(1-(R)-(3,5-(dimethyl)-4-(fluoro)phenyl)ethoxy)-3-(S)-phenyl-morpholine;
- 5 126) 2-(R)-(1-(R)-(3,5-(dimethyl)-4-(fluoro)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 127) 2-(R)-(1-(R)-(3,5-(dimethyl)-4-(chloro)phenyl)ethoxy)-3-(S)-phenyl-morpholine;
- 10 128) 2-(R)-(1-(R)-(3,5-(dimethyl)-4-(chloro)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 129) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)-4-(fluoro)phenyl)ethoxy)-3-(S)-phenyl-morpholine;
- 15 130) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)-4-(fluoro)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 20 131) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)-4-chloro)phenyl)ethoxy)-3-(S)-phenyl-morpholine;
- 132) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)-4-chloro)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 25 133) 2-(R)-(1-(R)-(3,5-(dichloro)phenyl)ethoxy)-3-(S)-phenyl-morpholine;
- 30 134) 2-(R)-(1-(R)-(3,5-(dichloro)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;

- 55 -

- 135) 2-(R)-(1-(R)-(3,5-(difluoro)phenyl)ethoxy)-3-(S)-phenyl-morpholine;
- 5 136) 2-(R)-(1-(R)-(3,5-(difluoro)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 137) 2-(R)-(1-(R)-(1-(naphthyl)ethoxy)-3-(S)-phenyl-morpholine;
- 10 138) 2-(R)-(1-(R)-(1-(naphthyl)ethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)morpholine;
- 139) 2-(R)-(1-(R)-(1-(4-(fluoro)naphthyl))ethoxy)-3-(S)-phenyl-morpholine;
- 15 140) 2-(R)-(1-(R)-(1-(4-(fluoro)naphthyl))ethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 141) 2-(R)-(1-(R)-(1-(3-(fluoro)naphthyl))ethoxy)-3-(S)-phenyl-morpholine;
- 20 142) 2-(R)-(1-(R)-(1-(3-(fluoro)naphthyl))ethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 25 143) 2-(R)-(1-(R)-(1-(3-(chloro)naphthyl))ethoxy)-3-(S)-phenyl-morpholine;
- 144) 2-(R)-(1-(R)-(1-(3-(chloro)naphthyl))ethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 30 145) 2-(R)-(1-(R)-(1-(3-(methyl)naphthyl))ethoxy)-3-(S)-phenyl-morpholine;
- 146) 2-(R)-(1-(R)-(1-(3-(methyl)naphthyl))ethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;

- 56 -

- 147) 2-(R)-(1-(R)-(1-(3-(trifluoromethyl)naphthyl))ethoxy)-3-(S)-phenyl-morpholine;
- 5 148) 2-(R)-(1-(R)-(1-(3-(trifluoromethyl)naphthyl))ethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 149) 2-(S)-(3,5-bis(trifluoromethyl)benzyloxy)-3-(S)-(4-hydroxy)-phenyl-morpholine;
- 10 150) 2-(S)-(3,5-bis(trifluoromethyl)benzyloxy)-3-(S)-(4-hydroxy)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 15 151) 2-(S)-(2-fluoro-5-trifluoromethyl)benzyloxy-3-(S)-phenyl-morpholine;
- 152) 2-(S)-(2-fluoro-5-trifluoromethyl)benzyloxy-3-(S)-(4-fluoro)phenyl-morpholine;
- 20 153) 2-(R)-(1-(R)-(2-fluoro-5-trifluoromethyl)phenylethoxy)-3-(S)-phenyl-morpholine;
- 154) 2-(R)-(1-(R)-(2-fluoro-5-trifluoromethyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-morpholine;
- 25 155) 2-(S)-(2-fluoro-5-trifluoromethyl)benzyloxy-3-(S)-phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 156) 2-(S)-(2-fluoro-5-trifluoromethyl)benzyloxy-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 30 157) 2-(R)-(1-(R)-(2-fluoro-5-trifluoromethyl)phenylethoxy)-3-(S)-phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;

- 57 -

- 158) 2-(R)-(1-(R)-(2-fluoro-5-trifluoromethyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 5 159) 2-(S)-(2-fluoro-5-trifluoromethyl)benzyloxy-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 160) 2-(S)-(2-fluoro-5-trifluoromethyl)benzyloxy-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 10 161) 2-(R)-(1-(R)-(2-fluoro-5-trifluoromethyl)phenylethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 162) 2-(R)-(1-(R)-(2-fluoro-5-trifluoromethyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 15 163) 2-(S)-(2-fluoro-5-trifluoromethyl)benzyloxy-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 20 164) 2-(S)-(2-fluoro-5-trifluoromethyl)benzyloxy-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 165) 2-(R)-(1-(R)-(2-fluoro-5-trifluoromethyl)phenylethoxy)-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 25 166) 2-(R)-(1-(R)-(2-fluoro-5-trifluoromethyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 30 167) 2-(S)-(2-fluoro-5-trifluoromethyl)benzyloxy-3-(S)-phenyl-4-(2-imidazolo)methyl-morpholine;
- 168) 2-(S)-(2-fluoro-5-trifluoromethyl)benzyloxy-3-(S)-(4-fluoro)phenyl-4-(2-imidazolo)methyl-morpholine;

- 58 -

- 169) 2-(R)-(1-(R)-(2-fluoro-5-trifluoromethyl)phenylethoxy)-3-(S)-phenyl-4-(2-imidazolo)methyl-morpholine;
- 5 170) 2-(R)-(1-(R)-(2-fluoro-5-trifluoromethyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(2-imidazolo)methyl-morpholine;
- 171) 2-(S)-(2-fluoro-5-trifluoromethyl)benzyloxy-3-(S)-phenyl-4-(4-imidazolo)methyl-morpholine;
- 10 172) 2-(S)-(2-fluoro-5-trifluoromethyl)benzyloxy-3-(S)-(4-fluoro)phenyl-4-(4-imidazolo)methyl-morpholine;
- 173) 2-(R)-(1-(R)-(2-fluoro-5-trifluoromethyl)phenylethoxy)-3-(S)-phenyl-4-(4-imidazolo)methyl-morpholine;
- 15 174) 2-(R)-(1-(R)-(2-fluoro-5-trifluoromethyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-imidazolo)methyl-morpholine;
- 20 175) 2-(S)-(2-fluoro-5-trifluoromethyl)benzyloxy-3-(S)-phenyl-4-(5-tetrazolo)methyl-morpholine;
- 176) 2-(S)-(2-fluoro-5-trifluoromethyl)benzyloxy-3-(S)-(4-fluoro)phenyl-4-(5-tetrazolo)methyl-morpholine;
- 25 177) 2-(R)-(1-(R)-(2-fluoro-5-trifluoromethyl)phenylethoxy)-3-(S)-phenyl-4-(5-tetrazolo)methyl-morpholine;
- 178) 2-(R)-(1-(R)-(2-fluoro-5-trifluoromethyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(5-tetrazolo)methyl-morpholine;
- 30 179) 2-(S)-(2-fluoro-5-trifluoromethyl)benzyloxy-3-(S)-phenyl-4-(2-oxo-5H-pyrrol-4-yl)methyl-morpholine;

- 59 -

- 180) 2-(S)-(2-fluoro-5-trifluoromethyl)benzyloxy-3-(S)-(4-fluoro)phenyl-4-(2-oxo-5H-pyrrol-4-yl)methyl-morpholine;
- 5 181) 2-(R)-(1-(R)-(2-fluoro-5-trifluoromethyl)phenylethoxy)-3-(S)-phenyl-4-(2-oxo-5H-pyrrol-4-yl)methyl-morpholine;
- 182) 2-(R)-(1-(R)-(2-fluoro-5-trifluoromethyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(2-oxo-5H-pyrrol-4-yl)methyl-morpholine;
- 10 183) 2-(S)-(2-chloro-5-trifluoromethyl)benzyloxy-3-(S)-phenyl-morpholine;
- 184) 2-(S)-(2-chloro-5-trifluoromethyl)benzyloxy-3-(S)-(4-fluoro)phenyl-morpholine;
- 15 185) 2-(R)-(1-(R)-(2-chloro-5-trifluoromethyl)phenylethoxy)-3-(S)-phenyl-morpholine;
- 186) 2-(R)-(1-(R)-(2-chloro-5-trifluoromethyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-morpholine;
- 20 187) 2-(S)-(2-chloro-5-trifluoromethyl)benzyloxy-3-(S)-phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 25 188) 2-(S)-(2-chloro-5-trifluoromethyl)benzyloxy-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 189) 2-(R)-(1-(R)-(2-chloro-5-trifluoromethyl)phenylethoxy)-3-(S)-phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 30 190) 2-(R)-(1-(R)-(2-chloro-5-trifluoromethyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;

- 60 -

- 191) 2-(S)-(2-chloro-5-trifluoromethyl)benzyloxy-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 5 192) 2-(S)-(2-chloro-5-trifluoromethyl)benzyloxy-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 10 193) 2-(R)-(1-(R)-(2-chloro-5-trifluoromethyl)phenylethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 194) 2-(R)-(1-(R)-(2-chloro-5-trifluoromethyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 15 195) 2-(S)-(2-chloro-5-trifluoromethyl)benzyloxy-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 20 196) 2-(S)-(2-chloro-5-trifluoromethyl)benzyloxy-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 197) 2-(R)-(1-(R)-(2-chloro-5-trifluoromethyl)phenylethoxy)-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 25 198) 2-(R)-(1-(R)-(2-chloro-5-trifluoromethyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 199) 2-(S)-(2-chloro-5-trifluoromethyl)benzyloxy-3-(S)-phenyl-4-(2-imidazolo)methyl)-morpholine;
- 30 200) 2-(S)-(2-chloro-5-trifluoromethyl)benzyloxy-3-(S)-(4-fluoro)phenyl-4-(2-imidazolo)methyl)-morpholine;
- 201) 2-(R)-(1-(R)-(2-chloro-5-trifluoromethyl)phenylethoxy)-3-(S)-phenyl-4-(2-imidazolo)methyl)-morpholine;

- 61 -

- 202) 2-(R)-(1-(R)-(2-chloro-5-trifluoromethyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(2-imidazolo)methyl-morpholine;
- 5 203) 2-(S)-(2-chloro-5-trifluoromethyl)benzyloxy-3-(S)-phenyl-4-(4-imidazolo)methyl-morpholine;
- 204) 2-(S)-(2-chloro-5-trifluoromethyl)benzyloxy-3-(S)-(4-fluoro)phenyl-4-(4-imidazolo)methyl-morpholine;
- 10 205) 2-(R)-(1-(R)-(2-chloro-5-trifluoromethyl)phenylethoxy)-3-(S)-phenyl-4-(4-imidazolo)methyl-morpholine;
- 206) 2-(R)-(1-(R)-(2-chloro-5-trifluoromethyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-imidazolo)methyl-morpholine;
- 15 207) 2-(S)-(2-chloro-5-trifluoromethyl)benzyloxy-3-(S)-phenyl-4-(5-tetrazolo)methyl-morpholine;
- 20 208) 2-(S)-(2-chloro-5-trifluoromethyl)benzyloxy-3-(S)-(4-fluoro)phenyl-4-(5-tetrazolo)methyl-morpholine;
- 209) 2-(R)-(1-(R)-(2-chloro-5-trifluoromethyl)phenylethoxy)-3-(S)-phenyl-4-(5-tetrazolo)methyl-morpholine;
- 25 210) 2-(R)-(1-(R)-(2-chloro-5-trifluoromethyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(5-tetrazolo)methyl-morpholine;
- 211) 2-(S)-(2-chloro-5-trifluoromethyl)benzyloxy-3-(S)-phenyl-4-(2-oxo-5H-pyrrol-4-yl)methyl-morpholine;
- 30 212) 2-(S)-(2-chloro-5-trifluoromethyl)benzyloxy-3-(S)-(4-fluoro)phenyl-4-(2-oxo-5H-pyrrol-4-yl)methyl-morpholine;

- 62 -

- 213) 2-(R)-(1-(R)-(2-chloro-5-trifluoromethyl)phenylethoxy)-3-(S)-phenyl-4-(2-oxo-5H-pyrrol-4-yl)methyl-morpholine;
- 5 214) 2-(R)-(1-(R)-(2-Chloro-5-trifluoromethyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(2-oxo-5H-pyrrol-4-yl)methyl-morpholine;
- 215) 2-(S)-(3-methyl)benzyloxy-3-(S)-phenyl-morpholine;
- 10 216) 2-(S)-(3-methyl)benzyloxy-3-(S)-(4-fluoro)phenyl-morpholine;
- 217) 2-(R)-(1-(R)-(3-methyl)phenylethoxy)-3-(S)-phenyl-morpholine;
- 218) 2-(R)-(1-(R)-(3-methyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-morpholine;
- 15 219) 2-(S)-(3-methyl)benzyloxy-3-(S)-phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 20 220) 2-(S)-(3-methyl)benzyloxy-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 221) 2-(R)-(1-(R)-(3-methyl)phenylethoxy)-3-(S)-phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 25 222) 2-(R)-(1-(R)-(3-methyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 223) 2-(S)-(3-methyl)benzyloxy-3-(S)-phenyl-4-(3-(5-oxo-1,2,4-triazolo)methyl)-morpholine;
- 30 224) 2-(S)-(3-methyl)benzyloxy-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1,2,4-triazolo)methyl)-morpholine;

- 63 -

- 225) 2-(R)-(1-(R)-(3-methyl)phenylethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1,2,4-triazolo)methyl)-morpholine;
- 5 226) 2-(R)-(1-(R)-(3-methyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1,2,4-triazolo)methyl)-morpholine;
- 227) 2-(S)-(3-methyl)benzyloxy-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 10 228) 2-(S)-(3-methyl)benzyloxy-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 229) 2-(R)-(1-(R)-(3-methyl)phenylethoxy)-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 15 230) 2-(R)-(1-(R)-(3-methyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 20 231) 2-(S)-(3-methyl)benzyloxy-3-(S)-phenyl-4-(2-imidazolo)methyl-morpholine;
- 232) 2-(S)-(3-methyl)benzyloxy-3-(S)-(4-fluoro)phenyl-4-(2-imidazolo)methyl-morpholine;
- 25 233) 2-(R)-(1-(R)-(3-methyl)phenylethoxy)-3-(S)-phenyl-4-(2-imidazolo)methyl-morpholine;
- 234) 2-(R)-(1-(R)-(3-methyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(2-imidazolo)methyl-morpholine;
- 30 235) 2-(S)-(3-methyl)benzyloxy-3-(S)-phenyl-4-(4-imidazolo)methyl-morpholine;

- 64 -

- 236) 2-(S)-(3-methyl)benzyloxy-3-(S)-(4-fluoro)phenyl-4-(4-imidazolo)methyl-morpholine;
- 5 237) 2-(R)-(1-(R)-(3-methyl)phenylethoxy)-3-(S)-phenyl-4-(4-imidazolo)methyl-morpholine;
- 238) 2-(R)-(1-(R)-(3-methyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-imidazolo)methyl-morpholine;
- 10 239) 2-(S)-(3-methyl)benzyloxy-3-(S)-phenyl-4-(5-tetrazolo)methyl-morpholine;
- 240) 2-(S)-(3-methyl)benzyloxy-3-(S)-(4-fluoro)phenyl-4-(5-tetrazolo)methyl-morpholine;
- 15 241) 2-(R)-(1-(R)-(3-methyl)phenylethoxy)-3-(S)-phenyl-4-(5-tetrazolo)methyl-morpholine;
- 242) 2-(R)-(1-(R)-(3-methyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(5-tetrazolo)methyl-morpholine;
- 20 243) 2-(S)-(3-methyl)benzyloxy-3-(S)-phenyl-4-(2-oxo-5H-pyrrol-4-yl)methyl-morpholine;
- 25 244) 2-(S)-(3-methyl)benzyloxy-3-(S)-(4-fluoro)phenyl-4-(2-oxo-5H-pyrrol-4-yl)methyl-morpholine;
- 245) 2-(R)-(1-(R)-(3-methyl)phenylethoxy)-3-(S)-phenyl-4-(2-oxo-5H-pyrrol-4-yl)methyl-morpholine;
- 30 246) 2-(R)-(1-(R)-(3-methyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(2-oxo-5H-pyrrol-4-yl)methyl-morpholine;
- 247) 2-(S)-(3-bromo)benzyloxy-3-(S)-phenyl-morpholine;

- 65 -

- 248) 2-(S)-(3-bromo)benzyloxy-3-(S)-(4-fluoro)phenyl-morpholine;
- 5 249) 2-(R)-(1-(R)-(3-bromo)phenylethoxy)-3-(S)-phenyl-morpholine;
- 250) 2-(R)-(1-(R)-(3-bromo)phenylethoxy)-3-(S)-(4-fluoro)phenyl-morpholine;
- 10 251) 2-(S)-(3-bromo)benzyloxy-3-(S)-phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 252) 2-(S)-(3-bromo)benzyloxy-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 15 253) 2-(R)-(1-(R)-(3-bromo)phenylethoxy)-3-(S)-phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 254) 2-(R)-(1-(R)-(3-bromo)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 20 255) 2-(S)-(3-bromo)benzyloxy-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 25 256) 2-(S)-(3-bromo)benzyloxy-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 257) 2-(R)-(1-(R)-(3-bromo)phenylethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 30 258) 2-(R)-(1-(R)-(3-bromo)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 259) 2-(S)-(3-bromo)benzyloxy-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;

- 66 -

- 260) 2-(S)-(3-bromo)benzyloxy-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 5 261) 2-(R)-(1-(R)-(3-bromo)phenylethoxy)-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 262) 2-(R)-(1-(R)-(3-bromo)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 10 263) 2-(S)-(3-bromo)benzyloxy-3-(S)-phenyl-4-(2-imidazolo)methyl-morpholine;
- 264) 2-(S)-(3-bromo)benzyloxy-3-(S)-(4-fluoro)phenyl-4-(2-imidazolo)methyl-morpholine;
- 15 265) 2-(R)-(1-(R)-(3-bromo)phenylethoxy)-3-(S)-phenyl-4-(2-imidazolo)methyl-morpholine;
- 266) 2-(R)-(1-(R)-(3-bromo)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(2-imidazolo)methyl-morpholine;
- 20 267) 2-(S)-(3-bromo)benzyloxy-3-(S)-phenyl-4-(4-imidazolo)methyl-morpholine;
- 25 268) 2-(S)-(3-bromo)benzyloxy-3-(S)-(4-fluoro)phenyl-4-(4-imidazolo)methyl-morpholine;
- 269) 2-(R)-(1-(R)-(3-bromo)phenylethoxy)-3-(S)-phenyl-4-(4-imidazolo)methyl-morpholine;
- 30 270) 2-(R)-(1-(R)-(3-bromo)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-imidazolo)methyl-morpholine;

- 67 -

- 271) 2-(S)-(3-bromo)benzyloxy-3-(S)-phenyl-4-(5-tetrazolo)methyl-morpholine;
- 5 272) 2-(S)-(3-bromo)benzyloxy-3-(S)-(4-fluoro)phenyl-4-(5-tetrazolo)methyl-morpholine;
- 273) 2-(R)-(1-(R)-(3-bromo)phenylethoxy)-3-(S)-phenyl-4-(5-tetrazolo)methyl-morpholine;
- 10 274) 2-(R)-(1-(R)-(3-bromo)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(5-tetrazolo)methyl-morpholine;
- 275) 2-(S)-(3-bromo)benzyloxy-3-(S)-phenyl-4-(2-oxo-5H-pyrrol-4-yl)methyl-morpholine;
- 15 276) 2-(S)-(3-bromo)benzyloxy-3-(S)-(4-fluoro)phenyl-4-(2-oxo-5H-pyrrol-4-yl)methyl-morpholine;
- 20 277) 2-(R)-(1-(R)-(3-bromo)phenylethoxy)-3-(S)-phenyl-4-(2-oxo-5H-pyrrol-4-yl)methyl-morpholine;
- 278) 2-(R)-(1-(R)-(3-bromo)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(2-oxo-5H-pyrrol-4-yl)methyl-morpholine;
- 25 279) 2-(S)-(3-chloro)benzyloxy-3-(S)-phenyl-morpholine;
- 280) 2-(S)-(3-chloro)benzyloxy-3-(S)-(4-fluoro)phenyl-morpholine;
- 30 281) 2-(R)-(1-(R)-(3-chloro)phenylethoxy)-3-(S)-phenyl-morpholine;
- 282) 2-(R)-(1-(R)-(3-chloro)phenylethoxy)-3-(S)-(4-fluoro)phenyl-morpholine;

- 68 -

- 283) 2-(S)-(3-chloro)benzyloxy-3-(S)-phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 5 284) 2-(S)-(3-chloro)benzyloxy-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 285) 2-(R)-(1-(R)-(3-chloro)phenylethoxy)-3-(S)-phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 10 286) 2-(R)-(1-(R)-(3-chloro)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 287) 2-(S)-(3-chloro)benzyloxy-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 15 288) 2-(S)-(3-chloro)benzyloxy-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 289) 2-(R)-(1-(R)-(3-chloro)phenylethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 20 290) 2-(R)-(1-(R)-(3-chloro)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 25 291) 2-(S)-(3-chloro)benzyloxy-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 292) 2-(S)-(3-chloro)benzyloxy-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 30 293) 2-(R)-(1-(R)-(3-chloro)phenylethoxy)-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;

- 69 -

- 294) 2-(R)-(1-(R)-(3-chloro)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 5 295) 2-(S)-(3-chloro)benzyloxy-3-(S)-phenyl-4-(2-imidazolo)methyl-morpholine;
- 296) 2-(S)-(3-chloro)benzyloxy-3-(S)-(4-fluoro)phenyl-4-(2-imidazolo)methyl-morpholine;
- 10 297) 2-(R)-(1-(R)-(3-chloro)phenylethoxy)-3-(S)-phenyl-4-(2-imidazolo)methyl-morpholine;
- 298) 2-(R)-(1-(R)-(3-chloro)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(2-imidazolo)methyl-morpholine;
- 15 299) 2-(S)-(3-chloro)benzyloxy-3-(S)-phenyl-4-(4-imidazolo)methyl-morpholine;
- 300) 2-(S)-(3-chloro)benzyloxy-3-(S)-(4-fluoro)phenyl-4-(4-imidazolo)methyl-morpholine;
- 20 301) 2-(R)-(1-(R)-(3-chloro)phenylethoxy)-3-(S)-phenyl-4-(4-imidazolo)methyl-morpholine;
- 25 302) 2-(R)-(1-(R)-(3-chloro)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-imidazolo)methyl-morpholine;
- 303) 2-(S)-(3-chloro)benzyloxy-3-(S)-phenyl-4-(5-tetrazolo)methyl-morpholine;
- 30 304) 2-(S)-(3-chloro)benzyloxy-3-(S)-(4-fluoro)phenyl-4-(5-tetrazolo)methyl-morpholine;

- 70 -

- 305) 2-(R)-(1-(R)-(3-chloro)phenylethoxy)-3-(S)-phenyl-4-(5-tetrazolo)methyl-morpholine;
- 5 306) 2-(R)-(1-(R)-(3-chloro)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(5-tetrazolo)methyl-morpholine;
- 307) 2-(S)-(3-chloro)benzyloxy-3-(S)-phenyl-4-(2-oxo-5H-pyrrol-4-yl)methyl-morpholine;
- 10 308) 2-(S)-(3-chloro)benzyloxy-3-(S)-(4-fluoro)phenyl-4-(2-oxo-5H-pyrrol-4-yl)methyl-morpholine;
- 309) 2-(R)-(1-(R)-(3-chloro)phenylethoxy)-3-(S)-phenyl-4-(2-oxo-5H-pyrrol-4-yl)methyl-morpholine;
- 15 310) 2-(R)-(1-(R)-(3-chloro)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(2-oxo-5H-pyrrol-4-yl)methyl-morpholine;
- 20 311) 2-(S)-(3-trifluoromethyl)benzyloxy-3-(S)-phenyl-morpholine;
- 312) 2-(S)-(3-trifluoromethyl)benzyloxy-3-(S)-(4-fluoro)phenyl-morpholine;
- 25 313) 2-(R)-(1-(R)-(3-trifluoromethyl)phenylethoxy)-3-(S)-phenyl-morpholine;
- 314) 2-(R)-(1-(R)-(3-trifluoromethyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-morpholine;
- 30 315) 2-(S)-(3-trifluoromethyl)benzyloxy-3-(S)-phenyl-4-(3-1H,4H-1,2,4-triazolo)methyl-morpholine;
- 316) 2-(S)-(3-trifluoromethyl)benzyloxy-3-(S)-(4-fluoro)phenyl-4-(3-1H,4H-1,2,4-triazolo)methyl-morpholine;

- 71 -

- 317) 2-(R)-(1-(R)-(3-trifluoromethyl)phenylethoxy)-3-(S)-phenyl-4-(3-1H,4H-1,2,4-triazolo)methyl-morpholine;
- 5 318) 2-(R)-(1-(R)-(3-trifluoromethyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl-morpholine;
- 319) 2-(S)-(3-trifluoromethyl)benzyloxy-3-(S)-phenyl-4-(3-(5-oxo)-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 10 320) 2-(S)-(3-trifluoromethyl)benzyloxy-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 321) 2-(R)-(1-(R)-(3-trifluoromethyl)phenylethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 15 322) 2-(R)-(1-(R)-(3-trifluoromethyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 20 323) 2-(S)-(3-trifluoromethyl)benzyloxy-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 324) 2-(S)-(3-trifluoromethyl)benzyloxy-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 25 325) 2-(R)-(1-(R)-(3-trifluoromethyl)phenylethoxy)-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 30 326) 2-(R)-(1-(R)-(3-trifluoromethyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 327) 2-(S)-(3-trifluoromethyl)benzyloxy-3-(S)-phenyl-4-(2-imidazolo)methyl-morpholine;

- 72 -

- 328) 2-(S)-(3-trifluoromethyl)benzyloxy-3-(S)-(4-fluoro)phenyl-4-(2-imidazolo)methyl-morpholine;
- 5 329) 2-(R)-(1-(R)-(3-trifluoromethyl)phenylethoxy)-3-(S)-phenyl-4-(2-imidazolo)methyl-morpholine;
- 330) 2-(R)-(1-(R)-(3-trifluoromethyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(2-imidazolo)methyl-morpholine;
- 10 331) 2-(S)-(3-trifluoromethyl)benzyloxy-3-(S)-phenyl-4-(4-imidazolo)methyl-morpholine;
- 332) 2-(S)-(3-trifluoromethyl)benzyloxy-3-(S)-(4-fluoro)phenyl-4-(4-imidazolo)methyl-morpholine;
- 15 333) 2-(R)-(1-(R)-(3-trifluoromethyl)phenylethoxy)-3-(S)-phenyl-4-(4-imidazolo)methyl-morpholine;
- 20 334) 2-(R)-(1-(R)-(3-trifluoromethyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-imidazolo)methyl-morpholine;
- 335) 2-(S)-(3-trifluoromethyl)benzyloxy-3-(S)-phenyl-4-(5-tetrazolo)methyl-morpholine;
- 25 336) 2-(S)-(3-trifluoromethyl)benzyloxy-3-(S)-(4-fluoro)phenyl-4-(5-tetrazolo)methyl-morpholine;
- 337) 2-(R)-(1-(R)-(3-trifluoromethyl)phenylethoxy)-3-(S)-phenyl-4-(5-tetrazolo)methyl-morpholine;
- 30 338) 2-(R)-(1-(R)-(3-trifluoromethyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(5-tetrazolo)methyl-morpholine;

- 73 -

- 339) 2-(S)-(3-trifluoromethyl)benzyloxy-3-(S)-phenyl-4-(2-oxo-5H-pyrrol-4-yl)methyl-morpholine;
- 5 340) 2-(S)-(3-trifluoromethyl)benzyloxy-3-(S)-(4-fluoro)phenyl-4-(2-oxo-5H-pyrrol-4-yl)methyl-morpholine;
- 341) 2-(R)-(1-(R)-(3-trifluoromethyl)phenylethoxy)-3-(S)-phenyl-4-(2-oxo-5H-pyrrol-4-yl)methyl-morpholine;
- 10 342) 2-(R)-(1-(R)-(3-trifluoromethyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(2-oxo-5H-pyrrol-4-yl)methyl-morpholine;
- 343) 2-(S)-(3-t-butyl)benzyloxy-3-(S)-phenyl-morpholine;
- 15 344) 2-(S)-(3-t-butyl)benzyloxy-3-(S)-(4-fluoro)phenyl-morpholine;
- 345) 2-(R)-(1-(R)-(3-t-butyl)phenylethoxy)-3-(S)-phenyl-morpholine;
- 20 346) 2-(R)-(1-(R)-(3-t-butyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-morpholine;
- 347) 2-(S)-(3-t-butyl)benzyloxy-3-(S)-phenyl-4-(3-(1,2,4-triazolo)methyl-morpholine;
- 25 348) 2-(S)-(3-t-butyl)benzyloxy-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl-morpholine;
- 349) 2-(R)-(1-(R)-(3-t-butyl)phenylethoxy)-3-(S)-phenyl-4-(3-(1,2,4-triazolo)methyl-morpholine;
- 30 350) 2-(R)-(1-(R)-(3-t-butyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl-morpholine;

- 74 -

- 351) 2-(S)-(3-t-butyl)benzyloxy-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl-morpholine;
- 5 352) 2-(S)-(3-t-butyl)benzyloxy-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl-morpholine;
- 353) 2-(R)-(1-(R)-(3-t-butyl)phenylethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 10 354) 2-(R)-(1-(R)-(3-t-butyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 355) 2-(S)-(3-t-butyl)benzyloxy-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl-morpholine;
- 15 356) 2-(S)-(3-t-butyl)benzyloxy-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl-morpholine;
- 20 357) 2-(R)-(1-(R)-(3-t-butyl)phenylethoxy)-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl-morpholine;
- 358) 2-(R)-(1-(R)-(3-t-butyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl-morpholine;
- 25 359) 2-(S)-(3-t-butyl)benzyloxy-3-(S)-phenyl-4-(2-imidazolo)methyl-morpholine;
- 360) 2-(S)-(3-t-butyl)benzyloxy-3-(S)-(4-fluoro)phenyl-4-(2-imidazolo)methyl-morpholine;
- 30 361) 2-(R)-(1-(R)-(3-t-butyl)phenylethoxy)-3-(S)-phenyl-4-(2-imidazolo)methyl-morpholine;

- 75 -

- 362) 2-(R)-(1-(R)-(3-t-butyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(2-imidazolo)methyl-morpholine;
- 5 363) 2-(S)-(3-t-butyl)benzyloxy-3-(S)-phenyl-4-(4-imidazolo)methyl-morpholine;
- 364) 2-(S)-(3-t-butyl)benzyloxy-3-(S)-(4-fluoro)phenyl-4-(4-imidazolo)methyl-morpholine;
- 10 365) 2-(R)-(1-(R)-(3-t-butyl)phenylethoxy)-3-(S)-phenyl-4-(4-imidazolo)methyl-morpholine;
- 366) 2-(R)-(1-(R)-(3-t-butyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-imidazolo)methyl-morpholine;
- 15 367) 2-(S)-(3-t-butyl)benzyloxy-3-(S)-phenyl-4-(5-tetrazolo)methyl-morpholine;
- 368) 2-(S)-(3-t-butyl)benzyloxy-3-(S)-(4-fluoro)phenyl-4-(5-tetrazolo)methyl-morpholine;
- 20 369) 2-(R)-(1-(R)-(3-t-butyl)phenylethoxy)-3-(S)-phenyl-4-(5-tetrazolo)methyl-morpholine;
- 25 370) 2-(R)-(1-(R)-(3-t-butyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(5-tetrazolo)methyl-morpholine;
- 371) 2-(S)-(3-t-butyl)benzyloxy-3-(S)-phenyl-4-(2-oxo-5H-pyrrol-4-yl)methyl-morpholine;
- 30 372) 2-(S)-(3-t-butyl)benzyloxy-3-(S)-(4-fluoro)phenyl-4-(2-oxo-5H-pyrrol-4-yl)methyl-morpholine;

- 76 -

- 373) 2-(R)-(1-(R)-(3-t-butyl)phenylethoxy)-3-(S)-phenyl-4-(2-oxo-5H-pyrrol-4-yl)methyl-morpholine;
- 5 374) 2-(R)-(1-(R)-(3-t-butyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(2-oxo-5H-pyrrol-4-yl)methyl-morpholine;
- 375) 4-(4-(imidazolo)methyl)-2-(3-(tert-butyl)-5-(trifluoromethyl)-benzyloxy)-3-phenyl-morpholine;
- 10 376) 2-(R)-(2,5-bis(trifluoromethyl)benzoyloxy)-3-(S)-(4-fluorophenyl)-4-benzyl-morpholine;
- 377) 2-(R)-(1-(2,5-bis(trifluoromethyl)phenyl)ethenyloxy)-3-(S)-(4-fluorophenyl)-4-benzyl-morpholine;
- 15 378) 2-(R)-(1-(R)-(2,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(4-fluorophenyl)-morpholine;
- 379) 2-(R)-(1-(R)-(2,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(4-fluorophenyl)-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 20 380) 2-(R)-(1-(R)-(2,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(4-fluorophenyl)-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 25 381) 2-(R)-(1-(R)-(2,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(4-fluorophenyl)-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 382) 2-(R)-(1-(R)-(3-(thiomethyl)phenyl)ethoxy)-3-(S)-phenyl-morpholine;
- 30 383) 2-(R)-(1-(R)-(3-(thiomethyl)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;

- 77 -

- 384) 2-(R)-(1-(R)-(3-(thiomethylphenyl)ethoxy)-3-(S)-phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 5 385) 2-(R)-(1-(R)-(3-(thiomethylphenyl)ethoxy)-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 386) 2-(R)-(1-(R)-(3-(thiomethyl)-5-(trifluoromethyl)phenyl)ethoxy)-3-(S)-phenyl-morpholine;
- 10 387) 2-(R)-(1-(R)-(3-(thiomethyl-5-(trifluoromethyl)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 15 388) 2-(R)-(1-(R)-(3-(thiomethyl-5-(trifluoromethyl)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 389) 2-(R)-(1-(R)-(3-(thiomethyl-5-(trifluoromethyl)phenyl)ethoxy)-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 20 390) 2-(R)-(1-(R)-(2,2-(dimethyl)-5-(thiomethyl)-2,3-dihydrobenzofuran-7-yl)ethoxy)-3-(S)-phenyl-morpholine;
- 391) 2-(R)-(1-(R)-(2,2-(dimethyl)-5-(thiomethyl)-2,3-dihydrobenzofuran-7-yl)ethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 25 392) 2-(R)-(1-(R)-(2,2-(dimethyl)-5-(thiomethyl)-2,3-dihydrobenzofuran-7-yl)ethoxy)-3-(S)-phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 30 393) 2-(R)-(1-(R)-(2,2-(dimethyl)-5-(thiomethyl)-2,3-dihydrobenzofuran-7-yl)ethoxy)-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;

- 78 -

- 394) 2-(R)-(1-(R)-(3,5-(dimethoxy)phenyl)ethoxy)-3-(S)-phenyl-morpholine;
- 5 395) 2-(R)-(1-(R)-(3,5-(dimethoxy)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 396) 2-(R)-(1-(R)-(3,5-(dimethoxy)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 10 397) 2-(R)-(1-(R)-(3,5-(dimethoxy)phenyl)ethoxy)-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 398) 2-(R)-(1-(R)-(3-(fluoro)-5-(trifluoromethyl)phenyl)-ethoxy)-3-(S)-(4-fluorophenyl)-morpholine;
- 15 399) 2-(R)-(1-(R)-(3-(fluorophenyl)-5-(trifluoromethyl)ethoxy)-3-(S)-(4-fluorophenyl)-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 20 400) 2-(R)-(1-(R)-(3-(fluorophenyl)-5-(trifluoromethyl)ethoxy)-3-(S)-(4-fluorophenyl)-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 401) 2-(R)-(1-(R)-(3-(fluorophenyl)-5-(trifluoromethyl)ethoxy)-3-(S)-(4-fluorophenyl)-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 25 402) 2-(R)-(1-(R)-(3-(chloro)-5-(trifluoromethyl)phenyl)-ethoxy)-3-(S)-(4-fluorophenyl)-morpholine;
- 403) 2-(R)-(1-(R)-(3-(chlorophenyl)-5-(trifluoromethyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 30 404) 2-(R)-(1-(R)-(3-(chlorophenyl)-5-(trifluoromethyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;

- 79 -

- 405) 2-(R)-(1-(R)-(3-(chlorophenyl)-5-(trifluoromethyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 5 406) 2-(R)-(1-(R)-(3,5-(dimethyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-morpholine;
- 407) 2-(R)-(1-(R)-(3,5-(dimethyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 10 408) 2-(R)-(1-(R)-(3,5-(dimethyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 15 409) 2-(R)-(1-(R)-(3,5-(dimethyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 410) 2-(R)-(1-(R)-(3-(fluoro)-5-(methyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-morpholine;
- 20 411) 2-(R)-(1-(R)-(3-(fluoro)-5-(methyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 25 412) 2-(R)-(1-(R)-(3-(fluoro)-5-(methyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 413) 2-(R)-(1-(R)-(3-(fluoro)-5-(methyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 30 414) 2-(R)-(1-(R)-(3-(chloro)-5-(methyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-morpholine;

- 80 -

- 415) 2-(R)-(1-(R)-(3-(chloro)-5-(methyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 5 416) 2-(R)-(1-(R)-(3-(chloro)-5-(methyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 417) 2-(R)-(1-(R)-(3-(chloro)-5-(methyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 10 418) 2-(R)-(1-(R)-(3-(bromo)-5-(methyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-morpholine;
- 419) 2-(R)-(1-(R)-(3-(bromo)-5-(methyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 15 420) 2-(R)-(1-(R)-(3-(bromo)-5-(methyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 20 421) 2-(R)-(1-(R)-(3-(bromo)-5-(methyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 422) 2-(R)-(1-(R)-(3-(isopropoxy)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-morpholine;
- 25 423) 2-(R)-(1-(R)-(3-(isopropoxy)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 30 424) 2-(R)-(1-(R)-(3-(isopropoxy)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;

- 81 -

- 425) 2-(R)-(1-(R)-(3-(isopropoxy)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 5 426) 2-(R)-(1-(R)-(3-(isopropoxy)-5-(trifluoromethyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-morpholine;
- 427) 2-(R)-(1-(R)-(3-(isopropoxy)-5-(trifluoromethyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 10 428) 2-(R)-(1-(R)-(3-(isopropoxy)-5-(trifluoromethyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 15 429) 2-(R)-(1-(R)-(3-(isopropoxy)-5-(trifluoromethyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 430) 2-(R)-(1-(R)-(3-(chloro)-5-(isopropoxy)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-morpholine;
- 20 431) 2-(R)-(1-(R)-(3-(chloro)-5-(isopropoxy)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 25 432) 2-(R)-(1-(R)-(3-(chloro)-5-(isopropoxy)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 433) 2-(R)-(1-(R)-(3-(chloro)-5-(isopropoxy)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 30 434) 2-(R)-(1-(R)-(3-(fluoro)-5-(isopropoxy)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-morpholine;

- 82 -

- 435) 2-(R)-(1-(R)-(3-(fluoro)-5-(isopropoxy)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 5 436) 2-(R)-(1-(R)-(3-(fluoro)-5-(isopropoxy)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 437) 2-(R)-(1-(R)-(3-(fluoro)-5-(isopropoxy)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 10 438) 2-(R)-(1-(R)-(3-(t-butyl)-5-(chloro)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-morpholine;
- 439) 2-(R)-(1-(R)-(3-(t-butyl)-5-(chloro)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 15 440) 2-(R)-(1-(R)-(3-(t-butyl)-5-(chloro)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 20 441) 2-(R)-(1-(R)-(3-(t-butyl)-5-(chloro)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 442) 2-(R)-(1-(R)-(3-(t-butyl)-5-(trifluoromethyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-morpholine;
- 25 443) 2-(R)-(1-(R)-(3-(t-butyl)-5-(trifluoromethyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 30 444) 2-(R)-(1-(R)-(3-(t-butyl)-5-(trifluoromethyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;

- 83 -

- 445) 2-(R)-(1-(R)-(3-(t-butyl)-5-(trifluoromethyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 5 446) 2-(R)-(1-(R)-(3,5-(dimethyl)-4-(fluoro)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-morpholine;
- 447) 2-(R)-(1-(R)-(3,5-(dimethyl)-4-(fluoro)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 10 448) 2-(R)-(1-(R)-(3,5-(dimethyl)-4-(fluoro)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 15 449) 2-(R)-(1-(R)-(3,5-(dimethyl)-4-(fluoro)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 450) 2-(R)-(1-(R)-(3,5-(dimethyl)-4-(chloro)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-morpholine;
- 20 451) 2-(R)-(1-(R)-(3,5-(dimethyl)-4-(chloro)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 25 452) 2-(R)-(1-(R)-(3,5-(dimethyl)-4-(chloro)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 453) 2-(R)-(1-(R)-(3,5-(dimethyl)-4-(chloro)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 30 454) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)-4-(fluoro)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-morpholine;

- 84 -

- 455) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)-4-(fluoro)phenyl)ethoxy)-
3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-
triazolo)methyl)-morpholine;
- 5 456) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)-4-(fluoro)phenyl)ethoxy)-
3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 457) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)-4-(fluoro)phenyl)ethoxy)-
3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-
10 morpholine;
- 458) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)-4-(chloro)phenyl)ethoxy)-
3-(S)-(4-fluoro)phenyl-morpholine;
- 15 459) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)-4-(chloro)phenyl)ethoxy)-
3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-
triazolo)methyl)-morpholine;
- 20 460) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)-4-(chloro)phenyl)ethoxy)-
3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 461) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)-4-(chloro)phenyl)ethoxy)-
3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-
25 morpholine;
- 462) 2-(R)-(1-(R)-(3,5-(dichloro)phenyl)ethoxy)-3-(S)-(4-
fluoro)phenyl-morpholine;
- 30 463) 2-(R)-(1-(R)-(3,5-(dichloro)phenyl)ethoxy)-3-(S)-(4-
fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-
morpholine;
- 464) 2-(R)-(1-(R)-(3,5-(dichloro)phenyl)ethoxy)-3-(S)-(4-
fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;

- 85 -

- 465) 2-(R)-(1-(R)-(3,5-(dichloro)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 5 466) 2-(R)-(1-(R)-(3,5-(difluoro)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-morpholine;
- 467) 2-(R)-(1-(R)-(3,5-(difluoro)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 10 468) 2-(R)-(1-(R)-(3,5-(difluoro)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 15 469) 2-(R)-(1-(R)-(3,5-(difluoro)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 470) 2-(R)-(1-(R)-(1-(naphthyl)ethoxy)-3-(S)-(4-fluoro)phenyl-morpholine;
- 20 471) 2-(R)-(1-(R)-(1-(naphthyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 472) 2-(R)-(1-(R)-(1-(naphthyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 25 473) 2-(R)-(1-(R)-(1-(naphthyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 30 474) 2-(R)-(1-(R)-(1-(4-(fluoro)naphthyl)ethoxy)-3-(S)-(4-fluoro)phenyl-morpholine;

- 86 -

- 475) 2-(R)-(1-(R)-(1-(4-(fluoro)naphthyl))ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 5 476) 2-(R)-(1-(R)-(1-(4-(fluoro)naphthyl))ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 477) 2-(R)-(1-(R)-(1-(4-(fluoro)naphthyl))ethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 10 478) 2-(R)-(1-(R)-(1-(3-(fluoro)naphthyl))ethoxy)-3-(S)-(4-fluoro)phenyl-morpholine;
- 479) 2-(R)-(1-(R)-(1-(3-(fluoro)naphthyl))ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 15 480) 2-(R)-(1-(R)-(1-(3-(fluoro)naphthyl))ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 20 481) 2-(R)-(1-(R)-(1-(3-(fluoro)naphthyl))ethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 482) 2-(R)-(1-(R)-(1-(3-(chloro)naphthyl))ethoxy)-3-(S)-(4-fluoro)phenyl-morpholine;
- 25 483) 2-(R)-(1-(R)-(1-(3-(chloro)naphthyl))ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 30 484) 2-(R)-(1-(R)-(1-(3-(chloro)naphthyl))ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;

- 87 -

- 485) 2-(R)-(1-(R)-(1-(3-(chloro)naphthyl))ethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 5 486) 2-(R)-(1-(R)-(1-(3-(methyl)naphthyl))ethoxy)-3-(S)-(4-fluoro)phenyl-morpholine;
- 487) 2-(R)-(1-(R)-(1-(3-(methyl)naphthyl))ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 10 488) 2-(R)-(1-(R)-(1-(3-(methyl)naphthyl))ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 15 489) 2-(R)-(1-(R)-(1-(3-(methyl)naphthyl))ethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 490) 2-(R)-(1-(R)-(1-(3-(trifluoromethyl)naphthyl))ethoxy)-3-(S)-(4-fluoro)phenyl-morpholine;
- 20 491) 2-(R)-(1-(R)-(1-(3-(trifluoromethyl)naphthyl))ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 25 492) 2-(R)-(1-(R)-(1-(3-(trifluoromethyl)naphthyl))ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 493) 2-(R)-(1-(R)-(1-(3-(trifluoromethyl)naphthyl))ethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 30 494) 2-(R)-(1-(R)-(3-(thiomethyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-morpholine;

- 88 -

- 495) 2-(R)-(1-(R)-(3-(thiomethylphenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 5 496) 2-(R)-(1-(R)-(3-(thiomethylphenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 497) 2-(R)-(1-(R)-(3-(thiomethylphenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 10 498) 2-(R)-(1-(R)-(3-(thiomethyl)-5-(trifluoromethyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-morpholine;
- 499) 2-(R)-(1-(R)-(3-(thiomethyl-5-(trifluoromethyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 15 500) 2-(R)-(1-(R)-(3-(thiomethyl-5-(trifluoromethyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 20 501) 2-(R)-(1-(R)-(3-(thiomethyl-5-(trifluoromethyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 25 502) 2-(R)-(1-(R)-(2,2-(dimethyl)-5-(thiomethyl)-2,3-dihydro-benzofuran-7-yl)ethoxy)-3-(S)-(4-fluoro)phenyl-morpholine;
- 503) 2-(R)-(1-(R)-(2,2-(dimethyl)-5-(thiomethyl)-2,3-dihydro-benzofuran-7-yl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 30 504) 2-(R)-(1-(R)-(2,2-(dimethyl)-5-(thiomethyl)-2,3-dihydro-benzofuran-7-yl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;

- 89 -

- 505) 2-(R)-(1-(R)-(2,2-(dimethyl)-5-(thiomethyl)-2,3-dihydro-benzofuran-7-yl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
5
- 506) 2-(R)-(1-(R)-(3,5-(dimethoxy)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-morpholine;
- 507) 2-(R)-(1-(R)-(3,5-(dimethoxy)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
10
- 508) 2-(R)-(1-(R)-(3,5-(dimethoxy)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
15
- 509) 2-(R)-(1-(R)-(3,5-(dimethoxy)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 510) 2-(R)-(1-(R)-(phenyl)ethoxy)-3-(S)-phenyl-morpholine;
20
- 511) 2-(R)-(1-(R)-(phenyl)ethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 512) 2-(R)-(1-(R)-(phenyl)ethoxy)-3-(S)-phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
25
- 513) 2-(R)-(1-(R)-(phenyl)ethoxy)-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 514) 2-(R)-(1-(R)-(phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl morpholine;
30
- 515) 2-(R)-(1-(R)-(phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;

- 90 -

- 516) 2-(R)-(1-(R)-(phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 5 517) 2-(R)-(1-(R)-(phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 518) 2-(R)-(1-(R)-(3-(fluoro)phenyl)ethoxy)-3-(S)-phenyl-morpholine;
- 10 519) 2-(R)-(1-(R)-(3-(fluoro)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 520) 2-(R)-(1-(R)-(3-(fluoro)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 15 521) 2-(R)-(1-(R)-(3-(fluoro)phenyl)ethoxy)-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 522) 2-(R)-(1-(R)-(3-(fluoro)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-morpholine;
- 20 523) 2-(R)-(1-(R)-(3-(fluoro)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 524) 2-(R)-(1-(R)-(3-(fluoro)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 25 525) 2-(R)-(1-(R)-(3-(fluoro)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 30 526) 2-(R)-(1-(R)-(4-(fluoro)phenyl)ethoxy)-3-(S)-phenyl-morpholine;
- 527) 2-(R)-(1-(R)-(4-(fluoro)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;

- 91 -

- 528) 2-(R)-(1-(R)-(4-(fluoro)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 5 529) 2-(R)-(1-(R)-(4-(fluoro)phenyl)ethoxy)-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 530) 2-(R)-(1-(R)-(4-(fluoro)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-morpholine;
- 10 531) 2-(R)-(1-(R)-(4-(fluoro)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 532) 2-(R)-(1-(R)-(4-(fluoro)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 15 533) 2-(R)-(1-(R)-(4-(fluoro)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 534) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(3-fluoro)phenyl-morpholine;
- 20 535) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(3-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 25 536) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(3-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 537) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(3-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 30 538) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(3,4-difluoro)phenyl-morpholine;

- 92 -

- 539) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(3,4-difluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 5 540) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(3,4-difluoro)phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 541) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(3,4-difluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 10 542) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(3,4-dichloro)phenyl-morpholine;
- 543) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(3,4-dichloro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 15 544) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(3,4-dichloro)phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 20 545) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(3,4-dichloro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 546) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(3,4-dimethyl)phenyl-morpholine;
- 25 547) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(3,4-dimethyl)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 30 548) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(3,4-dimethyl)phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;

- 93 -

- 549) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(3,4-dimethyl)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 5 550) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-3,4-methylenedioxyphenyl-morpholine;
- 551) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-3,4-methylenedioxyphenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 10 552) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-3,4-methylenedioxyphenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 15 553) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-3,4-methylenedioxyphenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 554) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(2-naphthyl)-morpholine;
- 20 555) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(2-naphthyl)-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 25 556) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(2-naphthyl)-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 557) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(2-naphthyl)-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 30 558) 2-(R)-(1-(R)-(3-(fluorophenyl)-5-(trifluoromethyl)ethoxy)-3-(S)-phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 559) 2-(R)-(1-(R)-(3-(fluorophenyl)-5-(trifluoromethyl)ethoxy)-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;

- 94 -

- 560) 2-(R)-(1-(R)-(3-(chlorophenyl)-5-(trifluoromethyl)ethoxy)-3-(S)-phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 5 561) 2-(R)-(1-(R)-(3-(chlorophenyl)-5-(trifluoromethyl)ethoxy)-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 562) 2-(R)-(1-(R)-(3,5-(dimethyl)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 10 563) 2-(R)-(1-(R)-(3,5-(dimethyl)phenyl)ethoxy)-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 564) 2-(R)-(1-(R)-(3-(fluoro)-5-(methyl)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 15 565) 2-(R)-(1-(R)-(3-(fluoro)-5-(methyl)phenyl)ethoxy)-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 20 566) 2-(R)-(1-(R)-(3-(chloro)-5-(methyl)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 567) 2-(R)-(1-(R)-(3-(chloro)-5-(methyl)phenyl)ethoxy)-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 25 568) 2-(R)-(1-(R)-(3-(bromo)-5-(methyl)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 569) 2-(R)-(1-(R)-(3-(bromo)-5-(methyl)phenyl)ethoxy)-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 30 570) 2-(R)-(1-(R)-(3-(isopropoxy)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;

- 95 -

- 571) 2-(R)-(1-(R)-(3-(isopropoxy)phenyl)ethoxy)-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 5 572) 2-(R)-(1-(R)-(3-(isopropoxy)-5-(trifluoromethyl)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 573) 2-(R)-(1-(R)-(3-(isopropoxy)-5-(trifluoromethyl)phenyl)ethoxy)-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 10 574) 2-(R)-(1-(R)-(3-(chloro)-5-(isopropoxy)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 575) 2-(R)-(1-(R)-(3-(chloro)-5-(isopropoxy)phenyl)ethoxy)-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 15 576) 2-(R)-(1-(R)-(3-(fluoro)-5-(isopropoxy)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 577) 2-(R)-(1-(R)-(3-(fluoro)-5-(isopropoxy)phenyl)ethoxy)-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 20 578) 2-(R)-(1-(R)-(3-(t-butyl)-5-(chloro)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 25 579) 2-(R)-(1-(R)-(3-(t-butyl)-5-(chloro)phenyl)ethoxy)-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 580) 2-(R)-(1-(R)-(3-(t-butyl)-5-(trifluoromethyl)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 30 581) 2-(R)-(1-(R)-(3-(t-butyl)-5-(trifluoromethyl)phenyl)ethoxy)-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;

- 96 -

- 582) 2-(R)-(1-(R)-(3,5-(dimethyl)-4-(fluoro)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 583) 2-(R)-(1-(R)-(3,5-(dimethyl)-4-(fluoro)phenyl)ethoxy)-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 584) 2-(R)-(1-(R)-(3,5-(dimethyl)-4-(chloro)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 585) 2-(R)-(1-(R)-(3,5-(dimethyl)-4-(chloro)phenyl)ethoxy)-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 586) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)-4-(fluoro)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 587) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)-4-(fluoro)phenyl)ethoxy)-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 588) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)-4-(chloro)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 589) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)-4-(chloro)phenyl)ethoxy)-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 590) 2-(R)-(1-(R)-(3,5-(dichloro)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 591) 2-(R)-(1-(R)-(3,5-(dichloro)phenyl)ethoxy)-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 592) 2-(R)-(1-(R)-(3,5-(difluoro)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;

- 97 -


- 593) 2-(R)-(1-(R)-(3,5-(difluoro)phenyl)ethoxy)-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 5 594) 2-(R)-(1-(R)-(1-(naphthyl)ethoxy)-3-(S)-phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 595) 2-(R)-(1-(R)-(1-(naphthyl)ethoxy)-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 10 596) 2-(R)-(1-(R)-(1-(4-(fluoro)naphthyl))ethoxy)-3-(S)-phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 597) 2-(R)-(1-(R)-(1-(4-(fluoro)naphthyl))ethoxy)-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 15 598) 2-(R)-(1-(R)-(1-(3-(fluoro)naphthyl))ethoxy)-3-(S)-phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 599) 2-(R)-(1-(R)-(1-(3-(fluoro)naphthyl))ethoxy)-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 20 600) 2-(R)-(1-(R)-(1-(3-(chloro)naphthyl))ethoxy)-3-(S)-phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 25 601) 2-(R)-(1-(R)-(1-(3-(chloro)naphthyl))ethoxy)-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;

and pharmaceutically acceptable salts thereof.


30

- 98 -

Representative examples of the nomenclature employed herein are given below:




- 15 439) 2-(R)-(1-(R)-(3-(t-butyl)-5-(chloro)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;



- 30 449) 2-(R)-(1-(R)-(3,5-(dimethyl)-4-(fluoro)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;

- 99 -



468) 2-(R)-(1-(R)-(3,5-(difluoro)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;

15

TACHYKININ ANTAGONISM ASSAY

The compounds of this invention are useful for antagonizing tachykinins, in particular substance P and neurokinin A in the treatment of gastrointestinal disorders, central nervous system disorders, inflammatory diseases, pain or migraine and asthma in a mammal in need of such treatment. This activity can be demonstrated by the following assays.

A. Receptor Expression in COS

25 To express the cloned human neurokinin-1 receptor (NK1R) transiently in COS, the cDNA for the human NK1R was cloned into the expression vector pCDM9 which was derived from pCDM8 (INVITROGEN) by inserting the ampicillin resistance gene (nucleotide 1973 to 2964 from BLUESCRIPT SK+) into the Sac II site.

30 Transfection of 20 ug of the plasmid DNA into 10 million COS cells was achieved by electroporation in 800 ul of transfection buffer (135 mM NaCl, 1.2 mM CaCl₂, 1.2 mM MgCl₂, 2.4 mM K₂HPO₄, 0.6 mM KH₂PO₄, 10 mM glucose, 10 mM HEPES pH 7.4) at 260 V and 950 uF using the IBI GENEZAPPER (IBI, New Haven, CT). The cells were incubated in 10% fetal calf serum, 2 mM glutamine, 100U/ml penicillin-

- 100 -

streptomycin, and 90% DMEM media (GIBCO, Grand Island, NY) in 5% CO₂ at 37°C for three days before the binding assay.

B. Stable Expression in CHO

5 To establish a stable cell line expressing the cloned human NK1R, the cDNA was subcloned into the vector pRcCMV (INVITROGEN). Transfection of 20 ug of the plasmid DNA into CHO cells was achieved by electroporation in 800 ul of transfection buffer supplemented with 0.625 mg/ml Herring sperm DNA at 300 V and 950
10 uF using the IBI GENEZAPPER (IBI). The transfected cells were incubated in CHO media [10 % fetal calf serum, 100 U/ml penicillin-streptomycin, 2 mM glutamine, 1/500 hypoxanthine-thymidine (ATCC), 90% IMDM media (JRH BIOSCIENCES, Lenexa, KS), 0.7 mg/ml G418 (GIBCO)] in 5% CO₂ at 37°C until colonies were visible. Each colony
15 was separated and propagated. The cell clone with the highest number of human NK1R was selected for subsequent applications such as drug screening.

C. Assay Protocol using COS or CHO

20 The binding assay of human NK1R expressed in either COS or CHO cells is based on the use of ¹²⁵I-substance P (¹²⁵I-SP, from DU PONT, Boston, MA) as a radioactively labeled ligand which competes with unlabeled substance P or any other ligand for binding to the human NK1R. Monolayer cell cultures of COS or CHO were
25 dissociated by the non-enzymatic solution (SPECIALTY MEDIA, Lavallette, NJ) and resuspended in appropriate volume of the binding buffer (50 mM Tris pH 7.5, 5 mM MnCl₂, 150 mM NaCl, 0.04 mg/ml bacitracin, 0.004 mg/ml leupeptin, 0.2 mg/ml BSA, 0.01 mM phosphoramidon) such that 200 ul of the cell suspension would give rise
30 to about 10,000 cpm of specific ¹²⁵I-SP binding (approximately 50,000 to 200,000 cells). In the binding assay, 200 ul of cells were added to a tube containing 20 ul of 1.5 to 2.5 nM of ¹²⁵I-SP and 20 ul of unlabeled substance P or any other test compound. The tubes were incubated at 4°C or at room temperature for 1 hour with gentle shaking.

- 101 -

The bound radioactivity was separated from unbound radioactivity by GF/C filter (BRANDEL, Gaithersburg, MD) which was pre-wetted with 0.1 % polyethylenimine. The filter was washed with 3 ml of wash buffer (50 mM Tris pH 7.5, 5 mM MnCl₂, 150 mM NaCl) three times
5 and its radioactivity was determined by gamma counter.

The activation of phospholipase C by NK1R may also be measured in CHO cells expressing the human NK1R by determining the accumulation of inositol monophosphate which is a degradation product of IP₃. CHO cells are seeded in 12-well plate at 250,000 cells per well.
10 After incubating in CHO media for 4 days, cells are loaded with 0.025 uCi/ml of ³H-myoinositol by overnight incubation. The extracellular radioactivity is removed by washing with phosphate buffered saline. LiCl is added to the well at final concentration of 0.1 mM with or without the test compound, and incubation is continued at 37°C for 15
15 min. Substance P is added to the well at final concentration of 0.3 nM to activate the human NK1R. After 30 min of incubation at 37°C, the media is removed and 0.1 N HCl is added. Each well is sonicated at 4°C and extracted with CHCl₃/methanol (1:1). The aqueous phase is applied to a 1 ml Dowex AG 1X8 ion exchange column. The column is washed
20 with 0.1 N formic acid followed by 0.025 M ammonium formate-0.1 N formic acid. The inositol monophosphate is eluted with 0.2 M ammonium formate-0.1 N formic acid and quantitated by beta counter.

The activity of the present compounds may also be demonstrated by the assay disclosed by Lei, et al., British J. Pharmacol.,
25 105, 261-262 (1992).

The compounds of the present invention are useful in the prevention and treatment of a wide variety of clinical conditions which are characterized by the presence of an excess of tachykinin, in particular substance P, activity. These conditions may include disorders
30 of the central nervous system such as anxiety, depression, psychosis and schizophrenia; epilepsy; neurodegenerative disorders such as dementia, including senile dementia of the Alzheimer type, Alzheimer's disease and Down's syndrome; demyelinating diseases such as multiple sclerosis (MS) and amyotrophic lateral sclerosis (ALS; Lou Gehrig's disease) and

- 102 -

other neuropathological disorders such as peripheral neuropathy, for example AIDS related neuropathy, diabetic neuropathy, chemotherapy-induced neuropathy, and postherpetic and other neuralgias; small cell carcinomas such as small cell lung cancer; respiratory diseases, particularly those associated with excess mucus secretion, such as chronic obstructive airways disease, bronchopneumonia, chronic bronchitis, acute bronchitis, diffuse panbronchitis, emphysema, cystic fibrosis, asthma, and bronchospasm; airways disease modulated by neurogenic inflammation; laryngopharyngitis; bronchiectasis; whooping cough; pulmonary tuberculosis; diseases associated with decreased glandular secretions, including lacrimation, such as Sjogren's syndrome, hyperlipoproteinemias IV and V, hemochromatosis, sarcoidosis, or amyloidosis; iritis; inflammatory diseases such as inflammatory bowel disease, inflammatory intestinal disease, psoriasis, fibrositis, ocular inflammation, osteoarthritis, rheumatoid arthritis, pruritis, and sunburn; hepatitis; allergies such as eczema and rhinitis; hypersensitivity disorders such as poison ivy; ophthalmic diseases such as conjunctivitis, vernal conjunctivitis, dry eye syndrome, and the like; ophthalmic conditions associated with cell proliferation such as proliferative vitreoretinopathy; cutaneous diseases such as contact dermatitis, atopic dermatitis, urticaria, and other eczematoid dermatitis; hemodialysis-associated itching; lichen planus; oedema, such as oedema caused by thermal injury; addiction disorders such as alcoholism; mental disease, particularly anxiety and depression; stress related somatic disorders; reflex sympathetic dystrophy such as shoulder/hand syndrome; dysthymic disorders; tenalgia attended to hyperlipidemia; postoperative neuroma, particularly of mastectomy; vulvar vestibulitis; adverse immunological reactions such as rejection of transplanted tissues and disorders related to immune enhancement or suppression, such as systemic lupus erythematosus; gastrointestinal (GI) disorders and diseases of the GI tract such as disorders associated with the neuronal control of viscera such as ulcerative colitis, Crohn's disease, irritable bowel syndrome, nausea, and emesis, including acute, delayed, post-operative, late-phase, and anticipatory emesis, such as emesis or nausea induced by

- 103 -

for example chemotherapy, radiation, surgery, migraine, toxins, such as metabolic or microbial toxins, viral or bacterial infections, pregnancy, vestibular disorder, motion, mechanical stimulation, gastrointestinal obstruction, reduced gastrointestinal motility, visceral pain,
5 psychological stress or disturbance, high altitude, weightlessness, opioid analgesics, intoxication, resulting for example from consumption of alcohol, and variations in intracranial pressure, in particular, for example, drug or radiation induced emesis or post-operative nausea and vomiting; disorders of bladder function such as cystitis, bladder
10 detrusor hyperreflexia, and incontinence; fibrosing and collagen diseases such as scleroderma and eosinophilic fasciitis; disorders of blood flow caused by vasodilation and vasospastic diseases such as angina, migraine and Reynaud's disease; and pain or nociception, for example, chronic pain or that attributable to or associated with any of
15 the foregoing conditions especially the transmission of pain in migraine, or such as headache, toothache, cancerous pain, back pain, and superficial pain on congelation, burn, herpes zoster or diabetic neuropathy. Hence, these compounds may be readily adapted to therapeutic use for the treatment of physiological disorders associated
20 with an excessive stimulation of tachykinin receptors, especially neurokinin-1, and as neurokinin-1 antagonists in the control and/or treatment of any of the aforesaid clinical conditions in mammals, including humans.

25 The compounds of the present invention are also of value in the treatment of a combination of the above conditions, in particular in the treatment of combined post-operative pain and post-operative nausea and vomiting.

The compounds of the present invention are particularly
30 useful in the treatment of nausea or emesis, including acute, delayed, post-operative, late-phase, and anticipatory emesis, such as emesis or nausea induced by for example chemotherapy, radiation, surgery, migraine, toxins, such as metabolic or microbial toxins, viral or bacterial infections, pregnancy, vestibular disorder, motion, mechanical stimulation, gastrointestinal obstruction, reduced gastrointestinal

- 104 -

motility, visceral pain, psychological stress or disturbance, high altitude, weightlessness, opioid analgesics, intoxication, resulting for example from consumption of alcohol, and variations in intracranial pressure. Most especially, this compound is of use in the treatment of emesis induced by antineoplastic (cytotoxic) agents including those routinely used in cancer chemotherapy.

Examples of such chemotherapeutic agents include alkylating agents, for example, nitrogen mustards, ethyleneimine compounds, alkyl sulfonates and other compounds with an alkylating action such as nitrosoureas, cisplatin, and dacarbazine; antimetabolites, for example, folic acid, purine or pyrimidine antagonists; mitotic inhibitors, for example, vinca alkaloids and derivatives of podophyllotoxin; and cytotoxic antibiotics.

Particular examples of chemotherapeutic agents are described, for example, by D. J. Stewart in "Nausea and Vomiting: Recent Research and Clinical Advances", Eds. J. Kucharczyk, *et al.*, CRC Press Inc., Boca Raton, Florida, USA (1991), pages 177-203, especially page 188. Commonly used chemotherapeutic agents include cisplatin, dacarbazine (DTIC), dactinomycin, mechlorethamine (nitrogen mustard), streptozocin, cyclophosphamide, carmustine (BCNU), lomustine (CCNU), doxorubicin (adriamycin), daunorubicin, procarbazine, mitomycin, cytarabine, etoposide, methotrexate, 5-fluorouracil, vinblastine, vincristine, bleomycin, and chlorambucil [R. J. Gralla, *et al.*, Cancer Treatment Reports, 68(1), 163-172 (1984)].

The compounds of the present invention are also of use in the treatment of emesis induced by radiation including radiation therapy such as in the treatment of cancer, or radiation sickness, and in the treatment of post-operative nausea and vomiting.

The compounds of the present invention are also of use in the prevention or treatment of disorders of the central nervous system such as anxiety, psychosis and schizophrenia; neurodegenerative disorders such as senile dementia of the Alzheimer type, Alzheimer's disease and Down's syndrome; respiratory diseases, particularly those associated with excess mucus secretion, such as chronic obstructive

- 105 -

airways disease, broncho-pneumonia, chronic bronchitis, cystic fibrosis and asthma, and bronchospasm; inflammatory diseases such as inflammatory bowel disease, osteoarthritis and rheumatoid arthritis; adverse immunological reactions such as rejection of transplanted
5 tissues; gastrointestinal (GI) disorders and diseases of the GI tract such as disorders associated with the neuronal control of viscera such as ulcerative colitis, Crohn's disease and incontinence; disorders of blood flow caused by vasodilation; and pain or nociception, for example, that attributable to or associated with any of the foregoing conditions or the
10 transmission of pain in migraine (both prophylaxis and acute treatment).

As calcium channel blocking agents the compounds of the present invention are useful in the prevention of treatment of clinical conditions which benefit from inhibition of the transfer of calcium ions across the plasma membrane of cells. These include diseases and
15 disorders of the heart and vascular system such as angina pectoris, myocardial infarction, cardiac arrhythmia, cardiac hypertrophy, cardiac vasospasm, hypertension, cerebrovascular spasm and other ischemic disease. Furthermore, these compounds may be capable of lowering elevated intraocular pressure when administered topically to the
20 hypertensive eye in solution in a suitable ophthalmic vehicle. Also, these compounds may be useful in the reversal of multidrug resistance in tumor cells by enhancing the efficacy of chemotherapeutic agents. In addition, the compounds may have activity in blocking calcium channels in insect brain membranes and so may be useful as insecticides.

25 The compounds of the present invention are particularly useful in the treatment of pain or nociception and/or inflammation and disorders associated therewith such as, for example: neuropathy, such as diabetic or peripheral neuropathy and chemotherapy-induced neuropathy; postherpetic and other neuralgias; asthma; osteoarthritis;
30 rheumatoid arthritis; and especially migraine. The compounds of the present invention are also particularly useful in the treatment of diseases characterized by neurogenic mucus secretion, especially cystic fibrosis.

In the treatment of the clinical conditions noted above, the compounds of this invention may be utilized in compositions such as

- 106 -

tablets, capsules or elixirs for oral administration, suppositories for rectal administration, sterile solutions or suspensions for parenteral or intramuscular administration, and the like.

5 The pharmaceutical compositions of this invention may be used in the form of a pharmaceutical preparation, for example, in solid, semisolid or liquid form, which contains one or more of the compounds of the present invention, as an active ingredient, in admixture with an organic or inorganic carrier or excipient suitable for external, enteral or parenteral applications. The active ingredient may be compounded,
10 for example, with the usual non-toxic, pharmaceutically acceptable carriers for tablets, pellets, capsules, suppositories, solutions, emulsions, suspensions, and any other form suitable for use. The carriers which can be used are water, glucose, lactose, gum acacia, gelatin, mannitol, starch paste, magnesium trisilicate, talc, corn starch, keratin, colloidal
15 silica, potato starch, urea and other carriers suitable for use in manufacturing preparations, in solid, semisolid, or liquid form, and in addition auxiliary, stabilizing, thickening and coloring agents and perfumes may be used. The active object compound is included in the pharmaceutical composition in an amount sufficient to produce the
20 desired effect upon the process or condition of the disease.

For preparing solid compositions such as tablets, the principal active ingredient is mixed with a pharmaceutical carrier, e.g. conventional tableting ingredients such as corn starch, lactose, sucrose, sorbitol, talc, stearic acid, magnesium stearate, dicalcium phosphate or
25 gums, and other pharmaceutical diluents, e.g. water, to form a solid preformulation composition containing a homogeneous mixture of a compound of the present invention, or a non-toxic pharmaceutically acceptable salt thereof. When referring to these preformulation compositions as homogeneous, it is meant that the active ingredient is
30 dispersed evenly throughout the composition so that the composition may be readily subdivided into equally effective unit dosage forms such as tablets, pills and capsules. This solid preformulation composition is then subdivided into unit dosage forms of the type described above containing from 0.1 to about 500 mg of the active ingredient of the

- 107 -

present invention. The tablets or pills of the novel composition can be coated or otherwise compounded to provide a dosage form affording the advantage of prolonged action. For example, the tablet or pill can comprise an inner dosage and an outer dosage component, the latter
5 being in the form of an envelope over the former. The two components can be separated by an enteric layer which serves to resist disintegration in the stomach and permits the inner component to pass intact into the duodenum or to be delayed in release. A variety of materials can be used for such enteric layers or coatings, such materials including a
10 number of polymeric acids and mixtures of polymeric acids with such materials as shellac, cetyl alcohol and cellulose acetate.

The liquid forms in which the novel compositions of the present invention may be incorporated for administration orally or by injection include aqueous solution, suitably flavoured syrups,
15 aqueous or oil suspensions, and emulsions with acceptable oils such as cottonseed oil, sesame oil, coconut oil or peanut oil, or with a solubilizing or emulsifying agent suitable for intravenous use, as well as elixirs and similar pharmaceutical vehicles. Suitable dispersing or suspending agents for aqueous suspensions include synthetic and
20 natural gums such as tragacanth, acacia, alginate, dextran, sodium carboxymethylcellulose, methylcellulose, polyvinylpyrrolidone or gelatin.

Compositions for inhalation or insufflation include solutions and suspensions in pharmaceutically acceptable, aqueous or
25 organic solvents, or mixtures thereof, and powders. The liquid or solid compositions may contain suitable pharmaceutically acceptable excipients as set out above. Preferably the compositions are administered by the oral or nasal respiratory route for local or systemic effect. Compositions in preferably sterile pharmaceutically acceptable
30 solvents may be nebulized by use of inert gases. Nebulized solutions may be breathed directly from the nebulizing device or the nebulizing device may be attached to a face mask, tent or intermittent positive pressure breathing machine. Solution, suspension or powder

- 108 -

compositions may be administered, preferably orally or nasally, from devices which deliver the formulation in an appropriate manner.

For the treatment of the clinical conditions and diseases noted above, the compounds of this invention may be administered
5 orally, topically, parenterally, by inhalation spray or rectally in dosage unit formulations containing conventional non-toxic pharmaceutically acceptable carriers, adjuvants and vehicles. The term parenteral as used herein includes subcutaneous injections, intravenous, intramuscular, intrasternal injection or infusion techniques.

10 For the treatment of certain conditions it may be desirable to employ a compound of the present invention in conjunction with another pharmacologically active agent. For example, a compound of the present invention may be presented together with another
15 therapeutic agent as a combined preparation for simultaneous, separate, or sequential use for the relief of emesis. Such combined preparations may be, for example, in the form of a twin pack. A preferred combination comprises a compound of the present invention with a chemotherapeutic agent such as an alkylating agent, antimetabolite, mitotic inhibitor, or cytotoxic antibiotic, as described above. In
20 general, the currently available dosage forms of the known therapeutic agents for use in such combinations will be suitable.

Similarly, for the treatment of respiratory diseases, such as asthma, a compound of the present invention may be used in conjunction with a bronchodilator, such as a β_2 -adrenergic receptor agonist or a
25 tachykinin antagonist which acts at neurokinin-2 receptors. Also, for the treatment of conditions that require antagonism of both neurokinin-1 and neurokinin-2, including disorders associated with bronchoconstriction and/or plasma extravasation in airways, such as asthma, chronic bronchitis, airways disease, or cystic fibrosis, a
30 compound of the present invention may be used in conjunction with a tachykinin antagonist which acts at neurokinin-2 receptors, or with tachykinin receptor antagonist which acts at both neurokinin-1 and neurokinin-2 receptors. Similarly, for the prevention or treatment of emesis a compound of the present invention may be used in conjunction

- 109 -

with other anti-emetic agents, especially 5HT₃ receptor antagonists, such as ondansetron, granisetron, tropisetron, decadron, and zatisetron. Likewise, for the prevention or treatment of migraine a compound of the present invention may be used in conjunction with other anti-migraine agents, such as ergotamines or 5HT₁ agonists, especially sumatriptan. Likewise, for the treatment of behavioral hyperalgesia, a compound of the present invention may be used in conjunction with an antagonist of N-methyl D-aspartate (NMDA), such as dizocilpine. For the prevention or treatment of inflammatory conditions in the lower urinary tract, especially cystitis, a compound of the present invention may be used in conjunction with an antiinflammatory agent, such as a bradykinin receptor antagonist. The compound of the present invention and the other pharmacologically active agent may be administered to a patient simultaneously, sequentially or in combination.

The compounds of this invention may be administered to patients (animals and humans) in need of such treatment in dosages that will provide optimal pharmaceutical efficacy. It will be appreciated that the dose required for use in any particular application will vary from patient to patient, not only with the particular compound or composition selected, but also with the route of administration, the nature of the condition being treated, the age and condition of the patient, concurrent medication or special diets then being followed by the patient, and other factors which those skilled in the art will recognize, with the appropriate dosage ultimately being at the discretion of the attendant physician.

In the treatment of a condition associated with an excess of tachykinins, an appropriate dosage level will generally be about 0.001 to 50 mg per kg patient body weight per day which may be administered in single or multiple doses. Preferably, the dosage level will be about 0.01 to about 25 mg/kg per day; more preferably about 0.05 to about 10 mg/kg per day. For example, in the treatment of conditions involving the neurotransmission of pain sensations, a suitable dosage level is about 0.001 to 25 mg/kg per day, preferably about 0.05 to 10 mg/kg per day, and especially about 0.1 to 5 mg/kg per day. A compound may be

- 110 -

administered on a regimen of 1 to 4 times per day, preferably once or twice per day. In the treatment of emesis using an injectable formulation, a suitable dosage level is about 0.001 to 10 mg/kg per day, preferably about 0.005 to 5 mg/kg per day, and especially about 0.05 to 5 mg/kg per day. A compound may be administered on a regimen of 1 to 4 times per day, preferably once or twice per day.

Several methods for preparing the compounds of this invention are illustrated in the following Schemes and Examples wherein R1, R2, R3, R4, R5, R6, R7, R8, R9, R10, R11, R12 and R13 are as defined above.

ABBREVIATIONS USED IN SCHEMES AND EXAMPLES

Table 1

15 Reagents:

Et ₃ N	triethylamine
Ph ₃ P	triphenylphosphine
TFA	trifluoroacetic acid
20 NaOEt	sodium ethoxide
DCC	N,N'-dicyclohexylcarbodiimide
DCU	N,N'-dicyclohexylurea
CDI	1,1'-carbonyldiimidazole
MCPBA	m-chloroperbenzoic acid
25 DBU	1,8-diazabicyclo[5.4.0]undec-7-ene
Cbz-Cl	benzyl chloroformate
ACE-Cl	alpha-chloroethyl chloroformate
iPr ₂ NEt or DIEA	N,N-diisopropylethylamine
NHS	N-hydroxysuccinimide
30 DIBAL	diisobutylaluminum hydride
Me ₂ SO ₄	dimethyl sulfate
HOBt	1-hydroxybenzotriazole hydrate
EDAC	1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride

- 111 -

Solvents:


	DMF	dimethylformamide
	THF	tetrahydrofuran
5	MeOH	methanol
	EtOH	ethanol
	AmOH	n-amyl alcohol
	AcOH	acetic acid
	MeCN	acetonitrile
10	DMSO	dimethylsulfoxide

Others:


	Ph	phenyl
15	Ar	aryl
	Me	methyl
	Et	ethyl
	iPr	isopropyl
	Am	n-amyl
20	Cbz	carbobenzyloxy (benzyloxy-carbonyl)
	BOC	<u>tert</u> -butoxycarbonyl
	PTC	phase transfer catalyst
	cat.	catalytic
	FAB-MS	fast atom bombardment mass spectrometry
25	rt	room temperature

30

- 112 -


SCHEME 1

- 113 -


SCHEME 2

- 114 -

SCHEME 3




- 115 -

SCHEME 4


- 116 -

SCHEME 5




- 117 -


SCHEME 6




- 118 -

SCHEME 7


- 119 -

SCHEME 8



- 120 -

SCHEME 8 (CONT'D)

- 121 -

SCHEME 9

- 122 -

SCHEME 9 (CONT'D) $S = H_2 \text{ or } R^{14}-CH_2-$  $S = H_2 \text{ or } R^{14}-CH_2-$

25

30

- 123 -

The compounds of the present invention in which $X = Y = O$ may be prepared by the general route outlined in Scheme 1. Thus, the appropriately substituted α -bromo-phenylacetaldehyde, dimethyl acetal I (prepared using the method of Jacobs in Journal of the American Chemical Society, 1953, 75, 5500) may be converted to the dibenzyl acetal II by stirring I and a slight excess of a benzyl alcohol in the presence of an acid catalyst with concomitant removal of methanol. Alkylation of a substituted amino alcohol by benzyl bromide II may give N-alkyl amino alcohol III; use of a chiral amino alcohol would result in the formation of diastereomers and these may be separated at this (or at a later) stage using standard chromatographic methods. N-Alkylation or N-acylation of III may give the dialkyl- or acyl/alkyl-amino alcohol IV in which the group R^1 may serve as a protecting group or be used as or elaborated into a substituent in the final target compound. Cyclization to give substituted morpholine V may be realized by warming a solution of IV and an acid catalyst.

Diastereomers of V that may be formed may be separated using standard chromatographic methods. If R^1 is a protecting group, it may be removed using known procedures (Greene, T.W., Wuts, P.G.M. Protective Groups in Organic Synthesis, 2nd ed., John Wiley & Sons, Inc., New York, 1991). If the preparation of I-V results in the formation of enantiomers, these may be resolved by alkylating or acylating V ($R^1 = H$) with a chiral auxiliary, separating the diastereomers thus formed using known chromatographic methods, and removing the chiral auxiliary to give the enantiomers of V. Alternatively, the diastereomers of V may be separated via fractional crystallization from a suitable solvent of the diastereomeric salts formed by V and a chiral organic acid.

The compounds of the present invention in which $X = O$ and $Y = CH_2$ may be prepared by the general route outlined in Scheme 2. Thus, the N-methoxy-N-methyl amide of a protected phenyl glycine VI (prepared from the carboxylic acid via the mixed anhydride according to the procedure of Rapoport in Journal of Organic Chemistry, 1985, 50, 3972) may be used to acylate the lithium enolate

- 124 -

of methyl diethylphosphonate to give the ketophosphonate VII. The sodium salt of VII may be condensed with an appropriately substituted benzaldehyde to give the α,β -unsaturated ketone VIII. Reduction of the ketone and removal of the t-butylcarbamate protecting group may give amino alcohol IX; diastereomers that may form may be separated at this (or at a later) stage using standard chromatographic techniques. Williamson etherification of IX using a substituted chloroacetate, followed by warming, may result in the formation of morpholinone X. Reduction of the double bond and amide carbonyl may be accomplished in a straightforward manner to give the substituted morpholine XI. If the preparation of VI-XI results in the formation of enantiomers, these may be resolved by alkylating or acylating XI ($R^1 = H$) with a chiral auxiliary, separating the diastereomers thus formed using known chromatographic methods, and removing the chiral auxiliary to give the enantiomers of XI. Alternatively, the diastereomers of XI may be separated via fractional crystallization from a suitable solvent of the diastereomeric salts formed by XI and a chiral organic acid. If it is desired that R^1 is other than H, the morpholine nitrogen of XI may be further functionalized using standard methods for the alkylation or acylation of secondary amines. If it is desired that R^2 is other than H, morpholinone X may be elaborated into the carbinolcarbamate ($R^1 = RO_2C$, $R^2 = OH$), an intermediate that could be alkylated and would allow for variation in R^2 .

The compounds of the present invention in which $X = S-(O)_n$ ($n = 0,1,2$) and $Y = O$ may be prepared by the general route outlined in Scheme 3. Thus, alcohol IV (prepared in Scheme 1) may be converted to thioacetate XII using known procedures (Volante, R.P. Tetrahedron Letters, 1981, 22, 3119). Cleavage of the ester moiety to afford thiol XIII may be effected with aqueous base or reductively, depending on the restraints imposed by the other functional groups present. Cyclization of XIII to thiomorpholine XIV may be done by warming a solution of XIII and an acid catalyst. Oxidation of XIV using sodium metaperiodate in acetic acid may afford sulfoxide or sulfone XV. Diastereomers of XIV or XV that may be formed may be

- 125 -

separated using standard chromatographic methods. If R¹ is a protecting group, it may be removed using known procedures (Greene, T.W., Wuts, P.G.M. *Protective Groups in Organic Synthesis*, 2nd ed., John Wiley & Sons, Inc., New York, 1991). If the preparation of XII -
5 XV results in the formation of enantiomers, these may be resolved by alkylating or acylating XIV or XV (R¹ = H) with a chiral auxiliary, separating the diastereomers thus formed using known chromatographic methods, and removing the chiral auxiliary to give the enantiomers of XIV or XV. Alternatively, the diastereomers of XIV or XV may be
10 separated via fractional crystallization from a suitable solvent of the diastereomeric salts formed by XIV or XV and a chiral organic acid.

The compounds of the present invention in which X = Y = O may also be prepared by the general route outlined in Scheme 4. Thus, the appropriately substituted α -bromo-acetaldehyde, dimethyl
15 acetal (prepared using the method of Jacobs in Journal of the American Chemical Society, 1953, 75, 5500) may be converted to the acetal by stirring and a slight excess of the appropriate alcohol in the presence of an acid catalyst with concomitant removal of methanol. Alkylation of
20 a substituted amino alcohol by a bromide may give the N-alkyl amino alcohol; use of a chiral amino alcohol would result in the formation of diastereomers and these may be separated at this (or at a later) stage using standard chromatographic methods. N-Alkylation or N-acylation may give the dialkyl- or acyl/alkyl-amino alcohol in which the group R¹
25 may serve as a protecting group or be used as or elaborated into a substituent in the final target compound. Cyclization to give substituted morpholine may be realized by warming a solution with an acid catalyst. Diastereomers that may be formed may be separated using standard chromatographic methods. If R¹ is a protecting group, it may
30 be removed using known procedures (Greene, T.W., Wuts, P.G.M. *Protective Groups in Organic Synthesis*, 2nd ed., John Wiley & Sons, Inc., New York, 1991). If the preparation of such compounds results in the formation of enantiomers, these may be resolved by alkylating or acylating the final product (R¹ = H) with a chiral auxiliary, separating the diastereomers thus formed using known chromatographic methods,

- 126 -

and removing the chiral auxiliary to give the desired enantiomers. Alternatively, the diastereomers may be separated via fractional crystallization from a suitable solvent of the diastereomeric salts formed by the compound of a chiral organic acid.

5 One method of synthesizing enantiomerically pure substituted morpholines is illustrated in Scheme 5. Protection of enantiomerically pure phenylglycine as the N-benzyl derivative followed by double alkylation with a 1,2-dibromoethane derivative leads to the morpholinone. Reduction with an active hydride reagent such as
10 diisobutyl aluminum hydride, lithium aluminum hydride, lithium tri(sec-butyl)-borohydride (L-Selectride®) or other reducing agents leads predominantly to the 2,3-trans morpholine derivatives. Alkylation of the alcohol, removal of the protecting group on nitrogen (for example, with a palladium hydrogenation catalyst or with 1-chloroethyl
15 chloroformate (Olofson in *J. Org. Chem.*, 1984, 2081 and 2795), and alkylation of the nitrogen (wherein R^1CH_2- or R^1CHO = appropriate definitions of R^1 , and LG is an appropriate leaving group) produces the 2,3-trans compounds.

20 One method of producing enantiomerically pure 2,3-cis morpholines is illustrated in Scheme 6. In the first step, formation of the trifluoromethane-sulfonate ester of the appropriate benzyl alcohol (especially benzyl alcohols which are substituted with electron-withdrawing groups such as $-NO_2$, $-F$, $-Cl$, $-Br$, $-COR$, $-CF_3$, etc) is carried out in the presence of an unreactive base, in an inert solvent.
25 Other leaving groups such as iodide, mesylate, tosylate, p-nitrophenyl-sulfonate and the like may also be employed. Appropriate bases include 2,6-di-t-butylpyridine, 2,6-di-t-butyl-4-methyl-pyridine, diisopropyl-ethylamine, potassium carbonate, sodium carbonate, and the like. Suitable solvents include toluene, hexanes, benzene, carbon tetra-
30 chloride, dichloromethane, chloroform, dichloroethane, and the like and mixtures thereof. The filtered solution of the triflate is then added to a solution of the intermediate formed when the morpholinone is contacted with an active hydride reagent such as diisobutyl aluminum hydride, lithium aluminum hydride, or lithium tri(sec-butyl)- borohydride (L-

- 127 -

Selectride®) at low temperature, preferably from -78°C to -20°C. After several hours at low temperature, workup and purification provides predominantly 2,3-cis substituted products, which may be carried on to final compounds as shown in Scheme 6.

Enantiomerically pure phenylglycines substituted on the phenyl ring may be prepared by the procedure shown in Scheme 7 (D.A. Evans, et al, J. Am. Chem. Soc., 1990, 112, 4011).

Methods for preparing the nitrogen alkylating agents R^1CH_2X used in Scheme 5 and Scheme 6 are based on known literature methods (for $R^1 = 3$ -(1,2,4-triazolyl) or 5-(1,2,4-triazol-3-one)-yl and $X = Cl$, see Yanagisawa, I.; Hirata, Y.; Ishii, Y. Journal of Medicinal Chemistry, 27, 849 (1984); for $R^1 = 4$ -((2H)-imidazol-2-one)-yl or 5-(4-ethoxycarbonyl-(2H)-imidazol-2-one)-yl and $X = Br$, see Ducschinsky, R., Dolan, L.A. Journal of the American Chemical Society, 70, 657 (1948)).

One method of producing enantiomerically pure 2,3-cis morpholines that are substituted at the α -position of the C2 benzyl ether is shown in Scheme 8. Thus, a substituted 2-morpholinone (prepared as described in Scheme 5) is reacted with an active hydride reagent, such as diisobutylaluminum hydride, lithium aluminum hydride, or lithium tri(sec-butyl)borohydride and the resulting reaction intermediate is quenched with a substituted benzoyl halide, anhydride, or other activated acyl transfer reagent. Aqueous work-up affords the 2-benzoyloxy compound shown in Scheme 8. This compound is converted to the corresponding enol ether using a "titanium ylide" generated from reagents such as μ -chloro- μ -methylene-[bis(cyclopentadienyl)titanium]dimethylaluminum ("Tebbe Reagent", Tebbe, F.N., Parshall, G.W., Reddy, G.S., Journal of the American Society, 100, 3611 (1978)), dimethyl titanocene (Petasis, N.A., Bzowej, E.I., Journal of the American Chemical Society, 112, 6392 (1990)) or the reagent prepared by the reduction of 1,1-dibromoalkanes with zinc and titanium tetrachloride in the presence of N,N,N',N'-tetramethylethylenediamine (Takai, K. et.al., Journal of Organic Chemistry, 52, 4412 (1987)), wherein $R^{14}-CH_2- = Z$. The resulting enol ether is

- 128 -

reduced to its saturated analog by hydrogenation in the presence of a rhodium based catalyst, such as rhodium on alumina or on carbon; if concomitant removal of the N-benzyl group on the morpholine nitrogen is desired, the hydrogenation may be carried out in the presence of palladium on carbon catalyst. If diastereomers are obtained at this juncture, they may be separated using chromatographic methods or by recrystallization of the mixture of diastereomers. Elaboration of the morpholines so obtained to the final product is carried out in manners analogous to those described in Schemes 5 and 6.

Methods by which the substitution on the C-3 phenyl ring of the morpholines of the present invention may be introduced or altered is shown in Scheme 9. Thus, a substituted morpholine may be prepared as described in Scheme 5, 6, or 8 from an enantiomerically pure benzyloxy-substituted aryl glycine (prepared as described in the literature (e.g. L-p-benzyloxyphenylglycine may be prepared according to the procedure of Kamiya, *et al.* Tetrahedron, **35**, 323 (1979)) or using the methods described in Scheme 7). Selective cleavage of the benzyl ether via hydrogenolysis or nonselective hydrogenolysis followed by the synthetic sequence shown in Scheme 9 may afford a suitably protected phenolic intermediate. The phenol may be converted to the corresponding aryl triflate (as shown, or using N-phenyl-trifluoromethane-sulfonimide in the presence of a tertiary amine base in methylene chloride) and the triflate converted to the desired functional group using the palladium- or nickel-catalyzed methods described in Ritter, Synthesis, 735 (1993) (and refs. therein). Elaboration to the desired final product may be carried out as described in Scheme 5 or 6.

The object compounds of Formula I obtained according to the reactions as explained above may be isolated and purified in a conventional manner, for example, extraction, precipitation, fractional crystallization, recrystallization, chromatography, and the like.

The compounds of the present invention are capable of forming salts with various inorganic and organic acids and bases and such salts are also within the scope of this invention. Examples of such acid addition salts include acetate, adipate, benzoate, benzenesulfonate,

- 129 -

bisulfate, butyrate, citrate, camphorate, camphorsulfonate, diphosphate, ethane-sulfonate, fumarate, hemisulfate, heptanoate, hexanoate, hydrochloride, hydrobromide, hydroiodide, 2-hydroxyethylsulfonate, methanesulfonate, lactate, malate, maleate, methanesulfonate, 2-naphthalenesulfonate, nitrate, oxalate, pamoate, persulfate, phosphate, picrate, pivalate, propionate, salicylate, stearate, succinate, sulfate, tartrate, tosylate, and undecanoate. Base salts include ammonium salts, alkali metal salts such as sodium, lithium and potassium salts, alkaline earth metal salts such as calcium and magnesium salts, salts with aluminum or zinc, salts with organic bases such as dicyclohexylamine salts, N-methyl-D-glucamine (meglumine), and salts with amino acids such as arginine, lysine and so forth. Also, the basic nitrogen-containing groups may be quaternized with such agents as: lower alkyl halides, such as methyl, ethyl, propyl, and butyl chloride, bromides and iodides; dialkyl sulfates like dimethyl, diethyl, dibutyl; diamyl sulfates; long chain halides such as decyl, lauryl, myristyl and stearyl chlorides, bromides and iodides; aralkyl halides like benzyl bromide and others. The non-toxic physiologically acceptable salts are preferred, although other salts are also useful, such as in isolating or purifying the product.

The salts may be formed by conventional means, such as by reacting the free base form of the product with one or more equivalents of the appropriate acid in a solvent or medium in which the salt is insoluble, or in a solvent such as water which is removed in vacuo or by freeze drying or by exchanging the anions of an existing salt for another anion on a suitable ion exchange resin.

Although the reaction schemes described herein are reasonably general, it will be understood by those skilled in the art of organic synthesis that one or more functional groups present in a given compound of formula I may render the molecule incompatible with a particular synthetic sequence. In such a case an alternative route, an altered order of steps, or a strategy of protection and deprotection may be employed. In all cases the particular reaction conditions, including reagents, solvent, temperature, and time, should be chosen so that they

- 130 -

are consistent with the nature of the functionality present in the molecule.

The following examples are given for the purpose of illustrating the present invention and shall not be construed as being limitations on the scope or spirit of the instant invention.

EXAMPLE 1

(+/-)- α -Bromo-phenylacetaldehyde, 3,5-bis(trifluoro-methyl)benzyl acetal

A solution of 2.50 g (10.2 mmol) of α -bromo-phenylacetaldehyde, dimethyl acetal, 8.00 g (32.8 mmol) of 3,5-bis(trifluoromethyl)benzyl alcohol and 0.50 g (2.6 mmol) of p-toluenesulfonic acid monohydrate in 10 mL of toluene was stirred under vacuum (35 mmHg) at rt for 3 days. The reaction mixture was partitioned between 100 mL of ether and 50 mL of saturated aqueous sodium bicarbonate solution and the layers were separated. The organic layer was washed with 25 mL of saturated aqueous sodium chloride solution, dried over magnesium sulfate, and concentrated in vacuo.

Flash chromatography on 200 g of silica gel using 9:1 v/v hexane/methylene chloride as the eluant afforded 5.41 g (81%) of the title compound as a solid, mp 79-82°C: ¹H NMR 4.47 and 4.62 (AB q, 2 H, J = 12.5), 4.78-4.93 (2 H), 5.09 and 5.21 (AB q, 2 H, J = 7.7), 7.31-7.44 (m, 7 H), 7.70 (app s, 1 H), 7.82 (app s, 1 H), 7.84 (app s 2 H);

IR (thin film) 1363, 1278, 1174, 1130, 704, 682.

Analysis Calcd for C₂₆H₁₇BrF₁₂O₂:

C, 46.76; H, 2.23; Br, 11.64; F, 33.70.

Found: C, 46.65; H, 2.56; Br, 11.94; F, 34.06.

EXAMPLE 2

(+/-)-N-(2-Hydroxyethyl)-phenylglycinal, 3,5-bis-(trifluoromethyl)-benzyl acetal

- 131 -

A solution of 1.50 g (2.2 mmol) of (+/-)- α -bromo-phenylacetaldehyde, 3,5-bis(trifluoromethyl)-benzyl acetal (Example 1), 100 mg (0.67 mmol) of sodium iodide and 3 mL of ethanolamine in 6 mL of isopropanol was heated at reflux for 20 h. The solution was cooled and concentrated to ~25% the original volume in vacuo. The concentrated solution was partitioned between 50 mL of ether and 20 mL of 2 N aqueous sodium hydroxide solution and the layers were separated. The organic layer was washed with 20 mL of saturated aqueous sodium chloride solution, dried over magnesium sulfate and concentrated in vacuo. Flash chromatography on 50 g of silica gel using 65:35 v/v ether/hexane as the eluant afforded 1.18 g (83%) of the title compound as an oil: ^1H NMR 2.66 (br s, 2 H), 2.61 and 2.68 (ddAB q, 2 H, $J_{AB} = 12.4$, $J_{2.61} = 6.8$, 6.2, $J_{2.68} = 6.2$, 6.2), 3.57 and 3.66 (ddAB q, 2 H, $J_{AB} = 10.8$, $J_{3.57} = 6.2$, 6.2), $J_{3.66} = 6.8$, 6.2), 4.02 (d, 1 H, $J = 7.0$), 4.37 and 4.64 (AB q, 2 H, $J = 12.5$), 4.80 and 4.87 (AB q, 2 H, $J = 12.8$), 4.87 (d, 1 H, $J = 7.0$), 7.31-7.40 (7 H), 7.73 (app s, 1 H), 7.81 (app s, 3 H); IR (neat) 3342, 1456, 1373, 1278, 1173, 1128, 704, 682; FAB-MS 650(M+1)⁺.
Analysis Calcd for $\text{C}_{28}\text{H}_{23}\text{F}_{12}\text{NO}_3$:
C, 51.78; H, 3.57; N, 2.16; F, 35.11.
Found: C, 51.80; H, 3.67; N, 2.10; F, 35.41.

EXAMPLE 3

(+/-)-N-(2-Hydroxyethyl)-N-(prop-2-enyl)-phenyl-glycinal, 3,5-bis(trifluoromethyl)benzyl acetal

A mixture of 1.45 g (2.2 mmol) of (+/-)-N-(2-hydroxyethyl)-phenylglycinal, 3,5-bis-(trifluoromethyl)benzyl acetal (Example 2), 1.0 g (7.2 mmol) of potassium carbonate, 3.0 mL (35.0 mmol) of allyl bromide and 15 mL of ethanol was stirred at 60 °C for 20 h. The mixture was cooled, partitioned between 100 mL of ether and 25 mL of water and the layers were separated. The organic layer was dried over magnesium sulfate. The aqueous layer was extracted

- 132 -

with 100 mL of ether; the ether extract was dried and combined with the original organic layer. The combined organic layers were concentrated in vacuo. Flash chromatography on 50 g of silica gel using 4:1 v/v hexane/ether as the eluant afforded 1.36 g (88%) of the title compound as an oil: ¹H NMR 2.40 (dt, 1 H, J = 13.2, 2.8), 2.93-3.08 (3 H), 3.30 (ddt, 1 H, J = 12.0, 2.8, 1.6), 3.54 (br m, 2 H), 3.65 (dt, 1 H, J = 10.0, 2.8), 4.23 (d, 1 H, J = 8.4), 4.52 and 4.58 (AB q, 2 H, J = 12.4), 4.85 and 4.95 (AB q, 2 H, J = 12.4), 5.25 (d, 1 H, J = 9.6), 5.28 (d, 1 H, J = 16.4), 5.39 (d, 1 H, J = 8.4), 5.81 (m, 1 H), 7.24-7.40 (7 H), 7.68 (s, 1 H), 7.83 (s, 1 H), 7.86 (s, 2 H); IR (neat) 3457, 1362, 1278, 1174, 1132, 1056, 759, 705, 682; FAB-MS 690(M+1)+.

Analysis Calcd for C₃₁H₂₇F₁₂NO₃:
C, 53.99; H, 3.95; N, 2.03; F, 33.07.
Found: C, 54.11; H, 4.08; N, 1.78; F, 32.75.

EXAMPLE 4

(+/-)-2-(3,5-Bis(trifluoromethyl)benzyloxy)-3-phenylmorpholine

Step A:

A solution of 850 mg (1.2 mmol) of (+/-)-N-(2-hydroxyethyl)-N-(prop-2-enyl)-phenyl-glycinal, 3,5-bis(trifluoromethyl)benzyl acetal (Example 3) and 700 mg (3.7 mmol) of p-toluenesulfonic acid monohydrate in 15 mL of toluene was heated at reflux for 1.5 h. The reaction mixture was cooled and partitioned between 100 mL of ether and 25 mL of saturated aqueous sodium bicarbonate solution. The layers were separated; the organic layer was washed with 25 mL of saturated aqueous sodium chloride solution, dried over magnesium sulfate, and concentrated in vacuo. Flash chromatography on 30 g of silica gel using 50:1 v/v hexane/ether as the eluant afforded 426 mg (78%) of the N-allyl morpholines which were used in the next step without further purification.

- 133 -

Step B:

A 50 mL 2-necked flask, equipped with a stopper and a short path distillation apparatus, was charged with a solution of the N-allyl morpholines (Example 4, Step A) (540 mg, 1.2 mmol) and 80 mg (0.09 mmol) tris(triphenylphosphine)rhodium chloride (Wilkinson's catalyst) in 25 mL of 4:1 v/v acetonitrile/water. The reaction mixture was heated to boiling and solvent was allowed to distill from the reaction mixture. The volume of the reaction mixture was maintained between 10 and 20 mL by adding solvent through the stoppered inlet. After 1 h and 4 h, the reaction was treated with additional 80 mg portions of the Wilkinson's catalyst. After 6 h, the reaction mixture cooled and partitioned between 75 mL of ether and 50 mL of water. The layers were separated and the organic layer was dried over magnesium sulfate. The aqueous layer was extracted with 75 mL of ether; the extract was dried and combined with the original organic layer. The combined organic layers were concentrated in vacuo. Flash chromatography on 35 g of silica gel using 1:1 v/v ether/hexane as the eluant afforded 200 mg of trans-isomer and 130 mg of a mixture of cis- and trans-isomers (68% total). Chromatography of the mixture on 8 g of silica gel using 4:1 v/v hexane/ether as the eluant afforded 64 mg of cis and 57 mg of a mixture of the cis- and trans-isomers of the title compound.

For trans: ^1H NMR 2.03 (br s, 1 H), 2.94 (ddd, 1 H, $J = 11.0, 2.5, 2.5$), 3.08 (dt, 1 H, $J = 11.0, 3.2$), 3.71 (d, 1 H, $J = 7.0$), 3.83 (dt, 1 H, $J = 11.2, 2.8$), 4.05 (ddd, 1 H, $J = 11.2, 3.2, 3.2$), 4.43 (d, 1 H, $J = 7.0$), 4.53 and 4.88 (AB q, 2 H, $J = 13.3$), 7.26-7.45 (7 H), 7.70 (s, 1 H); IR (neat) 3333, 2859, 1456, 1374, 1278, 1173, 1131, 1082, 757, 702, 682;

FAB-MS 406(M+1)+.

Analysis Calcd for $\text{C}_{19}\text{H}_{17}\text{F}_6\text{NO}_2$:

C, 56.30; H, 4.23; N, 3.46; F, 28.12.

Found: C, 56.39; H, 4.28; N, 3.36; F, 28.32.

For cis: ^1H NMR 2.10 (br s, 1 H), 3.13 (dd, 1 H, $J = 12.4, 3.0$), 3.26 (dt, 1 H, $J = 12.4, 3.6$), 3.65 (dd, 1 H, $J = 11.6, 3.6$), 4.07 (dt, 1 H, $J =$

- 134 -

11.6, 3.0), 4.14 (d, 1 H, J = 2.4), 4.52 and 4.82 (AB q, 2 H, J = 13.6), 4.76 (d, 1 H, J = 2.4), 7.30-7.42 (6 H), 7.70 (s, 1 H),
FAB-MS 406(M+1)+.

5

EXAMPLE 5

(+/-)-2-(3,5-Bis(trifluoromethyl)benzyloxy)-3-phenyl-4-methylcarbox-
amido morpholine

10 A solution of 105 mg (0.26 mmol) of the trans-isomer of
(+/-)-2-(3,5-bis(trifluoromethyl)benzyloxy)-3-phenyl-morpholine
(Example 4) and 0.09 mL (0.50 mmol) of N,N-diisopropylethylamine
in 3 mL of acetonitrile was treated with 90 mg (0.50 mmol) of
iodoacetamide and the resulting solution was stirred at rt for 16 h. The
15 solution was concentrated in vacuo and the residue was partitioned
between 20 mL of ethyl acetate and 10 mL of 0.5 N aqueous potassium
hydrogen sulfate solution. The layers were separated; the organic layer
was washed with 10 mL of 5% aqueous sodium thiosulfate solution, 10
mL of saturated aqueous sodium bicarbonate solution, 10 mL of
20 saturated aqueous sodium chloride solution, dried over magnesium
sulfate and concentrated in vacuo. Flash chromatography on 5 g of
silica gel using 2:1 v/v ethyl acetate/hexane as the eluant afforded 99 mg
(82%) of the trans-isomer of the title compound as an oil: ¹H NMR
2.56 (dt, 1 H, J = 3.2, 11.6), 2.67 and 3.16 (AB q, 2 H, J = 16.4), 2.96
(dt, 1 H, J = 12.0, 1.6), 3.30 (d, 1 H, J = 7.0), 3.86 (dt, 1 H, J = 3.2,
25 12.0), 4.08 (ddt, 1 H, J = 11.6, 3.2, 1.6), 4.48 and 4.84 (AB q, 2 H, J =
13.2), 4.49 (d, 1 H, J = 7.0), 5.98 (br s, 1 H), 6.83 (br s, 1 H), 7.33
(app s, 7 H), 7.70 (s, 1 H);
IR (neat) 3445, 2838, 1682, 1278, 1173, 1132, 760, 704, 682; FAB-MS
463 (M+1)+.

30 Analysis Calcd for C₂₁H₂₀F₆NO₃:
C, 54.54; H, 4.36; N, 6.06; F, 24.65.
Found: C, 54.54; H, 4.52; N, 5.61; F, 24.45.

A similar experiment was carried out on 40 mg (0.99
mmol) of the cis-isomer of (+/-)-2-(3,5-bis-(trifluoromethyl)-

- 135 -

benzyloxy)-3-phenyl-morpholine (Example 4) using 0.035 mL (0.2 mmol) of N,N-diisopropylethylamine and 37 mg (0.2 mmol) of iodoacetamide in the reaction. Work-up and flash chromatography afforded 30 mg (65%) of the cis-isomer of the title compound as an oil:
5 ¹H NMR 2.54 and 3.04 (AB q, 2 H, J = 16.8), 2.63 (dt, 1 H, J = 3.6, 12.0), 3.04 (d, 1 H, J = 11.6), 3.65 (d, 1 H, J = 2.8), 3.71 (ddt, 1 H, J = 11.6, 3.2, 1.2), 4.21 (dt, 1 H, J = 11.6, 2.4), 4.44 and 4.89 (AB q, 2 H, J = 13.6), 4.71 (d, 1 H, J = 2.8), 5.86 (br s, 1 H), 7.15 (br s, 1 H), 7.27-7.45 (7 H), 7.73 (s, 1 H); FAB-MS 463(M+1)+.

EXAMPLE 6

(+/-)-2-(3,5-Bis(trifluoromethyl)benzyloxy)-3-phenyl-4-(methoxycarbonylmethyl)morpholine
15 A solution of 150 mg (0.37 mmol) of the trans-isomer of (+/-)-2-(3,5-bis(trifluoromethyl)benzyloxy)-3-phenyl morpholine (Example 4) and 0.18 mL (1.00 mmol) of N,N-diisopropyl-ethyl-amine in 2 mL of acetonitrile was treated with 0.095 mL (1.00 mmol) of methyl bromoacetate and the resulting solution was stirred at rt for 20
20 h. The solution was concentrated in vacuo and the residue was partitioned between 20 mL of ethyl acetate and 5 mL of 0.5 N aqueous potassium hydrogen sulfate solution. The layers were separated; the organic layer was washed with 10 mL of saturated aqueous sodium chloride solution, dried over magnesium sulfate and concentrated in
25 vacuo. Flash chromatography on 10 g of silica gel using 4:1 v/v hexanes/ether as the eluant afforded 164 mg (93%) of the trans-isomer of the title compound as an oil: ¹H NMR 2.79 (dt, 1 H, J = 3.2, 11.2), 2.93 (dt, 1 H, J = 11.2, 1.6), 3.52 (d, 1 H, J = 7.2), 3.63 (s, 3 H), 3.92 (dt, 1 H, J = 2.8, 11.6), 4.04 (ddd, 1 H, J = 11.6, 3.2, 1.6), 4.45 and
30 4.84 (AB q, 2 H, J = 13.2), 4.46 (d, 1 H, J = 7.2), 7.31 - 7.38 (m, 6 H), 7.68 (s, 1 H);
IR (neat) 2861, 1744, 1455, 1375, 1346, 1278, 1170, 887, 759, 704, 682; FAB-MS 478(M+1)+.
Analysis Calcd for C₂₂H₂₁F₆NO₄:

- 136 -

C, 55.35; H, 4.43; N, 2.93; F, 23.88.
Found: C, 55.74; H, 4.50; N, 2.79; F, 24.01.

EXAMPLE 7

N-Methoxy-N-methyl-(N-t-butoxycarbonyl)-phenylglycinamide

A solution of 20.0 g (79.7 mmol) of (N-t-butoxycarbonyl)-phenylglycine in 150 mL of ethyl acetate at -10°C was treated with 8.8 mL (79.7 mmol) of 4-methylmorpholine. Isobutylchloroformate (10.3 mL, 79.7 mmol) was added dropwise over 10 minutes maintaining the temperature at -10°C; the resulting suspension was stirred cold for 15 min. The mixture was treated with 11.6 g (119.0 mmol) of N,O-Dimethyl-hydroxylamine • HCl. A second portion of 4-methylmorpholine (13.0 mL, 119.0 mmol) was added and the reaction was stirred at -10°C for 15 min and at 25°C for 2 h. The reaction mixture was partitioned between 100 mL of ethyl acetate and 100 mL of 10% aqueous citric acid solution and the layers were separated. The organic layer was washed with 100 mL of saturated aqueous sodium bicarbonate solution, 100 mL of saturated aqueous ammonium chloride solution, dried over magnesium sulfate and concentrated in vacuo. Crystallization from hexanes at -20°C for 72 h afforded 8.0 g (34%) of the title compound as a solid: ¹H NMR 1.40 (s, 9 H), 3.20 (s, 3 H), 3.40 (s, 3 H), 5.80 (m, 2 H), 7.40 (m, 5 H).

EXAMPLE 8

Diethyl (2-oxo-3-t-butoxycarbamido-3-phenyl)-propylphosphonate

A solution of 7.45 mL (51.0 mmol) of diethyl methylphosphonate in tetrahydrofuran at -78°C was treated with 31.8 mL (51.0 mmol) of 1.6 M n-butyllithium in hexanes solution and the resulting mixture was stirred cold for 30 min. A solution of 4.0 g (14.0 mmol) of N-methoxy-N-methyl-(N-t-butoxycarbonyl)phenylglycinamide (Example 7) in 20 mL of tetrahydrofuran was added and the reaction was stirred at -78°C for 15 min and at 25°C for 15 min.

- 137 -

The reaction was quenched with 150 mL of saturated aqueous ammonium chloride solution, diluted with 300 mL of ethyl acetate, and the layers were separated. The organic layer was dried over magnesium sulfate and concentrated in vacuo. Flash chromatography on silica gel using 7:3 v/v then 4:1 v/v ethyl acetate/hexanes as the eluant afforded 4.8 g (92%) of the title compound as an oil: ¹H NMR 1.20-1.42 (15 H), 2.84 (dd, 1 H), 3.20 (dd, 1 H), 4.00-4.20 (m, 4 H), 5.50 (d, 1 H), 5.94 (br s, 1 H), 7.32 (m, 5 H).

EXAMPLE 9

N-t-Butoxycarbonyl-1-phenyl-2-oxo-4-(3,5-bis(tri-fluoromethyl)-phenyl)-but-3-enamine

A solution of 4.80 g (12.5 mmol) of diethyl (2-oxo-3-t-butoxycarbamido-3-phenyl)propylphosphonate (Example 8) in 20 mL of THF was added dropwise to a suspension of 1.05 g (26.3 mmol, 60% dispersion in mineral oil) of sodium hydride in 30 mL of tetrahydrofuran at 0°C. After 15 min, 2.06 mL (12.5 mmol) of 3,5-bis(trifluoromethyl)benzaldehyde was slowly added and the resulting mixture was stirred cold for 15 min. The reaction was quenched with 50 mL of saturated aqueous ammonium chloride solution, diluted with 50 mL of ethyl acetate, and the layers were separated. The organic layer was dried over magnesium sulfate and concentrated in vacuo. Flash chromatography on silica gel using 19:1 v/v, then 9:1 v/v ethyl acetate/petroleum ether as the eluant afforded 3.30 g (56%) of the title compound as a solid: ¹H NMR 1.40 (s, 9 H), 5.38 (d, 1 H), 5.90 (d, 1 H), 6.80 (d, 1 H), 7.39 (m, 5 H), 7.70 (s, 1 H), 7.84 (s, 3 H).

EXAMPLE 10

1-Phenyl-2-hydroxy-4-(3,5-bis(trifluoromethyl)phenyl)-but-3-enamine • HCl

A solution of 1.00 g (2.1 mmol) of N-t-butoxycarbonyl-1-phenyl-2-oxo-4-(3,5-bis(trifluoromethyl)phenyl)-but-3-enamine

- 138 -

(Example 8) in 30 mL of methanol at 0°C was treated with 241 mg (6.3 mmol) of sodium borohydride. After 30 min, the reaction was quenched with 50 mL of water and concentrated in vacuo to remove the methanol. The mixture was partitioned between 100 mL of ethyl acetate and 50 mL of water and the layers were separated. The organic layer was dried over magnesium sulfate and concentrated in vacuo. Crystallization from ether/hexanes afforded 680 mg (68%) of the title compound as a 5:1 mixture of diastereomers (each protected as the t-butylcarbamate): ¹H NMR (* indicates the resonances of the minor diastereomer) 1.40 (s, 9 H), 4.60 (dd, 1 H), 4.90 (br s, 1 H), 5.20 (br d, 1 H), 6.30 (dd, 1 H), 6.40 (dd, 1 H*), 6.70 (dd, 1 H), 6.80 (dd, 1 H*), 7.40 (m, 5 H), 7.80 (m, 3 H).

A solution of BOC-protected title compound in methanol (saturated with HCl) was allowed to stand for 72 h. The solution was concentrated in vacuo. Recrystallization of the resulting solid from ether/hexane afforded 500 mg (80%) of the title compound • HCl as a solid: ¹H NMR 4.20 (br s, 1 H), 4.40 (d, 1 H), 6.20 (dd, 1 H), 6.60 (dd, 1 H), 7.30 (m, 5 H), 7.80 (m, 3 H).

The title compound • HCl was dissolved in ethyl acetate and 1 N aqueous sodium hydroxide solution. The layers were separated; the organic layer was dried over magnesium sulfate and concentrated in vacuo to afford the title compound as the free base.

EXAMPLE 11

2-(2-(3,5-Bis(trifluoromethyl)phenyl)ethenyl)-3-phenyl-5-oxo-morpholine

A solution of 1.95 g (5.2 mmol) of 1-phenyl-2-hydroxy-4-(3,5-bis(trifluoromethyl)phenyl)-but-3-enamine (Example 10) in 20 mL of toluene was added to a suspension of 250 mg (6.2 mmol, 60% dispersion in mineral oil) of sodium hydride in 30 mL of toluene and the resulting mixture was stirred at rt for 15 min. A solution of 0.60 mL (1.15 mol) of ethyl chloroacetate in 5 mL of toluene was slowly added and the resulting mixture was heated at reflux for 3 h. The

- 139 -

reaction was cooled, quenched with 50 mL of saturated aqueous ammonium chloride solution, diluted with 50 mL of ethyl acetate and the layers were separated. The organic layer was dried over magnesium sulfate and concentrated in vacuo. Flash chromatography using ethyl acetate/hexanes (4:1 v/v, then 3:1 v/v, then 1:1 v/v) then ethyl acetate as the eluant afforded 300 mg of trans-title compound and 800 mg of cis-title compound (55% total), both as solids. For the cis-isomer: ¹H NMR 1.20-1.40 (m, 1 H), 1.50-1.62 (m, 1 H), 2.60-2.98 (m, 2 H), 3.86 (dt, 1 H), 4.24 (d, 1 H), 4.34 (dd, 1 H), 4.45 (d, 1 H), 6.40 (br s, 1 H), 7.24 (m, 2 H), 7.40 (m, 3 H), 7.50 (s, 2 H), 7.70 (s, 1 H).

EXAMPLE 12

3-Phenyl-2-(2-(3,5-bis(trifluoromethyl)phenyl)ethyl)-morpholine

A solution of 95 mg (0.23 mmol) of 2-(2-(3,5-bis(trifluoromethyl)phenyl)ethenyl)-3-phenyl-5-oxo-morpholine (Example 11) in 10 mL of 1:1 v/v ethanol/ ethyl acetate was treated with 10 mg of palladium hydroxide and the resulting mixture was stirred under an atmosphere of hydrogen for 2 h. The catalyst was filtered and the filtrate was concentrated in vacuo. The crude product was used directly without further purification.

A solution of 65 mg of the crude morpholinone was dissolved in 10 mL of tetrahydrofuran was treated with 0.84 mL of 1 M borane-tetrahydrofuran complex solution in tetrahydrofuran and the resulting solution was heated at reflux for 16 h. The reaction was quenched by adding 10 mL of methanol and 70 mg of potassium carbonate and heating the resulting mixture at reflux for 3 h. All volatiles were removed in vacuo and the residue was partitioned between 20 mL of ethyl acetate and 10 mL of saturated ammonium chloride solution. The organic layer was separated, dried over sodium carbonate, and concentrated in vacuo. The residue was dissolved in saturated HCl in methanol and concentrated in vacuo. The residue was triturated with ether; the resulting solid was filtered and dried to afford 32 mg (46%) of the title compound • HCl, mp 114-116°C: ¹H NMR

- 140 -

1.42 (m, 1 H), 1.66-1.84 (m, 1 H), 2.70-2.94 (m, 2 H), 3.00 (m, 1 H),
3.30-3.46 (m, 1 H), 3.80-3.94 (m, 2 H), 4.10 (m, 1 H), 4.20 (d, 1 H),
7.40 (m, 3 H), 7.64 (m, 5 H); CI-MS 402(M+1)+.

5

EXAMPLE 13N-Benzyl-(S)-phenylglycine

A solution of 1.51 g (10.0 mmol) of (S)-phenylglycine in 5 mL of 2 N aqueous sodium hydroxide solution was treated with 1.0 mL
10 (10.0 mmol) of benzaldehyde and stirred at room temperature for 20 minutes. The solution was diluted with 5 mL of methanol, cooled to 0°C, and carefully treated with 200 mg (5.3 mmol) of sodium borohydride. The cooling bath was removed and the reaction mixture was stirred at room temperature for 1.5 hours. The reaction was
15 diluted with 20 mL of water and extracted with 2x25 mL of methylene chloride. The aqueous layer was acidified with concentrated hydrochloric acid to pH 6 and the solid that precipitated was filtered, washed with 50 mL of water, 50 mL of 1:1 v/v methanol/ethyl ether and 50 mL of ether, and dried to afford 1.83 g (76%) of product, mp
20 230-232°C.

Analysis Calcd for C₁₅H₁₅NO₂:

C, 74.66; H, 6.27; N, 5.81.

Found: C, 74.17; H, 6.19; N, 5.86.

25

EXAMPLE 143-(S)-Phenyl-4-benzyl-2-morpholinone

A mixture of 4.00 g (16.6 mmol) of N-benzyl-(S)-phenylglycine (from Example 13), 5.00 g (36.0 mmol) of potassium carbonate, 10.0 mL of 1,2-dibromoethane and 25 mL of N,N-dimethylformamide was stirred at 100°C for 20 hours. The mixture was cooled and partitioned between 200 mL of ethyl ether and 100 mL of water. The layers were separated and the organic layer was washed with 3x50 mL of water, dried over magnesium sulfate and concentrated.

- 141 -

in vacuo. The residue was purified by flash chromatography on 125 g of silica gel eluting with 9:1 v/v, then 4:1 v/v hexanes/ethyl ether to afford 2.41 g (54%) of the product as a solid, mp 98-100°C.

Mass Spectrum (FAB): m/Z 268 (M+H, 100%).

5 ¹H NMR (CDCl₃, 200 MHz, ppm): δ 2.54-2.68 (m, 1H), 2.96 (dt, J= 12.8, 2.8, 1H), 3.14 (d, J= 13.3, 1H), 3.75 (d, J= 13.3, 1H), 4.23 (s, 1H), 4.29-4.37 (m, 1H), 4.53 (dt, J= 3.2, 11.0), 7.20-7.56 (m, 10H).

Analysis Calcd for C₁₇H₁₇NO₂:

C, 76.38; H, 6.41; N, 5.24.

10 Found: C, 76.06; H, 6.40; N, 5.78.

EXAMPLE 15

15 2-(S)-(3,5-Bis(trifluoromethyl)benzyloxy)-3-(S)-phenylmorpholine

Step A: 3,5-Bis(trifluoromethyl)benzyl alcohol, trifluoromethanesulfonate ester

20 A solution of 1.00 g (4.1 mmole) of 3,5-bis(trifluoromethyl)benzyl alcohol and 1.05 g (5.12 mmole) of 2,6-di-t-butyl-4-methylpyridine in 45 mL of dry carbon tetrachloride under a nitrogen atmosphere was treated with 0.74 mL (4.38 mmole) of trifluoromethanesulfonic anhydride at room temperature. A white precipitate formed shortly after the addition of the anhydride. After 90 min, the slurry was filtered under nitrogen with a Schlenk filter, and the filtrate
25 was concentrated in vacuo. The residue, which was a two-phase oil, was dissolved under nitrogen in 10 mL of dry toluene. The resulting clear solution was used immediately in Step B below.

30 Step B: 4-Benzyl-2-(S)-(3,5-bis(trifluoromethyl)benzyloxy)-3-(S)-phenylmorpholine

A solution of 0.500 g (1.87 mmole) of N-benzyl-3-(S)-phenylmorpholin-2-one (from Example 14) in 10 mL of dry THF was cooled to -75°C under nitrogen and was treated dropwise with 2.06 mL (2.06 mmole) of a 1M solution of lithium tri(sec-butyl)-borohydride

- 142 -

(L-Selectride®) in THF. After stirring the solution at -75°C for 30 min, a solution of 3,5-bis(trifluoromethyl)benzyl alcohol, trifluoromethanesulfonate ester in toluene was added by cannula so that the internal temperature was maintained below -60°C. The resulting
5 solution was stirred at -75°C for 1 hr and then between -38°C and -50°C for 2 hr. The solution was then poured into a mixture of 25 mL of ethyl acetate and 20 mL of saturated aqueous sodium bicarbonate, and the layers were separated. The aqueous phase was extracted with 2x30
10 mL of ethyl acetate, the combined organic layers were dried over sodium sulfate, the mixture was filtered and the filtrate concentrated in vacuo. The residue was purified by flash chromatography on 130 g of silica eluting with 2 L of 100:5 hexanes:ethyl acetate to give 0.68 g (73%) of an oil, which by ¹H NMR is a 20:1 mixture of cis:trans morpholines.

15 ¹H NMR (CDCl₃, 400 MHz, ppm): δ major (cis) isomer: 2.37 (td, J= 12, 3.6, 1H), 2.86 (app t, J= 13, 2H), 3.57 (d, J= 2.6, 1H), 3.63 (dq, J= 11.3, 1.6, 1H), 3.89 (d, J= 13.3, 1H), 4.12 (td, J= 11.6, 2.4, 1H), 4.40 (d, J= 13.6, 1H), 4.69 (d, J= 2.9, 1H), 4.77 (d, J= 13.6), 7.2-7.4 (m, 8H), 7.43 (s, 2H), 7.55 (br d, 2H), 7.69 (s, 1H).
20

Step C: 2-(S)-(3,5-Bis(trifluoromethyl)benzyloxy)-3-(S)-phenylmorpholine

A mixture of 0.68 g (1.37 mmole) of 4-benzyl-2-(S)-(3,5-bis(trifluoromethyl)benzyloxy)-3-(S)-phenylmorpholine and 280 mg of
25 10% Pd/C in 36 mL of 97:3 ethanol:water was stirred under one atmosphere of hydrogen for 15 hr. The mixture was filtered through Celite, the filter cake was washed generously with ethanol, and the filtrate was concentrated in vacuo. The residue was purified by flash chromatography on 68 g of silica eluting with 1L of 33:67 hexanes:
30 diethyl ether, then 1L of 25:75 hexanes: diethyl ether to give 0.443 g (80%) of an oil, which by ¹H NMR was pure cis morpholine.

¹H NMR (CDCl₃, 400 MHz, ppm): δ 1.8 (br s, 1H), 3.10 (dd, J= 12.5, 2.9, 1H), 3.24 (td, J= 12.2, 3.6, 1H), 3.62 (dd, J= 11.3, 2.5, 1H), 4.04 (td, J= 11.7, 3, 1H), 4.11 (d, J= 2.4, 1H), 4.49 (d, J= 13.5, 1H), 4.74 (d,

- 143 -

J= 2.5, 1H), 4.80 (d, J= 13.3, 1H), 7.25-7.40 (m, 5H), 7.40 (s, 2H), 7.68 (s, 1H).

Analysis Calcd for C₁₉H₁₇F₆NO₂:

C, 56.30; H, 4.23; N, 3.46; F, 28.12.

5 Found: C, 56.20; H, 4.29; N, 3.34; F, 27.94.

EXAMPLE 16

10 2(R)-(3,5-Bis(trifluoromethyl)benzyloxy)-3(R)-phenyl-morpholine

The title compound was prepared from (R)-phenylglycine employing the procedures of Examples 13, 14 and 15.

EXAMPLE 17

15 4-(3-(1,2,4-Triazolo)methyl)-2-(S)-(3,5-bis(trifluoro-methyl)-benzyloxy)-3-(S)-phenylmorpholine

Step A: N-Formyl-2-chloroacetamidrazone

20 A solution of 5 g (66.2 mmole) of chloroacetonitrile in 30 mL of dry methanol was cooled to 0°C under nitrogen and was treated with 0.1g (1.8 mmole) of sodium methoxide. The mixture was allowed to warm to room temperature and was stirred for 30 min, and 0.106 mL (1.8 mmole) of acetic acid was added. To the resulting mixture was then added 3.9 g (64.9 mmole) of formic hydrazide, and the material
25 was stirred for 30 min. The reaction mixture was concentrated in vacuo to a solid, and was used as such in Step B below.

Step B: 4-(3-(1,2,4-Triazolo)methyl)-2-(S)-(3,5-bis-(trifluoro-methyl)benzyloxy)-3-(S)-phenyl-morpholine

30 A solution of 0.295g (0.73 mmole) of 2-(S)-(3,5-bis(trifluoromethyl)benzyloxy)-3-(S)-phenylmorpholine (from Example 15) in 10 mL of dry DMF was treated with 0.302g (2.18 mmole) of anhydrous potassium carbonate and then 0.168g (1.24 mmole) of N-formyl-2-chloroacetamidrazone (from Example 17, Step A) and the

- 144 -

suspension was stirred at 60°C for 4 hr. The mixture was then heated to 120°C for 4.5 hr. After cooling, the reaction was diluted with 80 mL of ethyl acetate and the organic layer was washed with 3x20 mL of water. The organic layer was dried over magnesium sulfate, filtered
5 and concentrated in vacuo. The residue was purified by flash chromatography on 67 g of silica eluting with 1.5 L of 100:2 methylene chloride:methanol to give 0.22 g of a yellow solid, which was recrystallized from hexanes/methylene chloride to give 0.213g (60%) of a white crystalline solid, mp 134-135°C.

10 Mass Spectrum (FAB): m/Z 487 (M+H, 100%), 259 (35%), 243 (65%), 227 (40%), 174 (25%).

¹H NMR (CDCl₃, 400 MHz, ppm): δ 2.67 (td, J= 11.9, 3.4, 1H), 2.90 (br d, J= 11.7, 1H), 3.43 (d, J= 15.2, 1H), 3.66 (app dd, J= 13, 1.9, 2H), 3.88 (d, J= 15.1, 1H), 4.17 (td, J= 11.7, 2.3, 1H), 4.42 (d, J= 13.5, 1H),
15 4.69 (d, J= 2.6, 1H), 4.77 (d, J= 13.5, 1H), 7.30-7.50 (m, 7H), 7.70 (s, 1H), 7.94 (s, 1H).

EXAMPLE 18

20 4-(3-(5-Oxo-1H,4H-1,2,4-triazolo)methyl)-2-(S)-(3,5-bis(trifluoromethyl) benzyloxy)-3-(S)-phenylmorpholine

Step A: N-Methylcarboxy-2-chloroacetamidrazone

A solution of 5.0 g (66.2 mmol) of chloroacetonitrile in 35
25 mL of dry methanol was cooled to 0°C and was treated with 0.105 g (1.9 mmol) of sodium methoxide. The ice-bath was removed and the mixture was allowed to stir at room temperature for 30 minutes. To the reaction was then added 0.110 mL (1.9 mmol) of acetic acid and then
30 5.8 g (64.9 mmol) of methyl hydrazinecarboxylate. After stirring 30 minutes at room temperature, the suspension was concentrated in vacuo, and placed on the high-vac line overnight, to give 10.5 g (98%) of a yellow powder, which was employed in Step C below.

¹H NMR (CD₃OD, 400 MHz, ppm): δ 3.71 (s, 3H), 4.06 (s, 2H).

- 145 -

Step B: 4-(2-(N-Methylcarboxy-acetamidrazono)-2-(S)-(3,5-bis(trifluoromethyl)benzyloxy)-3-(S)-phenylmorpholine

A solution of 2.30 g (5.7 mmol) of 2-(S)-(3,5-bis(trifluoromethyl)benzyloxy)-3-(S)-phenylmorpholine (from Example 15), 1.13 g (6.8 mmol) of N-methylcarboxy-2-chloroacetamidrazone (from Step A), and 1.50 mL (8.6 mmol) N,N-diisopropylethylamine in 25 mL of acetonitrile was stirred at room temperature for 20 hours. The product, which had precipitated, was filtered, washed with 5 mL of ice cold acetonitrile and dried to give 1.83 g of a white solid. The filtrate was concentrated in vacuo and the residue was partitioned between 50 mL of methylene chloride and 20 mL of water. The layers were separated and the organic layer was dried over magnesium sulfate. The aqueous layer was extracted with 50 mL of methylene chloride; the extract was dried, combined with the original organic layer, and the combined organics were concentrated in vacuo. The residue was purified by flash chromatography on 30 g of silica gel eluting with 50:1:0.1 v/v/v methylene chloride/methanol/ammonium hydroxide to afford an additional 1.09 g of product (96% total).
Mass Spectrum (FAB): m/Z 535 (M+H, 100%), 462 (16%), 291 (30%), 226 (35%), 173 (25%).
¹H NMR (CDCl₃, 400 MHz, ppm): δ 2.53 (dt, J= 3.5, 12.2, 1H), 2.59 (d, J= 14.6, 1H), 2.94 (d, J= 11.8, 1H), 3.37 (d, J= 14.6, 1H), 3.58 (d, J= 2.8), 1H), 3.62-3.72 (m, 1H), 3.75 (s, 3H), 4.16 (dt, J= 2.2, 11.8, 1H), 4.44 (d, J= 13.2, 1H), 4.70 (d, J= 2.8, 1H), 4.79 (d, J= 13.2), 5.55 (br s, 2H), 7.30-7.46 (m, 7H), 7.72 (s, 1H).

Step C: 2-(S)-(3,5-Bis(trifluoromethyl)benzyloxy)-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-3-(S)-phenylmorpholine

A solution of 2.89 g (5.4 mmol) of 4-(2-(N-methylcarboxyacetamidrazono)-2-(S)-(3,5-bis(trifluoromethyl) benzyloxy)-3-(S)-phenylmorpholine (from Step B) in 36 mL of xylenes was heated at reflux for 1.5 hours. The solution was cooled and concentrated in vacuo. The residue was taken up in 50 mL of 3:1 v/v hexanes/ethyl acetate which caused crystallization of the product. The product was

- 146 -

filtered and dried to afford 1.85 g of a solid. Recrystallization of the solid from 30 mL of 4:1 v/v hexanes/ethyl acetate afforded 1.19 g of pure product as a white solid, mp= 156-157°C. All of the crystallization liquors were combined and concentrated in vacuo. The residue was purified by flash chromatography on 30 g of silica gel eluting with 50:1:0.1 v/v/v methylene chloride/methanol/ammonium hydroxide to afford an additional 0.69 g of a solid. Three recrystallizations from 20 mL of 4:1 v/v hexanes/ethyl acetate afforded an additional 0.39 g of pure product as a white solid (58% total).
Mass Spectrum (FAB): m/Z 503 (M+H), 259 (55%), 226 (40%), 160 (30%).
¹H NMR (CDCl₃, 400 MHz, ppm): δ 2.57 (app t, J= 9.6, 1H), 2.87-2.97 (m, 2H), 3.58-3.71 (m, 3H), 4.18 (app t, J= 10.4, 1H), 4.46 (d, J= 13.6), 4.68 (d, J= 2.8, 1H), 4.85 (d, J= 13.6, 1H), 7.30-7.45 (m, 7H), 7.64 (s, 1H), 10.40 (br s, 1H), 10.73 (br s, 1H).

EXAMPLE 19

N-(2-(R)-Hydroxypropyl)-phenylglycinal, 3,5-bis(tri-fluoromethyl)-benzyl acetal

A mixture of 1.00 g (1.5 mmol) of (+/-)-a- bromo-phenylacetaldehyde, 3,5-bis(trifluoromethyl)-benzyl acetal (from Example 12), 1.25 mL of (R)-1-amino-2-propanol, 225 mg (1.5 mmol) of sodium iodide, and 3.75 mL of isopropanol was heated at reflux for 20 h. The solution was cooled and concentrated to ~25% the original volume in vacuo. The concentrated solution was partitioned between 50 mL of ether and 20 mL of 2 N aqueous sodium hydroxide solution and the layers were separated. The organic layer was washed with 20 mL of saturated aqueous sodium chloride solution, dried over magnesium sulfate and concentrated in vacuo. Flash chromatography on 50 g of silica gel using 65:35 v/v ether/hexane as the eluant afforded 948 mg (95%) of the product as a 1:1 mixture of inseparable diastereomers. Mass Spectrum (FAB): m/Z 664 (M+H, 25%), 420 (20%), 226 (100%).

- 147 -

EXAMPLE 20

N-(2-(S)-Hydroxypropyl)-phenylglycinal, 3,5-bis(tri-fluoromethyl)-
benzyl acetal

5 Substitution of (S)-1-amino-2-propanol for (R)-1-amino-2-propanol in an experiment identical to the preceding example afforded 940 mg (95%) of the product as a 1:1 mixture of diastereomers.

EXAMPLE 21

10 N-(2-(R)-Hydroxypropyl)-N-(prop-2-enyl)-(R)-phenyl-glycinal, 3,5-bis(trifluoromethyl)benzyl acetal and N-(2-(R)-Hydroxypropyl)-N-(prop-2-enyl)-(S)-phenyl-glycinal, 3,5-bis(trifluoromethyl)benzyl
acetal

15 A mixture of 933 mg (1.40 mmol) of N-(2-(R)-hydroxypropyl)-phenylglycinal, 3,5-bis(trifluoromethyl)-benzyl acetal (from Example 19), 1 mL of allyl bromide, 600 mg (4.3 mmol) of potassium carbonate, and 5 mL of ethanol was stirred at 60°C for 20 hours. The mixture was cooled, partitioned between 100 mL of ethyl ether and 25
20 mL of water and the layers were separated. Flash chromatography on 50 g of silica gel using 20:1 v/v ether/hexanes as the eluant afforded 380 mg of the (R,R)-amino alcohol ($R_f = 0.72$ with 3:2 v/v ether/hexanes as the eluant), 220 mg of the (R,S)-amino alcohol ($R_f = 0.62$ with 3:2 v/v ether/hexanes as the eluant), and 285 mg of a mixture of the
25 diastereomeric amino alcohols.

For the (R,R)-amino alcohol:

Mass Spectrum (FAB): m/Z 704(M+H).

30 IR (neat) 3476, 2932, 1624, 1454, 1361, 1278, 1175, 1132, 760, 704, 682.

1H NMR ($CDCl_3$, 400 MHz, ppm) 1.12 (d, 3 H, $J = 6.4$), 2.19 and 2.62 (dAB q, 2 H, $J_{AB} = 13.0$, $J_{2.19} = 2.3$, $J_{2.62} = 10.4$), 2.97 (dd, 1 H, $J = 14.0, 8.8$), 3.25 - 3.30 (m, 1 H), 3.76 (s, 1 H), 3.77 - 3.85 (m, 1 H), 4.21 (d, 1 H, $J = 8.8$), 4.49 and 4.55 (AB q, 2 H, $J = 12.4$), 4.86 and

- 148 -

4.92 (AB q, 2 H, J = 12.4), 5.27 - 5.33 (m, 2 H), 5.39 (d, 1 H, J = 8.8), 5.79 - 5.89 (m, 1 H), 7.21 - 7.26 (m, 4 H), 7.35 - 7.40 (m, 3 H), 7.67 (s, 1 H), 7.81 (s, 1 H), 7.85 (s, 2 H).

Analysis Calcd for C₃₂H₂₉F₁₂NO₃:

5 C, 54.63; H, 4.15; N, 1.99; F, 32.41.

Found: C, 54.72; H, 3.94; N, 1.95; F, 32.17.

For the (R,S)-amino alcohol:

Mass Spectrum (FAB): m/Z 704(M+1).

10 IR (neat) 3451, 2931, 1624, 1454, 1362, 1277, 704, 683.

¹H NMR (CDCl₃, 400 MHz, ppm) 1.09 (d, 3 H, J = 6.0), 2.48 and 2.71 (dAB q, 2 H, J_{AB} = 13.2, J_{2.48} = 9.6, J_{2.62} = 3.6), 3.05 (dd, 1 H, J = 14.4, 6.8), 3.34 - 3.39 (m, 1 H), 3.35 (s, 1 H), 3.76 - 3.81 (m, 1 H), 4.21 (d, 1 H, J = 8.4), 4.50 and 4.54 (AB q, 2 H, J = 12.8), 4.86 and 15 4.96 (AB q, 2 H, J = 12.4), 5.10 - 5.17 (m, 2 H), 5.39 (d, 1 H, J = 8.4), 5.68 - 5.78 (m, 1 H), 7.23 - 7.32 (m, 4 H), 7.34 - 7.39 (m, 3 H), 7.69 (s, 1 H), 7.83 (s, 1 H), 7.86 (s, 2 H).

Analysis Calcd for C₃₂H₂₉F₁₂NO₃:

20 C, 54.63; H, 4.15; N, 1.99; F, 32.41.

Found: C, 54.80; H, 4.16; N, 1.90; F, 32.36.

EXAMPLE 22

25 N-(2-(S)-Hydroxypropyl)-N-(prop-2-enyl)-(S)-phenyl-glycinal, 3,5-bis(trifluoromethyl)benzyl acetal and N-(2-(S)-Hydroxypropyl)-N-(prop-2-enyl)-(R)-phenyl-glycinal, 3,5-bis(trifluoromethyl)benzyl acetal

30 Substitution of 880 mg (1.33 mmol) of N-(2-(S)-hydroxypropyl)-phenylglycinal, 3,5-bis(trifluoro-methyl)benzyl acetal (Example 20) for the N-(2-(R)-hydroxypropyl)-phenylglycinal, 3,5-bis(trifluoro-methyl)benzyl acetal in the procedures of the preceding example afforded 281 mg of the (S,S)-amino alcohol (R_f = 0.72 with 3:2 v/v ether/hexanes as the eluant), 367 mg of the (S,R)-amino alcohol (R_f =

- 149 -

0.62 with 3:2 v/v ether/hexanes as the eluant), and 197 mg of a mixture of the diastereomeric amino alcohols.

EXAMPLE 23

5

2-(R)-(3,5-Bis(trifluoromethyl)benzyloxy)-3-(R)-phenyl-6-(R)-methyl morpholine and 2-(S)-(3,5-Bis(trifluoromethyl)benzyloxy)-3-(R)-phenyl-6-(R)-methyl morpholine

10 Step A: 2-(R)-(3,5-Bis(trifluoromethyl)benzyloxy)-3-(R)-phenyl-4-(2-propenyl)-6-(R)-methyl morpholine and 2-(S)-(3,5-bis(trifluoro-methyl)-benzyloxy)-3-(R)-phenyl-4-(2-propenyl)-6-(R)-methyl morpholine

15 A solution of 355 mg (0.50 mmol) of N-(2-(R)-hydroxy-propyl)-N-(2-propenyl)-(R)-phenylglycinal, 3,5-bis(trifluoromethyl)-benzyl acetal (from Example 21) and 285 mg (1.5 mmol) of p-toluensulfonic acid monohydrate in 5 mL of toluene was heated at reflux for 40 min. The solution was cooled and partitioned between 40 mL of ether and 15 mL of saturated aqueous sodium bicarbonate
20 solution. The layers were separated; the organic layer was washed with 10 mL of saturated aqueous sodium chloride solution, dried over magnesium sulfate, and concentrated in vacuo. Flash chromatography on 10 g of silica gel using 19:1 v/v hexanes/ether as the eluant afforded 122 mg of (2R,3R,6R) product ($R_f = 0.53$ with 4:1 v/v hexanes/ether as the eluant) and 62 mg of the (2S,3R,6R) product ($R_f = 0.23$ with 4:1
25 v/v hexanes/ether as the eluant).

For the (2R,3R,6R) product:

Mass Spectrum (FAB): m/Z 460 (M+H, 65%)

30 1H NMR ($CDCl_3$, 400 MHz, ppm) 1.35 (d, 3 H, $J = 6.4$), 2.53 and 2.63 (dAB q, 2 H, $J_{AB} = 12.0$, $J_{2.53} = 3.2$, $J_{2.63} = 6.8$), 2.83 - 2.96 (m, 2 H), 3.60 (d, 1 H, $J = 4.0$), 4.27 - 4.32 (m, 1 H), 4.57 and 4.84 (AB q, 2 H, $J = 13.2$), 4.87 (d, 1 H, $J = 4.0$), 5.08 - 5.13 (m, 2 H), 5.76 - 5.86

- 150 -

(m, 1 H), 7.31 - 7.37 (m, 3 H), 7.50 - 7.52 (m, 2 H), 7.58 (s, 2 H), 7.71 (s, 1 H).

For the (2S,3R,6R) product:

5 Mass Spectrum (FAB): m/Z 460 (M+H, 65%)
1H NMR (CDCl₃, 400 MHz, ppm) 1.37 (d, 3 H, $J = 6.8$), 2.48 - 2.50 (m, 2 H), 2.74 and 3.01 (dtAB q, 2 H, $J = 6.4, 1.2, 12.4$) 3.84 (d, 1 H, $J = 3.6$), 3.92 - 3.99 (m, 1 H), 4.70 and 4.93 (AB q, 2 H, $J = 13.6$), 4.97 (d, 1 H, $J = 3.6$), 5.08 - 5.14 (m, 2 H), 5.74 - 5.84 (m, 1 H), 7.28 - 7.36
10 (m, 3 H), 7.43 - 7.46 (m, 2 H), 7.64 (s, 2 H), 7.75 (s, 1 H).

Step B: 2-(R)-(3,5-Bis(trifluoromethyl)benzyloxy)-3-(R)-phenyl-6-(R)-methyl morpholine

15 A solution of 115 mg (0.25 mmol) of the 2-(R)-(3,5-bis(trifluoromethyl)benzyloxy)-3-(R)-phenyl-4-(2-propenyl)-6-(R)-methyl morpholine (from Example 23, Step A) and 230 mg (0.25 mmol) of tris(triphenylphosphine)rhodium chloride in 15 mL of 4:1 v/v acetonitrile/water was heated at reflux for 30 min. The reaction
20 was cooled and partitioned between 50 mL of ethyl acetate and 15 mL of water. The layers were separated and the organic layer was dried over magnesium sulfate. The aqueous layer was extracted with 2 x 25 mL of ethyl acetate; the extracts were dried and combined with the original organic layer. The combined organics were concentrated in vacuo. The residue was filtered through a pad of silica gel (~ 20 g)
25 using 2:1 v/v ether/hexanes as the solvent. The filtrate was concentrated; flash chromatography on 5 g of silica gel using 17:3 v/v hexanes/ether as the eluant afforded 67 mg (64%) of 2-(R)-(3,5-bis(trifluoromethyl)benzyloxy)-3-(R)-phenyl-6-(R)-methyl morpholine as an oil.

30 Mass Spectrum (FAB): m/Z 420 (M+H, 90%)
1H NMR (CDCl₃, 400 MHz, ppm) 1.21 (d, 3 H, $J = 6.4$), 2.02 (br s, 1 H), 2.67 and 2.77 (dAB q, 2 H, $J_{AB} = 13.2, J_{2.67} = 8.8, J_{2.77} = 3.2$), 3.89 (d, 1 H, $J = 2.4$), 4.07 - 4.15 (m, 1 H), 4.68 and 4.90 (AB q,

- 151 -

2 H, J = 12.8), 5.03 (d, 1 H, J = 2.4), 7.28 - 7.38 (m, 3 H), 7.51 - 7.53 (m, 2 H), 7.77 (s, 2 H), 7.79 (s, 1 H).

5 Step C: 2-(S)-(3,5-Bis(trifluoromethyl)benzyloxy)-3-(R)-phenyl-6-(R)-methyl morpholine

 A similar reaction was carried out using 55 mg (0.12 mmol) of 2-(S)-(3,5-bis(trifluoromethyl)benzyloxy)-3-(R)-phenyl-4-(2-propenyl)-6-(R)-methyl morpholine (from Example 23, Step A) and 111 mg (0.12 mmol) of tris(triphenylphosphine)rhodium chloride in 12 mL of 4:1 v/v acetonitrile/water. Flash chromatography on 4 g of silica gel using 50:1 v/v methylene chloride/acetonitrile as the eluant afforded 14 mg (28%) of 2-(S)-(3,5-bis(trifluoromethyl)benzyloxy)-3-(R)-phenyl-6-(R)-methyl morpholine as an oil.

Mass Spectrum (FAB): m/Z 420 (M+H, 90%)

15 ¹H NMR (CDCl₃, 400 MHz, ppm) 1.39 (d, 3 H, J = 6.8), 1.92 (br s, 1 H), 2.84 and 2.95 (dAB q, 2 H, J_{AB} = 12.8, J_{2.84} = 6.4, J_{2.95} = 3.6), 3.93 - 4.00 (m, 1 H), 4.07 (d, 1 H, J = 2.8), 4.68 and 4.95 (AB q, 2 H, J = 13.2), 4.93 (d, 1 H, J = 2.8), 7.28 - 7.37 (m, 3 H), 7.48 - 7.52 (m, 2 H), 7.55 (s, 2 H), 7.72 (s, 1 H).

20 EXAMPLE 24

25 2-(S)-(3,5-Bis(trifluoromethyl)benzyloxy)-3-(S)-phenyl-6-(S)-methyl morpholine and 2-(R)-(3,5-Bis-(trifluoromethyl)benzyloxy)-3-(S)-phenyl-6-(S)-methyl morpholine

 Substitution of 350 mg of N-(2-(S)-hydroxy-propyl)-N-(2-propenyl)-(S)-phenylglycinal, 3,5-bis-(trifluoromethyl)benzyl acetal (from Example 22) for N-(2-(R)-hydroxypropyl)-N-(2-propenyl)-(R)-phenyl-glycinal, 3,5-bis(trifluoromethyl)benzyl acetal in an experiment similar to the preceding example afforded 50 mg of 2-(S)-(3,5-bis(trifluoromethyl)benzyloxy)-3-(S)-phenyl-6-(S)-methyl morpholine and 14 mg of 2-(S)-(3,5-bis(trifluoromethyl)benzyloxy)-3-(S)-phenyl-6-(S)-methyl morpholine.

- 152 -

EXAMPLE 25

2-(R)-(3,5-Bis(trifluoromethyl)benzyloxy)-3-(S)-phenyl-6-(R)-methyl
morpholine and 2-(S)-(3,5-Bis-(trifluoromethyl)benzyloxy)-3-(S)-
5 phenyl-6-(R)-methyl morpholine

Step A: 2-(R)-(3,5-Bis(trifluoromethyl)benzyloxy)-3-(S)-phenyl-4-
(2-propenyl)-6-(R)-methyl morpholine and 2-(S)-(3,5-
10 bis(trifluoro-methyl)-benzyloxy)-3-(S)-phenyl-4-(2-pro-
penyl)-6-(R)-methyl morpholine

The title compounds were prepared in a manner similar to
Example 23, Step A. Cyclization of 300 mg (0.43 mmol) N-(2-(R)-
hydroxypropyl)-N-(prop-2-enyl)-(S)-phenylglycinal, 3,5-bis(trifluoro-
15 methyl)benzyl acetal (from Example 23) was effected using 246 mg
(1.29 mmol) of p-toluenesulfonic acid monohydrate and 5 mL of
toluene. Flash chromatography on 8 g of silica gel using 20:1 v/v
hexanes/ether as the eluant afforded 149 mg (75%) of the products as
inseparable diastereomers.

Mass Spectrum (FAB): m/Z 460 (M+H, 65%).

Step B: 2-(R)-(3,5-Bis(trifluoromethyl)benzyloxy)-3-(S)-phenyl-6-
(R)-methyl morpholine and 2-(S)-(3,5-Bis(trifluoro-
20 methyl)benzyloxy)-3-(S)-phenyl-6-(R)-methyl morpholine

A solution of 150 mg (0.33 mmol) of 2-(R)-(3,5-
25 bis(trifluoromethyl)benzyloxy)-3-(S)-phenyl-4-(2-propenyl)-6-(R)-
methyl morpholine and 2-(S)-(3,5-bis-(trifluoromethyl)-benzyloxy)-3-
(S)-phenyl-4-(2-propenyl)-6-(R)-methyl morpholine (from Example 25,
Step A) and 318 mg (0.32 mmol) of tris(triphenyl-phosphine)-rhodium
30 chloride in 20 mL of 4:1 v/v acetonitrile/water was heated at reflux for
1 h. Flash chromatography on 5 g of silica gel using 9:1 v/v
hexanes/ether as the eluant afforded 35 mg of the products as a mixture
and 26 mg of 2-(R)-(3,5-bis-(trifluoromethyl)benzyloxy)-3-(S)-phenyl-
6-(R)-methyl morpholine ($R_f = 0.22$ with 3:2 v/v hexanes/ether as the
eluant). Chromatography of the mixture on 5 g of silica gel using 20:1

- 153 -

v/v afforded 14 mg of 2-(S)-(3,5-bis(trifluoromethyl)benzyloxy)-3-(S)-phenyl-6-(R)-methyl morpholine (R_f = 0.14 with 3:2 v/v hexanes/ether as the eluant) and 17 mg of 2-(R)-(3,5-bis(trifluoromethyl)benzyloxy)-3-(S)-phenyl-6-(R)-methyl morpholine (41% total yield).

For the (2R,3S,6R) product:

Mass Spectrum (FAB): m/z 420 (M+H, 90%)

^1H NMR (CDCl_3 , 400 Mhz. ppm) 1.30 (d, 3 H, J = 6.4), 1.74 (br s, 1 H), 2.73 and 2.98 (dAB q, 2 H, J_{AB} = 11.6, $J_{2.73}$ = 10.0, $J_{2.98}$ = 2.4), 3.65 (d, 1 H, J = 7.2), 3.89 - 3.94 (m, 1 H), 4.45 (d, 1 H, J = 7.2), 4.53 and 4.90 (AB q, 2 H, J = 13.2), 7.28 - 7.38 (m, 3 H), 7.41 - 7.43 (m, 2 H), 7.45 (s, 2 H), 7.70 (s, 1 H).

For the (2S,3S,6R) product:

Mass Spectrum (FAB): m/z 420 (M+H, 90%)

^1H NMR (CDCl_3 , 400 Mhz. ppm) 1.20 (d, 3 H, J = 6.4), 2.04 (br s, 1 H), 2.84 and 3.15 (dAB q, 2 H, J_{AB} = 12.8, $J_{2.84}$ = 10.8, $J_{3.15}$ = 2.8), 4.08 (d, 1H, J = 2.8), 4.08 - 4.15 (m, 1 H), 4.53 and 4.80 (AB q, 2 H, J = 13.2), 4.79 (d, 1 H, J = 2.8), 7.28 - 7.38 (m, 5 H), 7.43 (s, 2 H), 7.70 (s, 1 H).

EXAMPLE 26

2-(S)-(3,5-Bis(trifluoromethyl)benzyloxy)-3-(R)-phenyl-6-(S)-methyl morpholine and 2-(R)-(3,5-Bis-(trifluoromethyl)benzyloxy)-3-(R)-phenyl-6-(S)-methyl morpholine

Substitution of 250 mg of N-(2-(S)- hydroxy-propyl)-N-(2-propenyl)-(S)-phenylglycinal, 3,5-bis-(trifluoromethyl)benzyl acetal (from Example 22) for N-(2-(R)-hydroxypropyl)-N-(2-propenyl)-(R)-phenyl-glycinal, 3,5-bis(trifluoromethyl)benzyl acetal in an experiment similar to the preceding example afforded 42 mg of 2-(S)-(3,5-bis-(trifluoromethyl)benzyloxy)-3-(R)-phenyl-6-(S)-methyl morpholine and 17 mg of 2-(S)-(3,5-bis(trifluoro- methyl)benzyloxy)-3-(R)-phenyl-6-(S)-methyl morpholine.

- 154 -

EXAMPLE 27

5 2-(R)-(3,5-Bis(trifluoromethyl)benzyloxy)-3-(S)-phenyl-5-(R)-methyl morpholine, 2-(S)-(3,5-Bis(trifluoromethyl)benzyloxy)-3-(S)-phenyl-5-(R)-methyl morpholine, 2-(R or S)-(3,5-Bis(trifluoromethyl)benzyloxy)-3-(R)-phenyl-5-(R)-methyldmorpholine, and 2-(S or R)-(3,5-Bis(trifluoromethyl) benzyloxy)-3-(R)-phenyl-5-(R)-methyldmorpholine

10 Execution of the sequence described in Example 19 substituting (R)-2-amino-1-propanol for (R)-1-amino-2-propanol provided a mixture of 55 mg of high R_f material and 56 mg of low R_f material. The high R_f material was processed according to Example 23, Step A above to provide 10 mg of high R_f material (2-(R)-(3,5-Bis(trifluoromethyl)benzyloxy)-3-(S)-phenyl-5-(R)-methyl morpholine and 7 mg of low R_f material (2-(S)-(3,5-Bis(trifluoromethyl)benzyloxy)-3-(S)-phenyl-5-(R)-methyl morpholine. The low R_f material (after being combined with an additional 30 mg of material) was processed according to Example 23, Step A to provide 24 mg of high R_f material (2-(R or S)-(3,5-Bis(trifluoromethyl)benzyloxy)-3-(R)-phenyl-5-(R)-methyl-morpholine and 18 mg of low R_f material (2-(S or R)-(3,5-Bis(trifluoromethyl)benzyloxy)-3-(R)-phenyl-5-(R)-methyldmorpholine.

25 2-(R)-(3,5-Bis(trifluoromethyl)benzyloxy)-3-(S)-phenyl-5-(R)-methyl morpholine

Mass Spectrum (FAB): m/Z 420 (M+H, 100%), 227 (50%), 192 (75%), 176 (65%).

30 NMR (CDCl₃, 400 MHz, ppm): δ 0.98 (d, 3H, J= 6.3 Hz), 3.16-3.20 (m, 1H), 3.43-3.47 (m, 1H), 3.79 (d, 1H, J= 7.5 Hz), 3.91 (dd, 1H, J= 3.2 & 11.5 Hz), 4.51 (d, 2H, J= 13.4 Hz), 4.85 (d, 1H, J= 13.2 Hz), 7.29-7.45 (m, 7H), 7.67 (s, 1H).

2-(S)-(3,5-Bis(trifluoromethyl)benzyloxy)-3-(S)-phenyl-5-(R)-methyl morpholine

- 155 -

Mass Spectrum (FAB): m/Z 420 (M+H, 48%), 227 (35%), 192 (39%), 176 (100%).

5 NMR (CDCl₃, 400 MHz, ppm): δ 1.10 (d, 3H, J= 6.4 Hz), 3.23-3.26 (m, 1H), 3.56-3.61 (m, 2H), 4.17 (d, 1H, J= 2.3 Hz), 4.51 (d, 1H, J= 13.7 Hz), 4.71 (d, 1H, J= 2.4 Hz), 4.78 (d, 1H, J= 13.5 Hz), 7.28-7.39 (m, 7H), 7.68 (s, 1H).

2-(R or S)-(3,5-Bis(trifluoromethyl)benzyloxy)-3-(R)-phenyl-5-(R)-methyl morpholine

10 Mass Spectrum (FAB): m/Z 281 (35%), 221 (55%), 207 (45%), 192 (40%), 147 (100%).

15 NMR (CDCl₃, 400 MHz, ppm): δ 1.13 (d, 3H, J= 6.6 Hz), 3.10-3.14 (m, 1H), 3.66 (dd, 1H, J= 6.6 & 11.4 Hz), 3.76 (dd, 1H, J= 3.5 & 11.2 Hz), 4.04 (d, 1H, J= 4.0 Hz), 4.61 (d, 1H, J= 13.2 Hz), 4.74 (d, 1H, J= 3.9 Hz), 4.89 (d, 1H, 13.2 Hz), 7.26-7.35 (m, 3H), 7.47-7.49 (m, 2H), 7.64 (s, 1H), 7.74 (s, 1H).

2-(R or S)-(3,5-Bis(trifluoromethyl)benzyloxy)-3-(R)-phenyl-5-(R)-methyl morpholine

20 NMR (CDCl₃, 400 MHz, ppm): δ 1.36 (d, 3H, J= 6.7 Hz), 3.27-3.31 (m, 1H), 3.39 (dd, 1H, J= 2.2 & 11.3 Hz), 4.16 (dd, 1H, J= 3.2 & 11.0 Hz), 4.37 (d, 1H, J= 2.3 Hz), 4.53 (d, 1H, J= 13.5 Hz), 4.75 (d, 1H, J= 2.5 Hz), 4.81 (d, 1H, 13.6 Hz), 7.26-7.35 (m, 3H), 7.26-7.43 (m, 7H), 7.68 (s, 1H).

25

EXAMPLE 28

2-(R or S)-(3,5-Bis(trifluoromethyl)-benzyloxy)-3-(S)-phenyl-5-(S)-methylmorpholine, 2-(S or R)-(3,5-Bis(trifluoromethyl)benzyloxy)-3-(S)-phenyl-5-(S)-methyl-morpholine, and 2-(R)-(3,5-Bis(trifluoromethyl)benzyl-oxy)-3-(R)-phenyl-5-(S)-methylmorpholine

30 Execution of the sequence described in Example 19 substituting (S)-2-amino-1-propanol for (R)-1-amino-2-propanol provided a mixture of 78 mg of high R_f material and 70 mg of low R_f

- 156 -

material. The high R_f material was processed according to Example 23, Step A above to provide less than 1 mg of high R_f material (2-(R)-(3,5-Bis(trifluoromethyl)benzyloxy)-3-(S)-phenyl-5-(S)-methylmorpholine) and 9 mg of low R_f material (2-(S)-(3,5-Bis(trifluoromethyl)benzyloxy)-3-(S)-phenyl-5-(S)-methyl morpholine. The low R_f material was processed according to Example 23, Step A to provide 20 mg of high R_f material (2-(R or S)-(3,5-Bis(trifluoromethyl)benzyloxy)-3-(S)-phenyl-5-(S)-methylmorpholine and 14 mg of low R_f material (2-(S or R)-(3,5-Bis(trifluoromethyl)benzyloxy)-3-(S)-phenyl-5-(S)-methylmorpholine.

2-(R or S)-(3,5-Bis(trifluoromethyl)benzyloxy)-3-(S)-phenyl-5-(S)-methyl morpholine

Mass Spectrum (FAB): m/Z 420 (M+H, 60%), 227 (68%), 192 (56%), 176 (100%).

NMR (CDCl₃, 400 MHz, ppm): δ 1.12 (d, 3H, J= 6.6 Hz), 3.09-3.14 (m, 1H), 3.65 (dd, 1H, J= 6.6 & 11.0 Hz), 3.75 (dd, 1H, J= 3.6 & 11.1 Hz), 4.04 (d, 1H, J= 3.9 Hz), 4.61 (d, 1H, J= 13.2 Hz), 4.73 (d, 1H, J= 3.9 Hz), 4.89 (d, 1H, 13.2 Hz), 7.28-7.35 (m, 3H), 7.47 (d, 2H, 7.0 Hz), 7.64 (s, 1H), 7.74 (s, 1H).

2-(S or R)-(3,5-Bis(trifluoromethyl)benzyloxy)-3-(S)-phenyl-5-(S)-methyl morpholine

Mass Spectrum (FAB): m/Z 420 (M+H, 50%), 227 (45%), 192 (40%), 176 (100%).

NMR (CDCl₃, 400 MHz, ppm): δ 1.36 (d, 3H, J= 6.9 Hz), 3.27-3.29 (m, 1H), 3.39 (dd, 1H, J= 2.2 & 11.1 Hz), 4.15 (dd, 1H, J= 3.3 & 11.1 Hz), 4.37 (d, 1H, J= 2.5 Hz), 4.52 (d, 1H, J= 13.3 Hz), 4.75 (d, 1H, J= 2.4 Hz), 4.81 (d, 1H, 13.5 Hz), 7.28-7.43 (m, 7H), 7.68 (s, 1H).

2-(R)-(3,5-Bis(trifluoromethyl)benzyloxy)-3-(R)-phenyl-5-(S)-methyl morpholine

NMR (CDCl₃, 400 MHz, ppm): δ 1.10 (d, 3H, J= 6.4 Hz), 3.22-3.25 (m, 1H), 3.55-3.60 (m, 2H), 4.17 (d, 1H, J= 2.3 Hz), 4.51 (d, 1H, J=

- 157 -

13.5 Hz), 4.71 (d, 1H, J= 2.4 Hz), 4.77 (d, 1H, J= 13.6 Hz), 7.28-7.38 (m, 7H), 7.67 (s, 1H).

EXAMPLE 29

5

2-(R)-(3,5-Bis(trifluoromethyl)benzyloxy)-3-(S)-phenyl-5-(R)-phenylmorpholine, 2-(S)-(3,5-Bis(tri-fluoromethyl)benzyloxy)-3-(S)-phenyl-5-(R)-phenyl-morpholine, and 2-(R or S)-(3,5-Bis(trifluoromethyl)-benzyloxy)-3-(R)-phenyl-5-(R)-phenylmorpholine

10

Execution of the sequence described in Example 19

substituting (R)-2-amino-2-phenylethanol for (R)-1-amino-2-propanol provided a mixture of 62 mg of high R_f material and 52 mg of low R_f material. The high R_f material was processed according to Example 23, Step A above to provide 16 mg of high R_f material (2-(R)-(3,5-Bis(trifluoromethyl)benzyloxy)-3-(S)-phenyl-5-(R)-phenylmorpholine and 4 mg of low R_f material (2-(S)-(3,5-Bis(trifluoromethyl)benzyloxy)-3-(S)-phenyl-5-(R)-phenylmorpholine . The low R_f material was processed according to Example 23, Step A to provide 4 mg of product (2-(R or S)-(3,5-Bis(trifluoromethyl)benzyloxy)-3-(R)-phenyl-5-(R)-phenylmorpholine.

15

20

2-(R)-(3,5-Bis(trifluoromethyl)benzyloxy)-3-(S)-phenyl-5-(R)-phenylmorpholine

25

NMR (CDCl₃, 400 MHz, ppm): δ 3.62 (t, 1H, J= 10.7 & 21.5 Hz), 3.93 (d, 1H, J= 7.4 Hz), 3.99 (dd, 1H, J= 3.1 & 11.2 Hz), 4.18 (dd, 1H, J= 3.0 & 10.2 Hz), 4.46 (d, 1H, J= 7.4 Hz), 4.53 (d, 1H, J= 13.5 Hz), 4.89 (d, 1H, J= 13.3 Hz), 7.28-7.55 (m, 12H), 7.69 (s, 1H).

30

2-(S)-(3,5-Bis(trifluoromethyl)benzyloxy)-3-(S)-phenyl-5-(R)-phenylmorpholine

NMR (CDCl₃, 400 MHz, ppm): δ 3.67 (dd, 1H, J= 3.5 & 11.0 Hz), 3.89 (d, 1H, J= 10.8 & 21.6 Hz), 4.25 (dd, 1H, J= 3.3 & 11.0 Hz), 4.34 (d, 1H, J= 2.2 Hz), 4.52 (d, 1H, J= 13.8 Hz), 4.78-4.87 (m, 2H), 7.28-7.51 (m, 12H), 7.69 (s, 1H).

- 158 -

2-(R or S)-(3,5-Bis(trifluoromethyl)benzyloxy)-3-(R)-phenyl-5-(R)-phenylmorpholine

5 NMR (CDCl₃, 400 MHz, ppm): δ 4.10-4.25 (m, 2H), 4.30-4.38 (m, 1H), 4.48-4.54 (m, 1H), 4.59-4.66 (m, 1H), 4.86-5.00 (m, 2H), 7.25-7.74 (m, 13H).

EXAMPLE 30

10 2-(S)-(3,5-Bis(trifluoromethyl)benzyloxy)-3-(R)-phenyl-5-(S)-phenylmorpholine, 2-(R)-(3,5-Bis(trifluoromethyl)benzyloxy)-3-(R)-phenyl-5-(S)-phenyl-morpholine, 2-(R or S)-(3,5-Bis(trifluoromethyl)benzyloxy)-3-(S)-phenyl-5-(S)-phenyl-morpholine, and 2-(R or S)-(3,5-Bis(trifluoromethyl)benzyloxy)-3-(S)-phenyl-5-(S)-phenylmorpholine

15 Execution of the sequence described in Example 19 substituting (S)-2-amino-2-phenylethanol for (R)-1-amino-2-propanol provided a mixture of 75 mg of high R_f material and 64 mg of low R_f material. The high R_f material was processed according to Example 23, Step A above to provide 23 mg of high R_f material (2-(S)-(3,5-Bis(trifluoromethyl)benzyloxy)-3-(R)-phenyl-5-(S)-phenylmorpholine [L-740, 930]) and 7 mg of low R_f material (2-(R)-(3,5-Bis(trifluoromethyl)benzyloxy)-3-(R)-phenyl-5-(S)-phenylmorpholine. The low R_f material was processed according to Example 23, Step A to provide 26 mg of higher R_f material (2-(R or S)-(3,5-Bis(trifluoromethyl)benzyloxy)-3-(S)-phenyl-5-(S)-phenylmorpholine and 6 mg of lower R_f material (2-(R or S)-(3,5-Bis(trifluoromethyl)benzyloxy)-3-(S)-phenyl-5-(S)-phenylmorpholine.

30 2-(S)-(3,5-Bis(trifluoromethyl)benzyloxy)-3-(R)-phenyl-5-(S)-phenyl-morpholine

NMR (CDCl₃, 400 MHz, ppm): δ 3.60-3.74 (m, 1H), 3.94 (d, 1H, J= 7.6 Hz), 4.00 (dd, 1H, J= 3.2 & 11.3 Hz), 4.18-4.21 (m, 1H), 4.50-4.55 (m, 2H), 4.89 (m, 1H), 7.26-7.55 (m, 12H), 7.69 (s, 1H).

- 159 -

2-(R)-(3,5-Bis(trifluoromethyl)benzyloxy)-3-(R)-phenyl-5-(S)-phenyl-morpholine

5 NMR (CDCl₃, 400 MHz, ppm): δ 3.68 (dd, 1H, J= 3.0 & 11.0 Hz), 3.88-3.94 (m, 1H), 4.26-4.30 (m, 1H), 4.36 (s, 1H), 4.52 (d, 1H, J= 13.5 Hz), 4.77-4.86 (m, 2H), 7.27-7.51 (m, 12H), 7.69 (s, 1H).

2-(R or S)-(3,5-Bis(trifluoromethyl)benzyloxy)-3-(S)-phenyl-5-(S)-phenylmorpholine

10 NMR (CDCl₃, 400 MHz, ppm): δ 3.93-3.95 (m, 1H), 4.06-4.21 (m, 2H), 4.38-4.42 (m, 1H), 4.59-4.68 (m, 2H), 4.83-4.94 (m, 2H), 7.25-7.81 (m, 13H).

2-(R or S)-(3,5-Bis(trifluoromethyl)benzyloxy)-3-(S)-phenyl-5-(S)-phenylmorpholine

15 NMR (CDCl₃, 400 MHz, ppm): δ 3.43-3.59 (m, 2H), 3.82 (d, 1H, J= 7.2 Hz), 4.25 (d, 1H, J= 12.5 Hz), 4.52-4.63 (m, 3H), 4.80-4.90 (br s, 1H), 7.11-7.81 (m, 13H).

20 EXAMPLE 31

2-(S)-(3,5-bis(trifluoromethyl)benzyloxy)-6-(R)-methyl-3-(S)-phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine

25 According to the procedure given in Example 17, Step B, 98 mg (0.24 mmole) of 2-(S)-(3,5-bis-(trifluoromethyl)benzyloxy)-3-(S)-phenyl-6-(R)-methyl morpholine (from Example 25 above), 38 mg (0.28 mmole) of N-formyl-2-chloroacetamidrazone (from Example 17, Step A above) and 97 mg (0.7 mmole) of anhydrous potassium carbonate gave, after flash chromatography on 28 g of silica eluting with 1 L of 100:4:0.5 methylene chloride:methanol: ammonia water, a
30 light yellow solid which after recrystallization from hexanes/methylene chloride provided 77 mg (66%) of 2-(S)-(3,5-bis(trifluoromethyl)benzyloxy)-6-(R)-methyl-3-(S)-phenyl-4-(3-(1,2,4-triazolo)methyl)morpholine as a white powder.

- 160 -

NMR (CDCl₃, 400 MHz, ppm): δ 1.17 (d, J= 6.3, 3H), 2.29 (t, J= 11.1, 1H), 2.92 (d, J= 11.1, 1H), 3.42 (d, J= 15.3, 1H), 3.58 (s, 1H), 3.88 (d, J= 15.4, 1H), 4.20-4.33 (m, 1H), 4.43 (d, 13.5, 1H), 4.71 (d, J= 2.4, 1H), 4.74 (d, J= 13.3, 1H), 7.30-7.55 (m, 7H), 7.69 (s, 1H), 7.95 (s, 1H).

EXAMPLE 32

2-(S)-(3,5-Bis(trifluoromethyl)benzyloxy)-6-(R)-methyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-3-(S)-phenylmorpholine

A mixture of 96 mg (0.23 mmole) of 2-(S)-(3,5-bis-(trifluoromethyl)benzyloxy)-3-(S)-phenyl-6-(R)-methyl morpholine (from Example 25 above), 46 mg (0.28 mmole) of N-methylcarboxy-2-chloroacetamidrazone and 95 mg (0.69 mmole) of anhydrous potassium carbonate in 3 mL of dry DMF was stirred at room temperature for 20 min, at 60°C for 90 min and then at 120°C for 2 hr. The mixture was cooled to room temperature, taken up in 15 mL of ethyl acetate and was washed with 3x10 mL of water. The combined aqueous layers were back-extracted with 10 mL of ethyl acetate, the combined organic layers were washed with 10 mL of brine, dried over sodium sulfate, filtered and concentrated in vacuo. The residue was purified by flash chromatography on 28 g of silica eluting with 1L of 100:4 methylene chloride: methanol to give 65 mg (55%) of 2-(S)-(3,5-bis(trifluoromethyl)benzyl-oxy)-6-(R)-methyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)-methyl)-3-(S)-phenylmorpholine as a light yellow powder.

NMR (CDCl₃, 400 MHz, ppm): δ 1.18 (d, J= 6.2, 3H), 2.15 (t, J= 11.1, 1H), 2.89 (d, J= 14, 2H), 3.49 (d, J= 2.2, 1H), 3.61 (d, J= 14.4, 1H), 4.20-4.30 (m, 1H), 4.45 (d, J= 13.6, 1H), 4.67 (d, J= 2.5, 1H), 4.79 (d, J= 13.5, 1H), 7.25-7.50 (m, 7H), 7.62 (s, 1H), 10.07 (s, 1H), 10.35 (s, 1H).

EXAMPLE 33

2-(S)-(3,5-Bis(trifluoromethyl)benzyloxy)-3-(R)-phenylmorpholine

- 161 -

Step A: 4-Benzyl-2-(S)-hydroxy-3-(R)-phenylmorpholine

A solution of 3.72 g (13.9 mmol) of 4-benzyl-3-(R)-phenyl-2-morpholinone, prepared from (R)-phenyl-glycine as described in Example 14, in 28 mL of CH₂Cl₂ was cooled in a -78°C bath under a N₂ atmosphere and 14 mL of a 1.5 M solution of DIBAL-H (21 mmol) in toluene were added. After stirring the resulting solution for 0.5 h, it was allowed to warm to -50°C and maintained at this temperature for 0.5 h. The reaction mixture was quenched by adding 10 mL of aqueous potassium sodium tartarate. The mixture was diluted with CH₂Cl₂ and the layers were separated. The aqueous layer was extracted 3 times with CH₂Cl₂. The CH₂Cl₂ layers were washed with brine, dried over Na₂SO₄ and filtered. Concentration of the filtrate furnished 3.32 g (88%) of 4-benzyl-2-(S)-hydroxy-3-(R)-phenylmorpholine suitable for use in the next step.

NMR (CDCl₃) 2.28 (m, 1H), 2.71 (m, 1H), 2.91 (d, J = 13 Hz, 1H), 3.09 (d, J = 6 Hz, 1H), 3.69 (d, J = 13 Hz, 1H), 3.82 (td, J = 10 Hz and 2 Hz, 1H), 3.91 (d, J = 10 Hz, 1H), 4.73 (t, J = 6 Hz, 1H), 7.2-7.52 (m, 10H).

Step B: 4-Benzyl-2-(S)-(3,5-bis(trifluoromethyl)-benzyloxy)-3-(R)-phenylmorpholine

To a suspension of 0.592 g (14.8 mmol) of NaH in 30 mL of dry THF at 0°C was added 3.32 g (12.3 mmol) of 4-benzyl-2-(S)-hydroxy-3-(R)-phenyl-morpholine prepared in step A. After 15 min 0.915 g of tetrabutylammonium iodide (2.47 mmol) and 2.4 mL (13 mmol) of 3,5-bis(trifluoromethyl)benzyl bromide were added. The resulting mixture was stirred at ice-bath temperature for 1 h, then poured into saturated NaHCO₃ solution and extracted with ethyl acetate (EtOAc). The organic layers were combined, washed with brine, dried over Na₂SO₄ and filtered. The filtrate was concentrated in vacuo and the residue was chromatographed on a Waters Prep500 HPLC system using 50% EtOAc/Hexane to isolate 3.6 g (59%) of 4-Benzyl-2-(S)-(3,5-bis(trifluoromethyl)benzyloxy)-3-(R)-phenylmorpholine.

- 162 -

1H NMR (CDCl₃) 2.3 (td, J = 11 Hz and 3 Hz, 1H), 2.71 (d, J = 11 Hz, 1H), 2.90 (d, J = 13 Hz, 1H), 3.22 (d, J = 7.3 Hz, 1H), 3.75 (m, 2H), 3.93 (m, 1H), 4.43 (d, J = 13 Hz, 1H), 4.45 (d, J = 7.3 Hz, 1H), 4.82 (d, J = 13 Hz, 1 H), 7.19-7.5 (m, 12H), 7.67 (s, 1H).

Step C: 2-(S)-(3,5-Bis(trifluoromethyl)benzyloxy)-3-(R)-phenylmorpholine

A solution of 3.6 g (7.27 mmol) of 4-benzyl-2-(S)-(3,5-bis(trifluoromethyl)benzyloxy)-3-(R)-phenylmorpholine in 100 mL of ethanol and 5 mL of water, containing 0.72 g of 10% Pd/C was hydrogenated on a Parr apparatus for 36 h. The catalyst was filtered and thoroughly washed with EtOAc. The filtrate was concentrated and the residue was partitioned between water and EtOAc. The EtOAc layer was washed with brine, dried over Na₂SO₄, filtered and concentrated. The residue was purified by flash chromatography using a gradient of 10-60 % EtOAc/hexane to isolate 2.05 g (70%) of 2-(S)-(3,5-bis(trifluoromethyl)benzyloxy)-3-(R)-phenylmorpholine. 1H NMR (CDCl₃) 1.92 (br s, 1H), 2.91 (m, 1H), 3.05 (td, J = 11 Hz and 3 Hz, 1H), 3.68 (d, J = 7 Hz, 1H), 3.81 (td, J = 11 Hz and 3 Hz, 1H), 4.01 (m, 1H), 4.44 (d, J = 7 Hz), 4.5 (d, J = 13 Hz, 1H), 4.85 (d, J = 13 Hz, 1 H), 7.28-7.42 (m, 7H), 7.67 (s, 1H).

EXAMPLE 34

4-(3-(1,2,4-Triazolo)methyl)-2-(S)-(3,5-bis(trifluoromethyl)benzyloxy)-3-(R)-phenylmorpholine

The title compound was prepared by the procedure of Example 17, step B employing the product of Example 33, step C as a starting material.

1H NMR (CDCl₃) 1.75 (br s, 1 H), 2.61 (td, J = 12 Hz and 2 Hz, 1H), 2.83 (d, J = 12 Hz, 1H), 3.33 (d, J = 7 Hz, 1H), 3.48 (d, J = 15 Hz, 1H), 3.78 (d, J = 15 Hz, 1H), 3.85 (m, 1H), 3.99 (m, 1H), 4.44 (d, J = 13 Hz, 1 H), 4.49 (d, J = 7 Hz, 1H), 4.81 (d, J = 13 Hz, 1H), 7.23-7.45 (m, 7H), 7.67 (s, 1H), 7.96 (s, 1H).

- 163 -

EXAMPLE 35

4-(3-(5-Oxo-1H,4H-1,2,4-triazolo)methyl)-2-(S)-(3,5-bis-(trifluoro-
methyl)benzyloxy)-3-(R)-phenyl-morpholine

The title compound was prepared by the procedure of Example 18, steps B & C employing the product of Example 33, step C as a starting material.

EXAMPLE 36

4-(2-(Imidazolo)methyl)-2-(S)-(3,5-bis(trifluoro-methyl)benzyloxy)-3-(S)-phenylmorpholine

A solution of 101 mg (0.25 mmol) of 2-(S)-(3,5-bis(trifluoromethyl)benzyloxy)-3-(S)-phenylmorpholine (Example 15), 98 mg (1.0 mmol) of imidazole-2-carboxaldehyde, and 5 drops of glacial acetic acid in 3 ml of methanol was treated with 1.5 ml of 1M sodium cyanoborohydride solution in THF. After 16 hr, the reaction was quenched with 5 ml of saturated aqueous sodium bicarbonate solution and partitioned between 40 ml of ethyl acetate and 20 ml of water. The organic layer was separated, dried over magnesium sulfate, and concentrated in vacuo. Flash chromatography on 8 g of silica gel using 50:1:0.1 methylene chloride/methanol/amonium hydroxide as the eluent afforded 54 mg (44% yield) of the title compound as a white solid.

¹H NMR (CDCl₃) 2.60 (dt, J = 3.2 Hz and 12.4 Hz, 1H), 2.85 (d, J = 12.4 Hz, 1H), 3.28 (d, J = 14.4 Hz, 1H), 3.59 (d, J = 2.8 Hz, 1H), 3.66 (dd, J = 2.0, 11.6 Hz, 1H), 3.84 (d, J = 14.4 Hz, 1H), 3.94 (app s, 2H), 4.14 (dt, J = 2.0, 12.0 Hz, 1H), 4.43 (d, J = 13.6 Hz, 1H), 4.71 (d, J = 2.8 Hz, 1H), 4.78 (d, J = 13.6 Hz, 1H), 6.99 (app s, 2H), 7.25-7.48 (m, 6H), 7.72 (s, 1H). Mass spectrum (FAB): m/z 486 (100%, M+H)

EXAMPLE 37

4-(2-(Imidazolo)methyl)-2-(S)-(3,5-bis(trifluoro-methyl)benzyloxy)-3-(R)-phenylmorpholine

- 164 -

The title compound was prepared by the procedure of Example 36 employing appropriate starting materials.

¹H NMR (CDCl₃) 2.53 (td, J = 11 Hz and 3 Hz, 1H), 2.74 (d, J = 12 Hz, 1H), 3.23 (d, J = 7 Hz, 1H), 3.32 (d, J = 15 Hz, 1H), 3.66 (d, J = 15 Hz, 1H), 3.77 (td, J = 11 Hz and 2 Hz, 1H), 3.99 (m, 1H), 4.44 (m, 2H), 4.8 (d, J = 13 Hz, 1H), 6.94 (s, 2H), 7.2-7.45 (m, 7H), 7.67 (s, 1H).

EXAMPLE 38

4-(5-(Imidazolo)methyl)-2-(S)-(3,5-bis(trifluoro-methyl)benzyloxy)-3-(R)-phenylmorpholine

The title compound was prepared by the procedure of Example 36 employing appropriate starting materials.

¹H NMR (CDCl₃) 2.47 (td, J = 12 Hz and 3 Hz, 1H), 2.83 (d, J = 12 Hz, 1H), 3.2 (m, 2H), 3.61 (d, J = 14 Hz, 1H), 3.79 (td, J = 12 Hz and 2 Hz, 1H), 3.96 (m, 1H), 4.44 (m, 2H), 4.80 (d, J = 13 Hz, 1H), 6.81 (s, 1H), 7.28-7.45 (m, 7H), 7.60 (s, 1H), 7.66 (s, 1H).

EXAMPLE 39

4-(Aminocarbonylmethyl)-2-(S)-(3,5-bis(trifluoro-methyl)benzyloxy)-3-(R)-phenylmorpholine

The title compound was prepared by the procedure of Example 15 employing appropriate starting materials.

¹H NMR (CDCl₃) 2.54 (td, J = 11 Hz and 2 Hz, 1H), 2.64 (d, J = 17 Hz, 1H), 2.93 (d, J = 12 Hz, 1H), 3.14 (d, J = 17 Hz, 1H), 3.27 (d, J = 7 Hz, 1H), 3.83 (td, J = 11 Hz and 2 Hz, 1H), 4.05 (m, 1H), 4.46 (m, 2H), 4.81 (d, J = 13 Hz, 1H), 5.62 (br s, 1H), 6.80 (br s, 1H), 7.28-7.32 (m, 7H), 7.67 (s, 1H).

EXAMPLES 40-43

4-(3-(1,2,4-Triazolo)methyl)-2-(3-(tert-butyl)-5-methylbenzyloxy)-3-phenyl-morpholine, 4-(3-(5-Oxo-1H,4H-1,2,4-triazolo)methyl)-2-(3-

- 165 -

(tert-butyl)-5-methylbenzyloxy)-3-phenyl-morpholine, 4-(2-(Imidazolo)methyl)-2-(3-(tert-butyl)-5-methyl-benzyloxy)-3-phenyl-morpholine, 4-(4-(Imidazolo)-methyl)-2-(3-(tert-butyl)-5-methyl-benzyloxy)-3-phenyl-morpholine

The title compounds are each prepared by the procedures of Examples 15, 17 & 18 employing appropriately substituted starting materials and reagents.

EXAMPLE 44

2-(S)-(3,5-Dichlorobenzyloxy)-3-(S)-phenylmorpholine

Step A: 3,5-Dichlorobenzyl alcohol, trifluoromethanesulfonate ester

A solution of 6.09 g (34.4 mmole) of 3,5-dichlorobenzyl alcohol and 8.48 g (41.3 mmole) of 2,6-di-t-butyl-4-methylpyridine in 280 mL of dry carbon tetrachloride under a nitrogen atmosphere was treated with 5.95 mL (35.4 mmole) of trifluoromethanesulfonic anhydride at room temperature. A white precipitate formed shortly after the addition of the anhydride. After 90 min, the slurry was filtered under nitrogen with a Schlenk filter, and the filtrate was concentrated in vacuo. The residue, which was a two-phase oil, was dissolved under nitrogen in 60 mL of dry toluene. The resulting solution was used immediately in Step B below.

Step B: 4-Benzyl-2-(S)-(3,5-dichlorobenzyloxy)-3-(S)-phenyl-morpholine

A solution of 5.11 g (19.1 mmole) of N-benzyl-3-(S)-phenylmorpholin-2-one (from Example 14) in 100 mL of dry THF was cooled to -75°C under nitrogen and was treated dropwise with 20.5 mL (20.5 mmole) of a 1M solution of lithium tri(sec-butyl)borohydride (L-Selectride®) in THF. After stirring the solution at -75°C for 30 min, a solution of 3,5-dichlorobenzyl alcohol, trifluoromethanesulfonate ester in toluene (from Example 44, Step A) was added by cannula so that the

- 166 -

internal temperature was maintained below -60°C. The resulting solution was stirred between -38°C and -50°C for 9 hr, and was then treated with 14 mL of aqueous ammonia and stored at -20°C for 12 hours. The solution was then poured into a mixture of 50 mL of ethyl acetate and 100 mL of water, and the layers were separated. The aqueous phase was extracted with 2x100 mL of ethyl acetate, each extract was washed with brine, the combined organic layers were dried over sodium sulfate, the mixture was filtered and the filtrate concentrated in vacuo. The residue was purified by flash chromatography on 235 g of silica eluting with 1.5 L of 100:2 hexanes:ethyl acetate, then 1.5 L of 100:3 hexanes:ethyl acetate and then 1.9 L of 100:5 hexanes:ethyl acetate to give 4.4 g (54%) of an oil, which by ¹H NMR is a 8:1 mixture of cis:trans morpholines.

Mass Spectrum (FAB): m/Z 430,428,426 (M+H, ~60%), 268 (M-ArCH₂, 100%), 252 (M-ArCH₂O, 75%), 222(20%), 159 (45%).

¹H NMR (CDCl₃, 400 MHz, ppm): δ major (cis) isomer: 2.32 (td, J= 12, 3.6, 1H), 2.84 (app t, J= 13, 2H), 3.52 (d, J= 2.6, 1H), 3.55 (dq, J= 11.3, 1.6, 1H), 3.91 (d, J= 13.3, 1H), 4.12 (td, J= 11.6, 2.4, 1H), 4.29 (d, J= 13.6, 1H), 4.59 (d, J= 2.9, 1H), 4.60 (d, J= 13.6), 6.70 (s, 2H), 7.13 (t, J= 1.9, 1H), 7.2-7.6 (m, 8H), 7.53 (br d, 2H).

Step C: 2-(S)-(3,5-Dichlorobenzyloxy)-3-(S)-phenyl-morpholine

A solution of 0.33 g (0.77 mmole) of 4-benzyl-2-(S)-(3,5-dichlorobenzyloxy)-3-(S)-phenylmorpholine (from Example 44, Step B) and 0.22 g (1.54 mmole) of 1-chloroethyl chloroformate in 4.5 mL of 1,2-dichloroethane was placed in a pressure vial which was lowered into an oil bath which was heated to 110°C. After stirring for 60 hr the solution was cooled and concentrated in vacuo. The residue was dissolved in 7 mL of methanol and the resulting solution was heated at reflux for 30 min. The mixture was cooled and treated with several drops of concentrated aqueous ammonia and the solution was concentrated. The residue was partly purified by flash chromatography

- 167 -

on 67 g of silica eluting with 1.5 L of 100:1 methylene chloride:
methanol, and the rich cuts were purified by flash chromatography on
32 g of silica eluting with 50:50 hexanes: ethyl acetate and then 50:50:5
hexanes:ethyl acetate:methanol to give 0.051 g (20%) of an oil, which
5 by ¹H NMR was pure cis morpholine.

Mass Spectrum (FAB): m/Z 468,466,464 (max 8%), 338,340 (M+H,
25%), 178 (20%), 162 (100%), 132 (20%),.

10 ¹H NMR (CDCl₃, 400 MHz, ppm): δ 1.89 (br s, 1H), 3.08 (dd, J= 12.5,
2.9, 1H), 3.23 (td, J= 12.2, 3.6, 1H), 3.59 (dd, J= 11.3, 2.5, 1H), 4.03
(td, J= 11.7, 3, 1H), 4.09 (d, J= 2.4, 1H), 4.37 (d, J= 13.5, 1H), 4.62 (d,
J= 13.3, 1H), 4.67 (d, J= 2.5, 1H), 6.72 (d, J= 1.8, 2H), 7.14 (t, J= 1.8,
1H), 7.25-7.40 (m, 5H).

15

EXAMPLE 45

20 2-(S)-(3,5-dichlorobenzyloxy)-4-(3-(5-oxo-1,2,4-triazolo)methyl)-3-
(S)-phenylmorpholine

Step A: N-Methylcarboxy-2-chloroacetamidrazone

A solution of 5.0 g (66.2 mmol) of chloroacetonitrile in 35
mL of dry methanol was cooled to 0°C and was treated with 0.105 g
(1.9 mmol) of sodium methoxide. The ice-bath was removed and the
25 mixture was allowed to stir at room temperature for 30 minutes. To the
reaction was then added 0.110 mL (1.9 mmol) of acetic acid and then
5.8 g (64.9 mmol) of methyl hydrazinecarboxylate. After stirring 30
minutes at room temperature, the suspension was concentrated in vacuo,
and placed on the high-vac line overnight, to give 10.5 g (98%) of a
30 yellow powder, a portion of which was employed in Step C below.

Step B: 4-(2-(N-Methylcarboxy-acetamidrazono)-2-(S)-(3,5-
dichlorobenzyloxy)-3-(S)-phenylmorpholine

A solution of 0.050 g (0.15 mmol) of 2-(S)-

- 168 -

(3,5-dichlorobenzyloxy)-3-(S)-phenylmorpholine (from Example 44, Step C), 0.034 g (0.21 mmol) of N-methyl-carboxy-2-chloro-acteamidrazone (from Step A), and 0.044 mL (0.25 mmol) N,N-diisopropylethylamine in 1 mL of acetonitrile was stirred at room temperature for 3 hours. The mixture was partitioned between 20 mL of methylene chloride and 10 mL of water. The layers were separated, the organic layer was dried over sodium sulfate and was then concentrated in vacuo. The residue was purified by flash chromatography on 35 g of silica eluting with 1L of 50:1: methylene chloride/methanol then 500 mL of 25:1:0.05 methylene chloride:methanol:aqueous ammonia to give 70 mg (~100%) of the product as a white solid.

Mass Spectrum (FAB): m/Z 469 (M+H, 60%), 467 (M+H, 100%), 291 (40%), 160 (20%), 158 (25%).

¹H NMR (CDCl₃, 400 MHz, ppm): δ 2.48 (td, J= 3.5, 12.2, 1H), 2.53 (d, J= 14.6, 1H), 2.90 (d, J= 11.8, 1H), 3.37 (d, J= 14.6, 1H), 3.52 (d, J= 2.8), 1H), 3.62 (dm, J= 11.4, 1H), 3.75 (s, 3H), 4.14 (td, J= 2.2, 11.8, 1H), 4.28 (d, J= 13.5, 1H), 4.58 (d, J= 13.6), 4.60 (d, J= 2.8, 1H), 5.45 (br s, 2H), 6.74 (d, J= 1.9, 2H), 7.15 (t, J= 1.9, 1H), 7.30-7.46 (m, 6H).

Step C: 2-(S)-(3,5-Dichlorobenzyloxy)-4-(3-(5-oxo-1,2,4-triazolo)-methyl)-3-(S)-phenylmorpholine

A solution of 0.069 g (0.15 mmol) of 4-(2-(N-methyl-carboxyacetamidrazono)-2-(S)-(3,5-dichlorobenzyloxy)-3-(S)-phenylmorpholine (from Step B) in 6 mL of xylenes was heated at reflux for 2 hours. The solution was cooled and concentrated in vacuo. The residue was purified by flash chromatography on 35 g of silica gel eluting with 500 mL of 50:1:0.1 methylene chloride/methanol/aqueous ammonia then 500 mL of 20:1:0.1 methylene chloride/methanol/aqueous ammonia to give 56 mg (88%) of the product as a white powder.

- 169 -

Mass Spectrum (FAB): m/z 437 (M+H, 65%), 435 (M+H, 100%), 259 (85%), 161 (55%).

¹H NMR (CDCl₃, 400 MHz, ppm): δ 2.53 (t, J= 11.7, 3.6, 1H), 2.88 (d, J= 11.6, 1H), 2.96 (d, J= 14.3, 1H), 3.54 (d, J= 2.6, 1H), 3.63 (dd, J= 11.6, 1.9, 1H), 3.68 (d, J= 14.6, 1H), 4.16 (t, J= 11.7, 2.2, 1H), 4.30 (d, J= 13.6), 4.58 (d, J= 2.7, 1H), 4.67 (d, J= 13.6, 1H), 6.65 (d, J= 1.8, 2H), 7.07 (t, J= 1.9, 1H), 7.29-7.44 (m, 5H), 10.25 (br s, 1H), 10.75 (br s, 1H).

EXAMPLE 46

2-(S)-(3,5-Bis(trifluoromethyl)benzyloxy)-4-(methoxy-carbonyl-methyl)-3-(S)-phenylmorpholine

A solution of 300 mg (0.74 mmole) of 2-(S)-(3,5-bis(trifluoromethyl)benzyloxy)-3-(S)-phenyl-morpholine (from Example 15, Step C) and 0.35 mL (2.0 mmole) of DIEA in 5 mL of acetonitrile was treated with 0.19 mL (2.0 mmole) of methyl bromoacetate and the mixture was stirred for 16 hr at room temperature. The solution was then concentrated *in vacuo* and the residue partitioned between 30 mL of ether and 15 mL of 0.5 N aqueous KHSO₄. The layers were separated and the organic phase was washed with 10 mL of brine and dried over magnesium sulfate. Following filtration, the organic phase was concentrated *in vacuo* and the residue purified by flash chromatography on 20 g of silica eluting with 80:20 hexanes:ether to give 351 mg (99%) of the product. $[\alpha]_D^{25} = +147.3^\circ$ (c= 1.6, CHCl₃).

Mass Spectrum (FAB): m/z 478 (M+H, 40%), 477 (65%), 418 (50%), 250 (95%), 234 (90%), 227 (100%).

¹H NMR (CDCl₃, 400 MHz, ppm): δ 3.02 (br d, 2H), 3.13 (d, J= 16.9, 1H), 3.36 (d, J= 16.8), 3.62 (s, 3H), 3.69 (dt, J= 11.7, 2.2, 1H), 4.03 (br

- 170 -

s, 1H), 4.23-4.32 (m, 1H), 4.44 (d, J= 13.3, 1H), 4.68, (d, J= 2.6, 1H),
4.81 (d, J= 13.5, 1H), 7.30-7.38 (m, 3H), 7.4-7.5 (m, 3H), 7.70 (s, 1H).

Analysis Calcd for C₂₂H₂₁F₆NO₄:

C, 55.35; H, 4.43; N, 2.93; F, 23.88

5 Found: C, 55.09; H, 4.43; N, 2.83; F, 24.05

EXAMPLE 47

10 2-(S)-(3,5-Bis(trifluoromethyl)benzyloxy)-4-(carboxymethyl)-3-(S)-phenylmorpholine

A solution of 0.016 g (0.034 mmole) of 2-(S)-(3,5-Bis(trifluoromethyl)benzyloxy)-4-(methoxy-carbonylmethyl)-3-(S)-phenylmorpholine (from Example 46) in 2 mL of THF and 0.5 mL of water was treated with 0.027 mL (0.067 mmole) of 2.5 N aqueous sodium hydroxide and the mixture was stirred at room temperature for 15 5 hr. The mixture was treated with 2 drops of 2N aqueous HCl and 3 mL of water and the solution was extracted with 15 mL of 1:1 hexanes:ethyl acetate. The organic phase was dried over magnesium sulfate, filtered and concentrated in vacuo. The residue was purified by 20 flash chromatography on 13 g of silica eluting with 250 mL of 100:3:0.1 methylene chloride:methanol:acetic acid then 100 mL of 50:2:0.1 methylene chloride:methanol:acetic acid to give 0.014 g (90%) of an oil.

25 Mass Spectrum (FAB): m/Z 464 (M+H, 90%), 420 (M-CO₂, 10%), 227 (ArCH₂, 35%), 220 (M-OCH₂Ar, 100%), 161 (20%).

¹H NMR (CDCl₃, 400 MHz, ppm): δ 2.9 (app d, 2H), 3.03 (d, 1H), 3.33 (d, 1H), 3.72 (d, 1H), 3.90 (d, 1H), 4.25 (t, 1H), 4.44 (d, 1H), 4.71 (d, 1H), 4.79 (d, 1H), 7.3-7.4 (m, 5H), 7.44 (s, 2H), 7.71 (s, 1H). 30

EXAMPLE 48

- 171 -

2-(S)-(3,5-Bis(trifluoromethyl)benzyloxy)-4-((2-aminoethyl)amino-carbonylmethyl)-3-(S)-phenylmorpholine hydrochloride

A solution of 54 mg (0.11 mmole) of 2-(S)-(3,5-bis(trifluoromethyl)benzyloxy)-4-(carboxymethyl)-3-(S)-phenylmorpholine (from Example 46) and 0.15 mL of ethylenediamine (2.3 mmole) in 1 mL of methanol was stirred at 55°C for 48 hr. The mixture was concentrated and the residue purified by flash chromatography on 16 g of silica eluting with 500 mL of 50:4:0.1 methylene chloride:methanol:aqueous ammonia to provide 57 mg(100%) of an oil. The oil was dissolved in ether and was treated with ether saturated with gaseous HCl. After concentration in vacuo, 58 mg (95%) of a rigid oil was obtained.

Mass Spectrum (FAB; free base): m/Z 506 (M+H, 100%), 418 (15%), 262(35%), 227 (30%), 173 (40%)
1H NMR (CDCl₃, 400 MHz, ppm): δ 2.56 (d, J= 15.5, 1H), 2.59 (td, J= 12.0, 3.6, 1H), 2.82 (t, J= 6.5, 2H), 2.96 (d, J= 11.8, 1H), 3.21 (d, J= 15.8, 1H), 3.25-3.40 (m, 2H), 3.65 (d, J= 2.6, 1H), 3.67 (app dt, J= 11.4, ~2, 1H), 4.18 (td, J= 11.8, 2.6, 1H), 4.33 (d, J= 13.5, 1H), 4.69 (d, J= 2.7, 1H), 4.79 (d, J= 13.5, 1H), 7.25-7.40 (m, 5H), 7.46 (s, 2H), 7.59 (br t, 1H), 7.71 (s, 1H).

EXAMPLE 49

2-(S)-(3,5-Bis(trifluoromethyl)benzyloxy)-4-((3-amino-propyl)amino carbonylmethyl)-3-(S)-phenylmorpholine hydrochloride

A solution of 59 mg (0.12 mmole) of 2-(S)-(3,5-bis(trifluoromethyl)benzyloxy)-4-(carboxymethyl)-3-(S)-phenylmorpholine (from Example 46) and 0.21 mL of 1,3-propylenediamine (2.5 mmole) in 1 mL of methanol was stirred at 55°C for 72 hr. The mixture was concentrated and the residue purified by flash chromatography on 16 g of silica eluting with 500 mL of 10:1:0.05 methylene chloride:methanol:aqueous ammonia to provide 56 mg (88%) of an oil. The oil was dissolved in methylene chloride and was treated with methylene chloride

- 172 -

saturated with gaseous HCl. After concentration in vacuo, a white paste was obtained.

Mass Spectrum (FAB; free base): m/Z 520 (M+H, 100%), 418 (10%), 276(30%), 227 (20%), 174 (30%)

5 ¹H NMR (CDCl₃, 400 MHz, ppm): δ 1.64 (pentet, J= 6.6, 2H), 2.53 (d, J= 15.5, 1H), 2.58 (td, J= 12.0, 3.6, 1H), 2.73 (t, J= 6.5, 2H), 2.92 (d, J= 11.8, 1H), 3.19 (d, J= 15.8, 1H), 3.25-3.40 (m, 2H), 3.62 (d, J= 2.6, 1H), 3.65 (app dt, J= 11.4, ~2, 1H), 4.16 (td, J= 11.8, 2.6, 1H), 4.41 (d, J= 13.5, 1H), 4.68 (d, J= 2.7, 1H), 4.79 (d, J= 13.5, 1H), 7.25-7.40 (m, 10 5H), 7.45 (s, 2H), 7.57 (br t, 1H), 7.70 (s, 1H).

EXAMPLE 50

15 4-benzyl-5-(S),6-(R)-dimethyl-3-(S)-phenylmorpholinone and 4-benzyl-5-(R),6-(S)-dimethyl-3-(S)-phenylmorpholinone

To a suspension of 1.7 g (7.0 mmole) of N-benzyl-(S)-phenylglycine (Example 13) in 15 ml of methylene chloride at 0°C was added 6.9 ml (13.9 mmole) of trimethylaluminum (2.0 M in toluene). After one hour at 0°C, 0.625 ml (7.0 mmole) of (+/-)-trans-2,3-epoxy butane (dissolved in 2.0 ml of methylene chloride) was added dropwise and then allowed to stir at 22°C for 16 hours. The reaction was then transferred to another flask containing 30 ml of 1:1 hexane:methylene chloride and 30 ml of 1M potassium sodium tartrate and stirred at 22°C for 2 hours. The layers were separated, and the aqueous layer was 25 extracted with methylene chloride (3 x 100 ml). The combined organic layers were washed with 25 ml of a saturated sodium chloride solution, dried over anhydrous sodium sulfate, filtered, and concentrated in vacuo.

30 The crude alcohol was dissolved in 25 ml of toluene, treated with 93 mg (0.49 mmole) of p-toluenesulfonic acid and heated at 50°C for 20 hours. The reaction was then cooled and concentrated in vacuo. The residue was partitioned between 15 ml of diethyl ether and 10 ml of saturated sodium bicarbonate. The layers were separated, and the organic layer was washed with water (3 x 10 ml). The combined

- 173 -

organic layers were washed with 25 ml of a saturated sodium chloride solution, dried over anhydrous magnesium sulfate, filtered, and concentrated in vacuo. Flash chromatography on 145 g of silica gel using 1:4 v/v ethyl acetate/ hexane as the eluant afforded 567 mg of the high R_f lactone (Isomer A) and 388 mg of the low R_f lactone (Isomer B).

¹H-NMR (400 MHz, CDCl₃) δ Isomer A: 1.04 (d, 3H, J = 8.0 Hz), 1.24 (d, 3H, J = 8.0 Hz), 2.92 (br qd, 1H), 3.41 (d, 1H, J = 16.0 Hz), 3.62 (d, 1H, J = 16.0 Hz), 4.38 (s, 1H), 4.96 (br qd, 1H), 7.20-7.42 (m, 8H), 7.58-7.64 (m, 2H); Isomer B: 1.04 (d, 3H, J = 10.0 Hz), 1.39 (d, 3H, J = 10.0 Hz), 3.06 (br qd, 1H), 3.53 (d, 1H, J = 16.0 Hz), 3.81 (d, 1H, J = 16.0 Hz), 4.33 (s, 1H), 4.67 (br qd, 1H), 7.18-7.50 (m, 10H). Mass Spectrum (FAB): m/z Isomer A: 296 (M+H, 100%), 294 (50%); Isomer B: 296 (M+H, 100%), 294 (50%).

EXAMPLE 51

2-(R)-(3,5-Bis(trifluoromethyl)benzyloxy)-[5-(S),6-(R) or 5-(R),6-(S)-dimethyl]-3-(S)-phenylmorpholinone

Step A: 4-Benzyl-2-(R)-(3,5-bis(trifluoromethyl)-benzyloxy)-[5-(S),6-(R) or 5-(R),6-(S)-dimethyl]-3-(S)-phenylmorpholinone

According to the procedure in Example 15, Step B, 251 mg (0.85 mmole) of Isomer A from Example 50 (4-benzyl-[5-(S),6-(R) or 5-(R)-6-(S)-dimethyl]-3-(S)-phenylmorpholinone) provided 238 mg (53%) of the product as an oil.

¹H-NMR (400 MHz, CDCl₃) δ 1.03 (d, 3H, J = 6.7 Hz), 1.13 (d, 3H, J = 6.6 Hz), 2.61 (qd, 1H, J = 2.2 & 6.6 Hz), 3.26 (d, 1H, J = 13.9 Hz), 3.55 (d, 1H, J = 13.9 Hz), 3.63 (d, 1H, J = 7.6 Hz), 4.01 (qd, 1H, J = 2.3 & 6.6 Hz), 4.44 (d, 1H, J = 13.1 Hz), 4.53 (d, 1H, J = 7.7 Hz), 4.71 (s, 1H), 4.85 (d, 1H, J = 13.2 Hz), 7.20-7.35 (m, 9H), 7.46-7.48 (m, 2H), 7.67 (s, 1H), 7.81 (s, 1H).

- 174 -

Mass Spectrum (FAB): m/z 523 (M+H, 100%), 296 (95%), 280 (40%), 227 (50%).

5 Step B: 2-(R)-(3,5-Bis(trifluoromethyl)benzyloxy)-[5-(S),6-(R) or 5-(R),6-(S)-dimethyl]-3-(S)-phenylmorpholinone

According to the procedure in Example 15, Step C, 260 mg of starting material from Step A [derived from Isomer A in Example 50 (4-Benzyl-2-(R)-(3,5-bis(trifluoromethyl)benzyloxy)-[5-(S),6-(R) or 5-(R),6-(S)-dimethyl]-3-(S)-phenylmorpholinone)] provided 122 mg
10 (57%) of the product as an oil.

$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 1.19 (d, 3H, $J = 6.5$ Hz), 1.27 (d, 3H, $J = 6.7$ Hz), 2.97 (qd, 1H, $J = 2.9$ & 6.9 Hz), 3.96 (d, 1H, $J = 7.7$ Hz), 4.08-4.11 (m, 2H), 4.39 (d, 1H, $J = 7.7$ Hz), 4.50 (d, 1H, $J = 13.3$ Hz), 4.88 (d, 1H, $J = 13.2$ Hz), 7.27-7.33 (m, 3H), 7.40-7.42 (m, 4H), 7.67 (s, 1H).
15

Mass Spectrum (FAB): m/z 434 (M+H, 45%), 227 (35%), 206 (40%), 190 (100%).

20 EXAMPLE 52

2-(S)-(3,5-Bis(trifluoromethyl)benzyloxy)-[5-(R),6-(S) or 5-(S),6-(R)-dimethyl]-3-(S)-phenylmorpholinone

25 Step A: 4-Benzyl-2-(S)-(3,5-bis(trifluoromethyl)-benzyloxy)-[5-(R),6-(S) or 5-(S),6-(R)-dimethyl]-3-(S)-phenylmorpholinone

According to the procedure in Example 15, Step B, 449 mg (1.52 mmole) of Isomer B from Example 50 (4-benzyl-[5-(R),6-(S) or 5-(S)-6-(R)-dimethyl]-3-(S)-phenylmorpholinone) provided 400 mg
30 (51%) of the product as an oil.

$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 0.90 (d, 3H, $J = 6.8$ Hz), 1.37 (d, 3H, $J = 6.6$ Hz), 2.86-2.89 (br qd, 1H), 3.47 (d, 1H, $J = 15.0$ Hz), 3.82-3.85 (m, 2H), 3.99-4.02 (br qd, 1H), 4.45 (d, 1H, $J = 13.6$ Hz), 4.81 (d, 1H, $J = 2.0$ Hz), 4.87 (d, 1H, $J = 13.5$ Hz), 7.17-7.83 (m, 13H).

- 175 -

Step B: 2-(S)-(3,5-Bis(trifluoromethyl)benzyloxy)-[5-(S),6-(R) or 5-(R),6-(S)-dimethyl]-3-(S)-phenylmorpholinone

According to the procedure in Example 15, Step C, 400 mg of starting material from Step A [derived from Isomer B in Example 50 (4-Benzyl-2-(S)-(3,5-bis(trifluoromethyl)benzyloxy)-[5-(R),6-(S) or 5-(S),6-(R)-dimethyl]-3-(S)-phenylmorpholinone)] provided 230 mg (69%) of the product as an oil.

¹H-NMR (400 MHz, CDCl₃) δ 1.08 (d, 3H, J = 6.7 Hz), 1.38 (d, 3H, J = 7.0 Hz), 3.41-3.45 (br qd, 1H), 3.85-3.89 (br qd, 1H), 4.16 (d, 1H, J = 2.9 Hz), 4.49 (d, 1H, J = 13.6 Hz), 4.71 (d, 1H, J = 2.9 Hz), 4.82 (d, 1H, J = 13.6 Hz), 7.25-7.36 (m, 7H), 7.66 (s, 1H).

Mass Spectrum (FAB): m/z 434 (M+H, 35%), 227 (40%), 206 (40%), 190 (100%).

EXAMPLE 53

2-(R)-(3,5-Bis(trifluoromethyl)benzyloxy)-4-(3-(1,2,4-triazolo)-methyl)-[5-(S),6-(R) or 5-(R),6-(S)-dimethyl]-3-(S)-phenylmorpholinone

A mixture of 62 mg (0.14 mmole) of 2-(R)-(3,5-Bis(trifluoromethyl)benzyloxy)-[5-(S),6-(R) or 5-(R),6-(S)-dimethyl]-3-(S)-phenylmorpholinone (from Example 51, Step B), 62 mg (0.45 mmole) of anhydrous potassium carbonate and 26 mg (0.19 mmole) of N-formyl-2-chloroacetamidrazone (from Example 17, Step A) in 2.0 ml of N,N-dimethylformamide was heated to 60°C for 2 hours and then 118°C for 1.5 hours. The mixture was then allowed to cool to room temperature and then quenched with 5 mls of water and diluted with 15 mls of ethyl acetate. The layers were separated and the organic layer was washed with ethyl acetate (2 x 10 mls). The combined organic layers were washed with 10 mls of brine, dried over anhydrous magnesium sulfate, filtered, and concentrated in vacuo. Flash chromatography on 42 g of silica gel using 95:5 v/v methylene chloride/methanol as the eluant afforded 42 mg (57%) of a clear oil.

- 176 -

¹H-NMR (400 MHz, CDCl₃) δ 1.13 (d, 3H, J = 6.5 Hz), 1.19 (d, 3H, J = 6.5 Hz), 2.65 (qd, 1H, J = 1.9 & 6.5 Hz), 3.58 (d, 1H, J = 15.5 Hz), 3.65 (d, 1H, J = 7.7 Hz), 3.75 (d, 1H, J = 15.4 Hz), 4.06 (qd, 1H, J = 2.2 & 6.6 Hz), 4.45 (d, 1H, J = 13.2 Hz), 4.54 (d, 1H, J = 7.7 Hz), 4.84 (d, 1H, J = 13.2 Hz), 7.28-7.37 (m, 7H), 7.67 (s, 1H), 7.89 (s, 1H).

Mass Spectrum (FAB): m/z 516 (M+H, 52%), 287 (28%), 271 (100%), 227 (40%), 202 (38%).

EXAMPLE 54

2-(R)-(3,5-Bis(trifluoromethyl)benzyloxy)-4-(3-(5-oxo-1,2,4-triazolo methyl)-[5-(S),6-(R) or 5-(R),6-(S)-dimethyl]-3-(S)-phenyl-morpholinone

A solution of 96 mg (0.22 mmole) of 2-(R)-(3,5-Bis(trifluoromethyl)benzyloxy)-[5-(S),6-(R) or 5-(R),6-(S)-dimethyl]-3-(S)-phenylmorpholinone (from Example 51, Step B), 92 mg (0.66 mmole) of potassium carbonate and 48 mg (0.29 mmole) of N-methylcarboxy-2-chloroacetamidrazone (from Example 18, Step A) in 4 mL of DMF was heated at 60°C for 1.5 hr and at 120°C for 3.5 hr. The mixture was cooled to room temperature and was partitioned between 15 mL of water and 25 mL of ethyl acetate. The aqueous layer was extracted with 3x10 mL of ethyl acetate, the combined organic layers were washed with 10 mL of brine, dried over sodium sulfate, filtered and concentrated in vacuo. The residue was partly purified by flash chromatography on 42 g of silica gel using 2L of 98:2 v/v methylene chloride/ methanol as the eluant and the rich cuts were purified under the same conditions to give 38 mg (33%) of a clear oil.

¹H-NMR (400 MHz, CDCl₃) δ 1.09 (d, 3H, J = 6.5 Hz), 1.20 (d, 3H, J = 6.6 Hz), 2.64 (qd, 1H, J = 2.4 & 6.6 Hz), 3.33 (s, 1H), 3.56 (d, 1H, J = 7.6 Hz), 4.11 (qd, 1H, J = 2.4 & 6.6 Hz), 4.41 (d, 1H, J = 13.2 Hz), 4.57 (d, 1H, J = 7.7 Hz), 4.82 (d, 1H, J = 13.2 Hz), 7.25-7.30 (m, 5H), 7.40 (d, 2H, J = 5.7 Hz), 7.65 (s, 1H), 9.46 (s, 1H), 10.51 (s, 1H).

Mass Spectrum (FAB): m/z 531 (M+H, 98%), 287 (100%), 227 (80%), 189 (65%).

- 177 -

EXAMPLE 55

5 2-(S)-(3,5-Bis(trifluoromethyl)benzyloxy)-4-(3-(1,2,4-triazolo)methyl)-
 [5-(R),6-(S) or 5-(S),6-(R)-dimethyl]-3-(S)-phenylmorpholinone

 According to the procedure in Example 53, 75 mg (0.17 mmole) of 2-(S)-(3,5-Bis(trifluoromethyl)-benzyloxy)-[5-(R),6-(S) or 5-(S),6-(R)-dimethyl]-3-(S)-phenylmorpholinone (from Example 52, Step B) provided, after flash chromatography on 73 g of silica gel using
10 98:2 v/v methylene chloride/ methanol as the eluant, 46 mg (52%) of a yellow oil.

¹H-NMR (400 MHz, CDCl₃) δ 1.04 (d, 3H, J = 6.6 Hz), 1.46 (d, 3H, J = 6.7 Hz), 3.05-3.08 (m, 1H), 3.74-3.81 (m, 2H), 3.91-3.95 (m, 2H), 4.41 (d, 1H, J = 13.2 Hz), 4.69 (d, 1H, J = 3.2 Hz), 4.82 (d, 1H, J = 13.5 Hz),
15 7.31-7.35 (m, 5H), 7.43-7.45 (m, 2H), 7.68 (s, 1H), 7.91 (s, 1H).

 Mass Spectrum (EI): m/z 432 (36%), 287 (60%), 270 (65%), 227 (30%), 187 (48%), 83 (100%).

EXAMPLE 56

20

2-(S)-(3,5-Bis(trifluoromethyl)benzyloxy)-4-(3-(5-oxo-1,2,4-triazolo)methyl)-[5-(R),6-(S) or 5-(S),6-(R)-dimethyl]-3-(S)-phenyl-
 morpholinone

 According to the procedure in Example 54, 86 mg (0.2 mmole) of 2-(S)-(3,5-Bis(trifluoromethyl)-benzyloxy)-[5-(R),6-(S) or 5-(S),6-(R)-dimethyl]-3-(S)-phenylmorpholinone (from Example 47, Step B) provided, after flash chromatography on 73 g of silica gel using
25 95:5 v/v methylene chloride/ methanol as the eluant, 32 mg (30%) of a yellow oil.

30 ¹H-NMR (400 MHz, CDCl₃) δ 1.03 (d, 3H, J = 6.7 Hz), 1.40 (d, 3H, J = 6.8 Hz), 3.00 (qd, 1H, J = 3.8 & 6.8 Hz), 3.44 (d, 1H, J = 16.1 Hz), 3.63 (d, 1H, J = 16.0 Hz), 3.82 (d, 1H, J = 3.3 Hz), 3.95 (qd, 1H, J = 3.7 & 6.7 Hz), 4.43 (d, 1H, J = 13.5 Hz), 4.73 (d, 1H, J = 3.3 Hz), 4.84 (d, 1H, J = 13.6 Hz), 7.28-7.47 (m, 7H), 7.68 (s, 1H), 9.52 (d, 2H).

- 178 -

Mass Spectrum (FAB): m/z 531 (M+H, 100%), 287 (55%), 227 (25%), 147 (50%).

EXAMPLE 57

5

2-(S)-(3,5-Bis(trifluoromethyl)benzyloxy)-4-(2-(1-(4-benzyl)-piperidino)ethyl)-3-(S)-phenylmorpholine

To a solution of 2-(S)-(3,5-bis(trifluoro-methyl)-benzyloxy)-3-(S)-phenylmorpholine (50 mg, 0.12 mmol) and 4-benzyl-10 1-(2-chloroethyl)piperidine hydrochloride (50 mg, 0.18 mmol) in acetonitrile (0.5 mL) was added diisopropylethylamine (0.065 mL, 0.36 mmol) at room temperature. After 60 hours, TLC (5% MeOH/2% Et₃N/93% EtOAc) indicated that the reaction was only partially complete. The reaction was diluted with methylene chloride and washed 15 with water, then brine, dried over sodium sulfate and evaporated. Prep TLC (5% MeOH/2% Et₃N/93% EtOAc) afforded 36 mg (50%) of the title compound as an oil.

¹H-NMR (400 MHz, CDCl₃) δ 1.1-1.4 (m, 2 H), 1.4-1.65 (2 m, 4 H), 1.65-2.05 (m, 3 H), 2.05-2.3 (m, 1H), 2.35-2.5 (m and d, J = 7 Hz, 3 20 H), 2.55 (br t, J = 11 Hz, 1 H), 2.65-2.8 (m, 2 H), 3.09 (d, J = 11 Hz, 1 H), 3.50 (d, J = 2.5 Hz, 1 H), 3.66 (dd, J = 2 and 11 Hz, 1 H), 4.15 (dt, J = 2 and 12 Hz, 1 H), 4.38 and 4.75 (AB q, J = 13 Hz, 2 H), 4.61 (d, J = 2.5 Hz, 1 H), 7.06 (d, J = 7 Hz, 2 H), 7.15 (t, J = 7 Hz, 1 H), 7.2-7.35 (m, 5 H), 7.36 (m, 4 H), 7.75 (s, 1H).

25

EXAMPLE 58

(S)-(4-Fluorophenyl)glycine

Via Chiral Synthesis:

30

Step A: 3-(4-Fluorophenyl)acetyl-4-(S)-benzyl-2-oxazolidinone

An oven-dried, 1 L 3-necked flask, equipped with a septum, nitrogen inlet, thermometer, and a magnetic stirring bar, was flushed with nitrogen and charged with a solution of 5.09 g (33.0 mmol) of 4-fluorophenylacetic acid in 100 mL of anhydrous ether. The

- 179 -

solution was cooled to -10°C and treated with 5.60 mL (40.0 mmol) of triethylamine followed by 4.30 mL (35.0 mmol) of trimethylacetyl chloride. A white precipitate formed immediately. The resulting mixture was stirred at -10°C for 40 minutes, then cooled to -78°C.

5 An oven-dried, 250 mL round bottom flask, equipped with a septum and a magnetic stirring bar, was flushed with nitrogen and charged with a solution of 5.31 g (30.0 mmol) of 4-(S)-benzyl-2-oxazolidinone in 40 mL of dry THF. The solution was stirred in a dry ice/acetone bath for 10 minutes, then 18.8 mL of 1.6 M n-butyllithium
10 solution in hexanes was slowly added. After 10 minutes, the lithiated oxazolidinone solution was added, via cannula, to the mixture in the 3-necked flask. The cooling bath was removed from the resulting mixture and the temperature was allowed to rise to 0°C. The reaction was quenched with 100 mL of saturated aqueous ammonium chloride
15 solution, transferred to a 1 L flask, and the ether and THF were removed in vacuo. The concentrated mixture was partitioned between 300 mL of methylene chloride and 50 mL of water and the layers were separated. The organic layer was washed with 200 mL of 2 N aqueous hydrochloric acid solution, 300 mL of saturated aqueous sodium
20 bicarbonate solution, dried over magnesium sulfate and concentrated in vacuo. Flash chromatography on 400 g of silica gel using 3:2 v/v hexanes/ether as the eluant afforded 8.95 g of an oil that slowly solidified on-standing. Recrystallization from 10:1 hexanes/ether afforded 7.89 g (83%) of the title compound as a white solid, mp 64-
25 66°C.

Mass Spectrum (FAB): m/Z 314 (M+H, 100%), 177 (M-ArCH₂CO+H, 85%).

¹H-NMR (400 MHz, CDCl₃): δ 2.76 (dd, 1 H, J = 13.2, 9.2), 3.26 (dd, J = 13.2, 3.2), 4.16-4.34 (m, 4 H), 4.65-4.70 (m, 1 H), 7.02-7.33 (m, 9
30 H).

Analysis: Calcd for C₁₈H₁₆FNO₃:

C, 69.00; H, 5.15; N, 4.47; F, 6.06

Found: C, 68.86; H, 5.14; N, 4.48; F, 6.08

- 180 -

Step B: 3-((S)-Azido-(4-fluorophenyl))acetyl-4-(S)-benzyl-2-oxazolidinone

An oven-dried, 1 L 3-necked flask, equipped with a septum, nitrogen inlet, thermometer, and a magnetic stirring bar, was
5 flushed with nitrogen and charged with a solution of 58.0 mL of 1 M potassium bis(trimethylsilyl)amide solution in toluene and 85 mL of THF and was cooled to -78°C. An oven-dried, 250 mL round-bottomed flask, equipped with a septum and a magnetic stirring bar, was flushed
10 with nitrogen and charged with a solution of 7.20 g (23.0 mmol) of 3-(4-fluorophenyl)acetyl-4-(S)-benzyl-2-oxazolidinone (from Example 58, Step A) in 40 mL of THF. The acyl oxazolidinone solution was stirred in a dry ice/acetone bath for 10 minutes, then transferred, via cannula, to the potassium bis(trimethylsilyl)amide solution at such a rate
15 that the internal temperature of the mixture was maintained below -70°C. The acyl oxazolidinone flask was rinsed with 15 mL of THF and the rinse was added, via cannula, to the reaction mixture and the resulting mixture was stirred at -78°C for 30 minutes. An oven-dried, 250 mL round-bottomed flask, equipped with a septum and a
20 magnetic stirring bar, was flushed with nitrogen and charged with a solution of 10.89 g (35.0 mmol) of 2,4,6-triisopropylphenylsulfonyl azide in 40 mL of THF. The azide solution was stirred in a dry ice/acetone bath for 10 minutes, then transferred, via cannula, to the reaction mixture at such a rate that the internal temperature of the
25 mixture was maintained below -70°C. After 2 minutes, the reaction was quenched with 6.0 mL of glacial acetic acid, the cooling bath was removed and the mixture was stirred at room temperature for 18 hours. The quenched reaction mixture was partitioned between 300 mL of ethyl acetate and 300 mL of 50% saturated aqueous sodium bicarbonate solution. The organic layer was separated, dried over magnesium
30 sulfate, and concentrated in vacuo. Flash chromatography on 500 g of silica gel using 2:1 v/v, then 1:1 v/v hexanes/methylene chloride as the eluant afforded 5.45 g (67%) of the title compound as an oil.
IR Spectrum (neat, cm⁻¹): 2104, 1781, 1702.

- 181 -

¹H-NMR (400 MHz, CDCl₃): δ 2.86 (dd, 1 H, J = 13.2, 9.6), 3.40 (dd, 1 H, J = 13.2, 3.2), 4.09-4.19 (m, 2 H), 4.62-4.68 (m, 1 H), 6.14 (s, 1 H), 7.07-7.47 (m, 9 H).

Analysis: Calcd for C₁₈H₁₅FN₄O₃:

C, 61.01; H, 4.27; N, 15.81; F, 5.36

Found: C, 60.99; H, 4.19; N, 15.80; F, 5.34

Step C: (S)-Azido-(4-fluorophenyl)acetic acid

A solution of 5.40 g (15.2 mmol) of 3-((S)-azido-(4-fluorophenyl))acetyl-4-(S)-benzyl-2-oxazolidinone (from Example 58, Step B) in 200 mL of 3:1 v/v THF/water was stirred in an ice bath for 10 minutes. 1.28 g (30.4 mmol) of lithium hydroxide monohydrate was added in one portion and the resulting mixture was stirred cold for 30 minutes. The reaction mixture was partitioned between 100 mL of methylene chloride and 100 mL of 25% saturated aqueous sodium bicarbonate solution and the layers were separated. The aqueous layer was washed with 2 x 100 mL of methylene chloride and acidified to pH 2 with 2 N aqueous hydrochloric acid solution. The resulting mixture was extracted with 2 x 100 mL of ethyl acetate; the extracts were combined, washed with 50 mL of saturated aqueous sodium chloride solution, dried over magnesium sulfate, and concentrated in vacuo to afford 2.30 g (77%) of the title compound as an oil that was used in the following step without further purification.

IR Spectrum (neat, cm⁻¹): 2111, 1724.

¹H-NMR (400 MHz, CDCl₃): δ 5.06 (s, 1 H), 7.08-7.45 (m, 4 H), 8.75 (br s, 1 H).

Step D: (S)-(4-Fluorophenyl)glycine

A mixture of 2.30 g (11.8 mmol) of (S)-azido-(4-fluorophenyl)acetic acid (from Example 58, Step C), 250 mg 10% palladium on carbon catalyst and 160 mL 3:1 v/v water/acetic acid was stirred under an atmosphere of hydrogen for 18 hours. The reaction mixture was filtered through Celite and the flask and filter cake were rinsed well with ~1L of 3:1 v/v water/acetic acid. The filtrate was

- 182 -

concentrated in vacuo to about 50 mL of volume. 300 mL of toluene was added and the mixture concentrated to afford a solid. The solid was suspended in 1:1 v/v methanol/ether, filtered and dried to afford 1.99 g (100%) of the title compound.

5 $^1\text{H-NMR}$ (400 MHz, D_2O + NaOD): δ 3.97 (s, 1 H), 6.77 (app t, 2 H, J = 8.8), 7.01 (app t, 2 H, J = 5.6).

Via Resolution:

Step A': 4-Fluorophenylacetyl chloride

10 A solution of 150 g (0.974 mol) of 4-fluorophenylacetic acid and 1 mL of N,N -dimethylformamide in 500 mL of toluene at 40 °C was treated with 20 mL of thionyl chloride and heated to 40°C. An additional 61.2 mL of thionyl chloride was added dropwise over 1.5 hours. After the addition, the solution was heated at 50 °C for 1 hour,
15 the solvent was removed in vacuo and the residual oil was distilled at reduced pressure (1.5 mmHg) to afford 150.4 g (89.5%) of the title compound, bp = 68-70 °C.

Step B': Methyl 2-bromo-2-(4-fluoro)phenylacetate

20 A mixture of 150.4 g (0.872 mol) of 4-fluorophenylacetyl chloride (from Example 58, Step A') and 174.5 g (1.09 mol) of bromine was irradiated at 40-50 °C with a quartz lamp for 5 hours. The reaction mixture was added dropwise to 400 mL of methanol and the solution was stirred for 16 hours. The solvent was removed in vacuo
25 and the residual oil was distilled at reduced pressure (1.5 mmHg) to afford 198.5 g (92%) of the title compound, bp = 106-110 °C.

Step C': Methyl (\pm)-(4-fluorophenyl)glycine

30 A solution of 24.7 g (0.1 mol) of methyl 2-bromo-2-(4-fluoro) phenylacetate (from Example 58, Step B') and 2.28 g (0.01 mol) of benzyl triethylammonium chloride in 25 mL of methanol was treated with 6.8 g (0.105 mol) of sodium azide and the resulting mixture was stirred 20 hours at room temperature. The reaction mixture was filtered; the filtrate was diluted with 50 mL of methanol and

- 183 -

hydrogenated in the presence of 0.5 g of 10% Pd/C at 50psi for 1 hour. The solution was filtered and the solvent removed in vacuo. The residue was partitioned between 10% aqueous sodium carbonate solution and ethyl acetate. The organic phase was washed with water, saturated
5 aqueous sodium chloride solution, dried over magnesium sulfate and concentrated in vacuo to afford 9.8 g of the title compound as an oil.

Step D': Methyl (S)-(4-fluorophenyl)glycinate

A solution of 58.4 g of methyl (\pm)-4-fluorophenylglycinate
10 (from Example 58, Step C') in 110 mL of 7:1 v/v ethanol/water was mixed with a solution of 28.6 g (0.0799 mol) of O,O'-(+)-dibenzoyltartaric acid ((+)-DBT) (28.6g, 0.0799mol) in 110 mL of 7:1 v/v ethanol: water and the resulting solution was allowed to age at room temperature. Ethyl acetate (220ml) was added after crystallization was
15 complete and the resulting mixture was cooled to -20 °C and filtered to afford 32.4 g of methyl (S)-(4-fluorophenyl) glycinate, (+)-DBT salt (ee = 93.2%). The mother liquors were concentrated in vacuo and the free base was liberated by partitioning between ethyl acetate and aqueous sodium carbonate solution. A solution of free base, so
20 obtained, in 110 mL of 7:1 v/v ethanol/water was mixed with a solution of 28.6 g (0.0799 mol) of O,O'-(-)-dibenzoyltartaric acid ((-)-DBT) (28.6g, 0.0799mol) in 110 mL of 7:1 v/v ethanol: water and the resulting solution was allowed to age at room temperature. Ethyl acetate (220ml) was added after crystallization was complete and the
25 resulting mixture was cooled to -20 °C and filtered to afford 47.0 g of methyl (R)-(4-fluorophenyl) glycinate, (-)-DBT salt (ee = 75.8%). Recycling of the mother liquors and addition of (+)-DBT gave a second crop of 7.4 g of (S)-(4-fluorophenyl) glycinate, (+)-DBT salt (ee =
96.4%). The two crops of the (S)-amino ester (39.8 g) were combined
30 in 200 mL of 7:1 v/v ethanol/water, heated for 30 minutes and cooled to room temperature. Addition of ethyl acetate, cooling, and filtration afforded 31.7 g of (S)-(4-fluorophenyl) glycinate, (+)-DBT salt (ee>98%). Enantiomeric excesses was determined by chiral HPLC

- 184 -

(Crownpak CR(+)) 5% MeOH in aqHClO₄ pH2 1.5ml/min 40°C
200nm).

5 A mixture of 17.5 g of (S)-(4-fluorophenyl) glycinate, (+)-
DBT salt and 32 mL of 5.5 N HCl (32ml) was heated at reflux for 1.5
hours. The reaction mixture was concentrated in vacuo and the residue
was dissolved in 40 mL of water. The aqueous solution was washed 3 x
30 mL of ethyl acetate and the layers were separated. The pH of the
aqueous layer was adjusted to 7 using ammonium hydroxide and the
10 precipitated solid was filtered to afford 7.4 g of the title compound (ee
= 98.8%).

EXAMPLE 59

15 3-(S)-(4-Fluorophenyl)-4-benzyl-2-morpholinone

Step A: N-Benzyl (S)-(4-fluorophenyl)glycine

A solution of 1.87 g (11.05 mmol) of (S)-(4-fluorophenyl)-
glycine (from Example 58) and 1.12 mL (11.1 mmol) of benzaldehyde
20 in 11.1 mL of 1 N aqueous sodium hydroxide solution and 11 mL of
methanol at 0°C was treated with 165 mg (4.4 mmol) of sodium
borohydride. The cooling bath was removed and the resulting mixture
was stirred at room temperature for 30 minutes. Second portions of
benzaldehyde (1.12 mL (11.1 mmol)) and sodium borohydride 165 mg
25 (4.4 mmol) were added to the reaction mixture and stirring was
continued for 1.5 hours. The reaction mixture was partitioned between
100 mL of ether and 50 mL of water and the layers were separated.
The aqueous layer was separated and filtered to remove a small amount
of insoluble material. The filtrate was acidified to pH 5 with 2 N
aqueous hydrochloric acid solution and the solid that had precipitated
30 was filtered, rinsed well with water, then ether, and dried to afford 1.95
g of the title compound.
¹H-NMR (400 MHz, D₂O + NaOD): δ 3.33 (AB q, 2 H, J = 8.4), 3.85
(s, 1 H), 6.79-7.16 (m, 4 H).

- 185 -

Step B: 3-(S)-(4-Fluorophenyl)-4-benzyl-2-morpholinone

A mixture of 1.95 g (7.5 mmol) of N-benzyl (S)-(4-fluorophenyl)glycine, 3.90 mL (22.5 mmol) of N,N-diisopropylethylamine, 6.50 mL (75.0 mmol) of 1,2-dibromoethane and 40 mL of N,N-dimethylformamide was stirred at 100°C for 20 hours (dissolution of all solids occurred on warming). The reaction mixture was cooled and concentrated in vacuo. The residue was partitioned between 250 mL of ether and 100 mL of 0.5 N potassium hydrogen sulfate solution and the layers were separated. The organic layer was washed with 100 mL of saturated aqueous sodium bicarbonate solution, 3 x 150 mL of water, dried over magnesium sulfate, and concentrated in vacuo. Flash chromatography on 125 g of silica gel using 3:1 v/v hexanes/ether as the eluant afforded 1.58 g (74%) of the title compound as an oil.

¹H-NMR (400 MHz, CDCl₃): δ 2.65 (dt, 1 H, J = 3.2, 12.8), 3.00 (dt, 1 H, J = 12.8, 2.8), 3.16 (d, 1 H, J = 13.6), 3.76 (d, 1 H, J = 13.6), 4.24 (s, 1 H), 4.37 (dt, 1 H, J = 13.2, 3.2), 4.54 (dt, 1 H, J = 2.8, 13.2), 7.07-7.56 (m, 9 H).

EXAMPLE 602-(S)-(3,5-Bis(trifluoromethyl)benzyloxy)-3-(S)-(4-fluorophenyl)-4-benzylmorpholine

The title compound was prepared in 72% yield from 3-(S)-(4-fluorophenyl)-4-benzyl-2-morpholinone (from Example 59) using procedures analogous to those in Example 15, Steps A and B.

¹H-NMR (200 MHz, CDCl₃): δ 2.37 (dt, 1 H, J = 3.6, 11.8), 2.83-2.90 (m, 2 H), 3.55-3.63 (m, 2 H), 3.85 (d, 1 H, J = 13.4), 4.14 (dt, 1 H, J = 2.0, 11.8), 4.44 (d, 1 H, J = 13.6), 4.66 (d, 1 H, J = 2.8), 4.79 (d, 1 H, J = 13.4), 7.00-7.70 (12 H).

EXAMPLE 612-(S)-(3,5-Bis(trifluoromethyl)benzyloxy)-3-(S)-(4-fluorophenyl)morpholine

- 186 -

The title compound was prepared in 70% yield from 2-(S)-(3,5-bis(trifluoromethyl)benzyloxy)-3-(S)-(4-fluorophenyl)-4-benzylmorpholine (from Example 60) using a procedure analogous to that in Example 15, Step C.

- 5 Mass Spectrum (FAB): m/z 424 (M+H, 40%).
1H-NMR (400 MHz, CDCl₃): δ 1.80 (br s, 1 H), 3.11 (app dd, 1 H, J = 2.2, 12.4), 3.25 (dt, 1 H, J = 3.6, 12.4), 3.65 (app dd, 1 H, J = 3.6, 11.4), 4.05 (dt, 1 H, J = 2.2, 11.8), 4.11 (d, 1 H, J = 2.2), 4.53 (d, 1 H, J = 13.6), 4.71 (d, 1 H, J = 2.2), 4.83 (d, 1 H, J = 13.6), 7.04 (t, 2 H, J = 7.2), 7.33-7.37 (m, 2 H), 7.42 (s, 2 H), 7.72 (s, 1 H).
- 10

EXAMPLE 62

- 15 2-(S)-(3,5-Bis(trifluoromethyl)benzyloxy)-3-(S)-(4-fluorophenyl)-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methylmorpholine

The title compound was prepared in 69% yield from 2-(S)-(3,5-bis(trifluoromethyl)benzyloxy)-3-(S)-(4-fluorophenyl)morpholine (from Example 61) using a procedure analogous to that in Example 18.

- 20 Mass Spectrum (FAB): m/z 521 (M+H, 100%).
1H-NMR (400 MHz, CDCl₃): δ 2.55 (dt, 1 H, J = 3.6, 12.0), 2.91 (d, 1 H, J = 11.6), 2.93 (d, 1 H, J = 14.4), 3.57 (d, 1 H, J = 2.8), 3.59 (d, 1 H, J = 14.4), 3.67-3.70 (m, 1 H), 4.18 (dt, 1 H, J = 2.4, 11.6), 4.48 (d, 1 H, J = 13.6), 4.65 (d, 1 H, J = 2.8), 4.84 (d, 1 H, J = 13.6), 7.07 (t, 2 H, J = 8.4), 7.40 (s, 2 H), 7.45-7.48 (m, 2 H), 7.68 (s, 1 H), 10.04 (br s, 1 H), 10.69 (br s, 1 H).
- 25

Analysis: Calcd for C₂₂H₁₉F₇N₄O₃:

C, 50.78; H, 3.68; N, 10.77; F, 25.55

Found: C, 50.89; H, 3.76; N, 10.62; F, 25.56

30

EXAMPLE 63

2-(S)-(3,5-Bis(trifluoromethyl)benzyloxy)-4-((3-pyridyl)methyl carbonyl)-3-(R)-phenylmorpholine

- 187 -

A solution of 55 mg (0.315 mmol) of 4-pyridylacetic acid in 1 mL of CH₂Cl₂, containing 0.079 mL (0.715 mmol) of N-methylmorpholine, 53 mg (0.37 mmol) of HOBt and 73 mg (0.37 mmol) of EDC was stirred for 10 min. A solution of 2-(S)-(3,5-bis(trifluoromethyl)benzyloxy)-3-(R)-phenylmorpholine (from Example 33) in 1 mL of CH₂Cl₂ was added. After stirring the mixture for 2 h, it was partitioned between water and CH₂Cl₂. The organic layer was washed with water, brine and dried by filtering through Na₂SO₄. The filtrate was concentrated and the residue was purified by flash chromatography using 70% EtOAc/hexane to furnish 152 mg (100 % yield) of the product.

¹H-NMR (400 MHz, CDCl₃): δ 3.0-3.85 (m, 5H), 3.95 & 4.4 (br s, 1H), 4.66 (d, J = 13 Hz, 1H), 4.82 (d, J = 13 Hz, 1H), 5.0 & 5.9 (br s, 1H), 5.23 (s, 1H), 7.1-7.65 (m, 7H), 7.8 (m, 3 H), 8.43 (br s, 2H).

EXAMPLE 64

2-(S)-(3,5-Bis(trifluoromethyl)benzyloxy)-4-(methoxycarbonylpentyl)-3-(R)-phenylmorpholine

To a solution of 0.259 g (0.64 mmol) of 2-(S)-(3,5-bis(trifluoromethyl)benzyloxy)-3-(R)-phenylmorpholine (from example 33) in 2 mL of DMF were added 0.16 g (0.77 mmol) of methyl 6-bromohexanoate, 0.155 g (1.12 mmol) of K₂CO₃ and 2 crystals of nBu₄NI. The resulting solution was heated in a 60°C bath for 36 h, at which time a tlc indicated incomplete reaction. The bath temperature was raised to 100°C. After 3 h the reaction mixture was cooled and diluted with EtOAc. The EtOAc solution was washed with water (2x), brine and dried over Na₂SO₄. The filtrate was concentrated and the residue was chromatographed using 30% EtOAc/hexane to isolate 220 mg (65%) of the product.

¹H-NMR (400 MHz, CDCl₃): δ 1.0-1.4 (m, 4 H), 1.47 (m, J = 8 Hz, 2H), 1.95 (m, 1H), 2.2 (t, J = 8 Hz, 2H), 2.35 (m, 2H), 2.9 (d, J = 13 Hz, 1H), 3.07 (d, J = 7 Hz, 1H), 3.62 (s, 3H), 3.81 (td, J = 8 Hz and 2 Hz, 1 H), 4.04 (dd, J = 10 Hz and 2 Hz, 1H), 4.36 (d, J = 7 Hz, 1H), 4.4

- 188 -

(d, J = 13 Hz, 1H), 4.79 (d, J = 13 Hz, 1H), 7.2-7.4 (m, 7H), 7.66 (s, 1H).

EXAMPLE 65

5 2-(S)-(3,5-Bis(trifluoromethyl)benzyloxy)-4-(carboxypentyl)-3-(R)-phenylmorpholine

A solution of 0.15 g (0.28 mmol) of 2-(S)-(3,5-Bis(trifluoromethyl)benzyloxy)-4-(methoxycarbonylpentyl)-3-(R)-phenylmorpholine (from Example 64) in 3 mL of MeOH was saponified
10 by treating with 0.5 mL of 5 N NaOH for 40 min at 65°C. The solution was cooled, concentrated and the residue was diluted with water. The aqueous solution was adjusted to pH 6 by adding 2 N HCl and it was extracted with EtOAc. The organic layer was washed with brine, dried and concentrated. The residue upon chromatography on a flash column
15 with 50% EtOAc/hexane furnished 0.13g (89%) of the product
1H-NMR (400 MHz, CDCl₃): δ 1.0-1.5 (m, 4H), 1.5 (m, 2H), 2.2 (m, 2H), 2.35 (m, 2H), 2.9 (d, J = 13 Hz, 1H), 3.08 (d, J = 7 Hz, 1H), 3.82 (t, J = 8 Hz, 1H), 4.09 (d, J = 7 Hz, 1H), 4.38 (s, 1H), 4.4 (d, J = 13 Hz, 1H), 4.79 (d, J = 13 Hz, 1H), 7.2-7.4 (m, 7H), 7.66 (s, 1H).
20

EXAMPLE 66

25 2-(S)-(3,5-Bis(trifluoromethyl)benzyloxy)-4-(methylaminocarbonylpentyl)-6-oxo-hexyl)-3-(R)-phenylmorpholine

A solution of 116 mg (0.22 mmol) of 2-(S)-(3,5-Bis(trifluoromethyl)benzyloxy)-4-(carboxypentyl)-3-(R)-phenylmorpholine (from Example 65) in 1 mL of CH₂Cl₂ was treated with 40 mg (0.29 mmol) of HOBt, 57 mg (0.29 mmol) of EDC and 0.037 mL of N-methylmorpholine. After 10 min 0.027 mL (0.3
30 mmol) of aqueous methylamine (40%) was added and the resulting mixture was stirred for 4 h. The reaction mixture was diluted with water and extracted with CH₂Cl₂. The combined CH₂Cl₂ layer was washed with water, brine and dried over Na₂SO₄, and the filtrate was

- 189 -

concentrated. Purification of the residue on a flash column with EtOAc furnished 0.10 g of the product.

¹H-NMR (400 MHz, CDCl₃): δ 1.0-1.4 (m, 4 H), 1.47 (m, 2H), 1.95 (m, 1H), 2.04 (t, J = 8 Hz, 2H), 2.35 (m, 2H), 2.74 (d, J = 5 Hz, 3 H), 2.89(d, J =12 Hz, 1H) 3.08 (d, J = 7 Hz, 1H), 3.81 (t, J = 7 Hz, 1H), 4.02 (d, J = 11 Hz, 1H), 4.36 (d, J = 7 Hz, 1H), 4.39 (d, J = 13 Hz, 1H), 4.79 (d, J =13 Hz, 1H), 5.03 (br s, 1H), 7.2-7.4 (m, 7H), 7.65 (s, 1H).

EXAMPLE 67

2-(R)-(3,5-Bis(trifluoromethyl)benzoyloxy)-3-(S)-phenyl-4-benzyl morpholine

A solution of 2.67 g (10.0 mmol) of 3-(S)-phenyl-4-benzyl-2-morpholinone (from Example 14) in 40 mL of dry THF was cooled to -78°C. The cold solution was treated with 12.5 mL of 1.0 M L-Selectride®, solution in THF, maintaining the internal reaction temperature below -70°C. The resulting solution was stirred cold for 45 minutes and the reaction was charged with 3.60 mL (20.0 mmol) of 3,5-bis(trifluoro-methyl)benzoyl chloride. The resulting yellow mixture was stirred cold for 30 minutes and the reaction was quenched with 50 mL of saturated aqueous sodium bicarbonate solution. The quenched mixture was partitioned between 300 mL of ether and 50 mL of water and the layers were separated. The organic layer was dried over magnesium sulfate. The aqueous layer was extracted with 300 mL of ether; the extract was dried and combined with the original organic layer. The combined organics were concentrated *in vacuo*. Flash chromatography on 150 g of silica gel using 37:3 v/v hexanes/ether as the eluant afforded 4.06 g (80%) of the title compound as a solid.

¹H NMR (CDCl₃, 200 MHz, ppm): δ 2.50 (dt, J= 3.4, 12.0, 1H), 2.97 (app d, J= 12.0, 1H), 2.99 (d, J= 13.6, 1H), 3.72-3.79 (m, 1H), 3.82 (d, J= 2.6, 1H), 4.00 (d, J= 13.6, 1H), 4.20 (dt, J= 2.4, 11.6), 6.22 (d, J= 2.6, 1H), 7.22-7.37 (m, 7H), 7.57 (app d, J= 6.8, 2H), 8.07 (s, 1H), 8.47 (s, 2H).

Analysis Calcd for C₂₆H₂₁F₆NO₃:

- 190 -

Found: C, 61.29; H, 4.16; N, 2.75; F, 22.38.
C, 61.18; H, 4.14; N, 2.70; F, 22.13.

EXAMPLE 68

5

2-(R)-(1-(3,5-Bis(trifluoromethyl)phenyl)ethenyloxy)-3-(S)-phenyl-4-benzyl morpholine

10 Step A: Dimethyl titanocene

A solution of 2.49 g (10.0 mmol) of titanocene dichloride in 50 mL of ether in the dark at 0°C was treated with 17.5 mL of 1.4 M methyllithium solution in ether maintaining the internal temperature below 5°C. The resulting yellow/orange mixture was stirred at room temperature for 30 minutes and the reaction was quenched by slowly adding 25 g of ice. The quenched reaction mixture was diluted with 50 mL of ether and 25 mL of water and the layers were separated. The organic layer was dried over magnesium sulfate and concentrated in vacuo to afford 2.03 g (98%) of the title compound as a light-sensitive solid. The dimethyl titanocene could be stored as a solution in toluene at 0°C for at least 2 weeks without apparent chemical degradation. ¹H NMR (CDCl₃, 200 MHz, ppm): δ -0.15 (s, 6H), 6.06 (s, 10H).

25 Step B: 2-(R)-(1-(3,5-Bis(trifluoromethyl)phenyl)ethenyloxy)-3-(S)-phenyl-4-benzyl morpholine

A solution of 2.50 g (4.9 mmol) of 2-(R)-(3,5-bis(trifluoro-methyl)benzoyloxy)-3-(S)-phenyl-4-benzyl morpholine (from Example 67) and 2.50 g (12.0 mmol) of dimethyl titanocene (from Example 68, Step A) in 35 mL of 1:1 v/v THF/toluene was stirred in an oil bath at 80°C for 16 hours. The reaction mixture was cooled and concentrated in vacuo. Flash chromatography on 150 g of silica gel using 3:1 v/v hexanes/methylene chloride as the eluant afforded 1.71 g (69%) of the title compound as a solid. Mass Spectrum (FAB): m/Z 508 (M+H, 25%).

- 191 -

¹H NMR (CDCl₃, 400 MHz, ppm): δ 2.42 (dt, J= 3.6, 12.0, 1H), 2.89 (app d, J= 11.6), 2.92 (d, J= 13.6, 1H), 3.61-3.66 (m, 1H), 3.73 (d, J= 2.8), 1H), 4.00 (d, J= 13.6, 1H), 4.09 (dt, J= 2.4, 11.6, 1H), 4.75 (d, J= 2.8, 1H), 4.79 (d, J= 2.8, 1H), 5.36 (d, J= 2.4, 1H), 7.23-7.41 (m, 7H), 7.63 (app d, J= 7.2, 2H), 7.79 (s, 1H), 7.91 (s, 2H).

Analysis Calcd for C₂₇H₂₃F₆NO₂:

C, 63.90; H, 4.57; N, 2.76; F, 22.46.

Found: C, 63.71; H, 4.53; N, 2.68; F, 22.66.

EXAMPLE 69

2-(R)-(1-(S)-(3,5-Bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-phenyl morpholine and 2-(S)-(1-(R)-(3,5-Bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-phenyl morpholine

A mixture of 1.50 g (2.9 mmol) of 2-(R)-(1-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-phenyl-4-benzyl morpholine (from Example 68) and 750 mg 10% palladium on carbon catalyst in 25 mL of 3:2 v/v isopropanol/ethyl acetate was stirred under an atmosphere of hydrogen for 48 hours. The catalyst was filtered onto a pad of Celite; the reaction flask and filter pad were rinsed with 500 mL of ethyl acetate. The filtrate was concentrated in vacuo. Flash chromatography on 60 g of silica gel using 2:1 v/v hexanes/ether, then 2:1 v/v hexanes/ether afforded 106 mg of 2-(R)-(1-(S)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-phenyl morpholine and 899 mg of 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-phenyl morpholine, both as oils (84% total yield).

For 2-(R)-(1-(R)-(3,5-Bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-phenyl morpholine:

Mass Spectrum (CI): m/Z 420 (M⁺, 20%), 178 (100%).

¹H NMR (CDCl₃, 400 MHz, ppm): δ 1.46 (d, J= 6.8), 1.92 (br s, 1H), 3.13 (dd, J= 3.0, 12.6, 1H), 3.24 (dt, J= 3.6, 12.6, 1H), 3.62 (dd, J= 3.6, 11.2), 4.04 (d, J= 2.4, 1H), 4.14 (dt, J= 3.0, 11.2, 1H), 4.48 (d, J= 2.4, 1H), 4.90 (q, J= 6.8, 1H), 7.21-7.32 (m, 7H), 7.64 (s, 1H).

Analysis Calcd for C₂₀H₁₉F₆NO₂:

- 192 -

Found: C, 57.28; H, 4.57; N, 3.34; F, 27.18.
C, 57.41; H, 4.61; N, 3.29; F, 27.23.

EXAMPLE 70

5

2-(R)-(1-(R)-(3,5-Bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1,2,4-triazolo)methyl)morpholine

10 Step A: 2-(R)-(1-(R)-(3,5-Bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-phenyl-4-(2-(N-methylcarboxy-acetamidrazono)morpholine

A solution of 945 mg (2.3 mmol) of 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-phenyl morpholine (from Example 69), 447 mg (2.7 mmol) of N-methylcarboxy-2-chloro-acetamidrazone (from Example 45, Step A), and 0.78 mL (4.5 mmol) of N,N-diisopropylethylamine in 17 mL of acetonitrile was stirred at room temperature for 20 hours. The reaction was concentrated in vacuo and the residue was partitioned between 50 mL of methylene chloride and 25 mL of water. The organic layer was separated, dried over magnesium sulfate and concentrated in vacuo. Flash chromatography on 50 g of silica gel using 50:1:0.1 methylene chloride/methanol/ammonium hydroxide as the eluant afforded 1.12 g (90%) of the title compound as a foam.

25 Step B: 2-(R)-(1-(R)-(3,5-Bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1,2,4-triazolo)methyl)morpholine

A solution of 1.01 g (1.8 mmol) of 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-phenyl-4-(2-(N-methylcarboxyacetamidrazono)morpholine (from Example 70, Step A) in 15 mL of xylenes was heated at reflux for 2 hours. The reaction was cooled and concentrated in vacuo. Flash chromatography on 50 g of silica gel using 50:1:0.1 methylene chloride/methanol/ammonium hydroxide as the eluant afforded 781 mg (76%) of the title compound as a solid.

30

- 193 -

Mass Spectrum (FAB) m/Z 517 (M+H, 18%), 178 (100%).

^1H NMR (CDCl_3 , 400 MHz, ppm): δ 1.47 (d, $J=6.8$), 2.01-2.05 (m, 2H), 2.55 (dt, $J=3.6, 12.0$, 1H), 2.91 (d, $J=10.8$, 1H), 2.95 (d, $J=14.8$, 1H), 3.49 (d, $J=2.4$, 1H), 3.65 (d, $J=14.8$, 1H), 3.69 (d, $J=10.8$, 1H), 4.29 (dt, $J=2.4, 10.0$), 4.38 (d, $J=2.8$, 1H), 4.88 (q, $J=6.8$, 1H), 7.14 (s, 2H), 7.33-7.40 (m, 5H), 7.62 (s, 1H), 9.91 (br s, 1H), 10.16 (br s, 1H).

Analysis Calcd for $\text{C}_{23}\text{H}_{22}\text{F}_6\text{N}_4\text{O}_3$:

C, 53.49; H, 4.06; N, 10.85; F, 22.07.

Found: C, 53.64; H, 4.33; N, 10.81; F, 22.27.

EXAMPLE 71

2-(R)-(1-(S)-(3,5-Bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1,2,4-triazolo)methyl)morpholine

The title compound was prepared in 32% yield from 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-phenyl morpholine (from Example 69) using a procedure analogous to Example 70.

Mass Spectrum (FAB): m/Z 517 (M+H, 100%), 259 (50%).

^1H NMR (CDCl_3 , 400 MHz, ppm): δ 1.09 (d, $J=6.4$, 3H), 2.47-2.53 (m, 1H), 2.83 (app d, $J=11.6$, 1H), 2.95 (d, $J=14.0$, 1H), 3.51-3.65 (m, 3H), 4.01 (app t, $J=11.6$, 1H), 4.60 (q, $J=6.4$, 1H), 4.84 (d, $J=2.4$, 1H), 7.33-7.51 (m, 5H), 7.74 (s, 2H), 7.76 (s, 1H), 9.51 (br s, 1H), 10.00 (br s, 1H).

EXAMPLE 72

2-(R)-(3,5-Bis(trifluoromethyl)benzoyloxy)-3-(S)-(4-fluoro)phenyl-4-benzyl morpholine

The title compound was prepared in 83% yield from 3-(R)-(4-fluoro)phenyl-4-benzyl-2-morpholinone (from Example 59) using a procedure analogous to Example 67.

Mass Spectrum (FAB): m/Z 528 (M+H, 25%), 270 (100%).

- 194 -

¹H NMR (CDCl₃, 400 MHz, ppm): δ 2.50 (dt, J= 3.2, 12.0, 1H), 2.96 (app d, J= 12.0, 1H), 2.98 (d, J= 13.6, 1H), 3.74-3.78 (m, 1H), 3.81 (d, J= 2.8, 1H), 3.94 (d, J= 13.6, 1H), 4.19 (dt, J= 2.0, 12.0), 6.20 (d, J= 2.8, 1H), 6.99 (t, J= 8.4, 2H), 7.27-7.38 (m, 5H), 7.52-7.56 (m, 2H), 8.09 (s, 1H), 8.46 (s, 2H).

EXAMPLE 73

2-(R)-(1-(3,5-Bis(trifluoromethyl)phenyl)ethenyloxy)-3-(S)-(4-fluoro)-phenyl-4-benzyl morpholine

The title compound was prepared in 60% yield from 2-(R)-(3,5-bis(trifluoromethyl)benzoyloxy)-3-(S)-(4-fluoro)phenyl-4-benzyl morpholine (Example 72) using a procedure analogous to Example 68. Mass Spectrum (FAB): m/Z 526 (M+H, 75%), 270 (100%).

¹H NMR (CDCl₃, 400 MHz, ppm): δ 2.42 (dt, J= 3.6, 12.0), 2.90 (app d, J= 12.0, 1H), 2.91 (d, J= 13.6, 1H), 3.62-3.66 (m, 1H), 3.72 (d, J= 2.6), 3.94 (d, J= 13.6, 1H), 4.09 (dt, J= 2.4, 12.0, 1H), 4.75 (d, J= 3.2, 1H), 4.82 (d, J= 3.2, 1H), 5.32 (d, J= 2.6, 1H), 7.09 (t, J= 8.8, 2H), 7.24-7.33 (m, 5H), 7.58-7.62 (m, 2H), 7.80 (s, 1H), 7.90 (s, 2H).

EXAMPLE 74

2-(R)-(1-(S)-(3,5-Bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl morpholine and 2-(S)-(1-(R)-(3,5-Bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl morpholine

A mixture of 1.83 g (3.5 mmol) of 2-(R)-(1-(3,5-bis(trifluoromethyl)phenyl)ethenyloxy)-3-(S)-(4-fluoro)phenyl-4-benzyl morpholine (from Example 73) and 800 mg 5% rhodium on alumina catalyst in 40 mL of absolute ethanol was stirred under an atmosphere of hydrogen for 24 hours. The catalyst was filtered onto a pad of Celite; the reaction flask and filter cake were rinsed with 200 mL of ethyl acetate. The filtrate was concentrated in vacuo and the residue was pumped under high vacuum (1 mmHg, room temperature) to dryness.

- 195 -

The residue was redissolved in 40 mL of isopropanol; 800 mg of 10% palladium on carbon catalyst was added and the resulting mixture was stirred under an atmosphere of hydrogen for 24 hours. The catalyst was filtered onto a pad of Celite; the reaction flask and filter cake were rinsed with 200 mL of ethyl acetate. The filtrate was concentrated in vacuo. Flash chromatography on 50 g of silica gel using 2:1 v/v hexanes/ether, then 3:2 v/v ether/hexanes as the eluant afforded 283 mg of 2-(R)-(1-(S)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl morpholine and 763 mg of 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl morpholine, both as oils (total yield 68%).

For 2-(R)-(1-(S)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl morpholine:

Mass Spectrum (FAB) m/Z 438 (M+H, 65%), 180 (100%).

¹H NMR (CDCl₃, 400 MHz, ppm): δ 1.47 (d, J= 6.8, 3H), 1.87 (br s, 1H), 3.03 (dd, J= 2.8, 12.8), 3.17 (dt, J= 4.0, 12.4, 1H), 3.43-3.47 (m, 1H), 3.80 (dt, J= 3.2, 11.6), 4.10 (d, J= 2.2, 1H), 4.70 (q, J= 6.8, 1H), 4.87 (d, J= 2.2, 1H), 6.99-7.03 (m, 2H), 7.23-7.27 (m, 2H), 7.63 (s, 2H), 7.66 (s, 1H).

For 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl morpholine:

Mass Spectrum (FAB) m/Z 438 (M+H, 75%), 180 (100%).

¹H NMR (CDCl₃, 400 MHz, ppm): δ 1.16 (d, J= 6.8), 1.80 (br s, 1H), 3.13 (dd, J= 3.2, 12.4), 3.23 (dt, J= 3.6, 12.4), 3.63 (dd, J= 2.4, 11.2), 4.01 (d, J= 2.4, 1H), 4.13 (dt, J= 3.2, 12.0), 4.42 (d, J= 2.4, 1H), 4.19 (q, J= 6.8, 1H), 7.04-7.09 (m, 2H), 7.27-7.40 (m, 4H), 7.73 (s, 1H).

EXAMPLE 75

2-(R)-(1-(R)-(3,5-Bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(4-fluoro)-phenyl-4-(3-(5-oxo-1,2,4-triazolo)methyl)morpholine

The title compound was prepared in 79% yield from 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl

- 196 -

morpholine (from Example 74) using a procedure analogous to Example 70.

Mass Spectrum (FAB): m/z 535 (M+H, 100%), 277 (60%).

^1H NMR ($\text{CDCl}_3 + \text{CD}_3\text{OD}$, 400 MHz, ppm): δ 1.48 (d, $J = 6.8$, 3H),
2.52 (app t, $J = 10.4$, 1H), 2.85-2.88 (m, 2H), 3.47 (d, $J = 2.8$, 1H), 3.63
(d, $J = 14.4$, 1H), 3.70 (dd, $J = 2.0$, 11.6, 1H), 4.24 (app t, $J = 10.8$, 1H),
4.35 (d, $J = 2.8$, 1H), 4.91 (q, $J = 6.8$, 1H), 7.07 (app t, $J = 8.4$, 2H), 7.15
(s, 2H), 7.37-7.40 (m, 2H), 7.65 (s, 1H).

Analysis Calcd for $\text{C}_{23}\text{H}_{21}\text{F}_7\text{N}_4\text{O}_3$:
C, 51.69; H, 3.96; N, 10.48; F, 24.88.
Found: C, 51.74; H, 4.04; N, 10.50; F, 24.59.

EXAMPLE 76

2-(R)-(1-(S)-(3,5-Bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(4-fluoro)-
phenyl-4-(3-(5-oxo-1,2,4-triazolo)methyl)morpholine

The title compound was prepared in 60% yield from 2-(R)-(1-(S)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl
morpholine (from Example 74) using a procedure analogous to
Example 70.

Mass Spectrum (FAB): m/z 535 (M+H, 50%), 293 (100%).

^1H NMR ($\text{CDCl}_3 + \text{CD}_3\text{OD}$, 400 MHz, ppm): δ 1.11 (d, $J = 6.4$, 3H),
2.49 (dt, $J = 2.4$, 11.2), 2.83 (app d, $J = 11.2$, 1H), 2.95 (d, $J = 14.4$, 1H),
2.48-2.58 (m, 3H), 3.99 (app t, $J = 9.6$, 1H), 4.61 (q, $J = 6.4$, 1H), 4.81
(d, $J = 2.4$, 1H), 7.09 (t, $J = 8.8$, 2H), 7.50-7.53 (m, 2H), 7.75 (app s,
3H), 10.40 (br s, 1H), 11.00 (br s, 1H).

EXAMPLE 77

2-(R)-(1-(R)-(3-(Trifluoromethyl)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(5-
oxo-1,2,4-triazolo)methyl)morpholine

Step A: 2-(R)-(1-(R)-(3-(Trifluoromethyl)phenyl)ethoxy)-3-(S)-
phenyl morpholine

- 197 -

The title compound was prepared in 25% yield from 3-(S)-phenyl-4-benzyl-2-morpholinone (from Example 14) using procedures analogous to Examples 67-69.

5 ¹H NMR (CDCl₃, 400 MHz, ppm): δ 1.39 (d, J= 6.6, 3H), 1.93 (br s, 1H), 3.10 (dd, J= 3.0, 12.7, 1H), 3.20 (dt, J= 3.6, 12.4, 1H), 3.58 (ddd, J= 1.1, 3.8, 11.2, 1H), 4.00 (d, J= 2.4, 1H), 4.12 (dt, J= 3.0, 11.2, 1H), 4.44 (d, J= 2.4, 1H), 4.79 (q, J= 6.6, 1H), 6.72 (d, J=7.7, 1H), 7.01 (s, 1H), 7.09 (t, J=7.7, 1H), 7.18-7.25 (m, 2H), 7.25-7.3 (m, 3H), 7.34 (d, 10 J=7.7, 1H).

Analysis:

Calcd for C₁₉H₁₉F₃N₁O₂: C-65.14 H-5.47 N-4.00 F-16.27

Found: C-64.89 H-5.73 N-3.83 F-15.95

15 Step B: 2-(R)-(1-(R)-(3-(Trifluoromethyl)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1,2,4-triazolo)methylmorpholine

The title compound was prepared in 90% yield from 2-(R)-(1-(R)-(3-(trifluoromethyl)phenyl)ethoxy)-3-(S)-phenyl morpholine (from Example 77, Step A) using a procedure analogous to Example 70.

20 ¹H NMR (CDCl₃, 400 MHz, ppm): δ 1.40 (d, J= 6.3, 3H), 2.53 (br t, J=11.2, 1H), 2.86 (app d, J= 12.2, 1H), 2.94 (d, J= 14.3, 1H), 3.44 (br s, 1H), 3.63 (br d, J=14, 2H), 4.27 (app t, J= 11.5, 1H), 4.34 (d, J=2.1, 1H), 4.76 (q, J= 6.7, 1H), 6.63 (d, J=7.7, 1H), 6.93 (s, 1H), 7.06 (t, J= 25 7.6, 1H), 7.25-7.45 (m, 6H), 9.63 (br s, 1H), 9.74 (br s, 1H).

Analysis:

Calcd for C₂₂H₂₂F₃N₄O₃: C-59.06 H-4.96 N-12.52 F-12.74

Found: C-58.84 H-5.17 N-12.37 F-12.50

30

EXAMPLE 78

2-(R)-(1-(R)-(3-(Fluoro)-5-(trifluoromethyl)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1,2,4-triazolo)methylmorpholine

- 198 -

Step A: 2-(R)-(1-(R)-(3-(Fluoro)-5-(trifluoromethyl)phenyl)-
ethoxy)-3-(S)-phenyl-morpholine

The title compound was prepared in 44% yield from 3-(S)-phenyl-4-benzyl-2-morpholinone (from Example 14) using
5 procedures analogous to Examples 67-69.

¹H NMR (CDCl₃, 400 MHz, ppm): δ 1.38 (d, J= 6.6, 3H), 1.90 (br s, 1H), 3.17 (dd, J= 3.0, 12.7, 1H), 3.18 (dt, J= 3.6, 12.7, 1H), 3.58 (ddd, J= 1.1, 3.8, 11.2, 1H), 4.02 (d, J= 2.3, 1H), 4.11 (dt, J= 3.0, 11.2, 1H),
10 4.44 (d, J= 2.3, 1H), 4.78 (q, J= 6.6, 1H), 6.29 (d, J=9.2, 1H), 6.85 (s, 1H), 7.03 (d, J=8.4, 1H), 7.18-7.26 (m, 2H), 7.26-7.3 (m, 3H).

Analysis:

Calcd for C₁₉H₁₈F₄N₁O₂: C-61.95 H-4.93 N-3.80 F-20.63
15 Found: C-61.78 H-5.14 N-3.71 F-20.35

Step B: 2-(R)-(1-(R)-(3-(Fluoro)-5-(trifluoromethyl)phenyl)-
ethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1,2,4-triazolo)methyl)-
morpholine

The title compound was prepared in 77% yield from 2-(R)-(1-(R)-(3-(fluoro)-5-(trifluoromethyl)phenyl)ethoxy)-3-(S)-phenyl
20 morpholine (from Example 78, Step A) using a procedure analogous to Example 70.

¹H NMR (CDCl₃, 400 MHz, ppm): δ 1.40 (d, J= 6.3, 3H), 2.54 (br t, J=11, 1H), 2.87 (app d, J= 12, 1H), 2.94 (app d, J= 14, 1H), 3.47 (br s, 1H), 3.63 (br t, J=14, 2H), 4.25 (app t, J= 11, 1H), 4.35 (d, J=1.5, 1H),
25 4.75 (q, J= 6.3, 1H), 6.62 (d, J=6.7, 1H), 6.78 (s, 1H), 7.01 (d J= 8.4, 1H), 7.24 (d, J=3.9, 1H), 7.35 (br s, 4H), 9.61 (br s, 1H), 9.89 (br s, 1H).

Analysis:

30 Calcd for C₂₂H₂₁F₄N₄O₃: C-56.77 H-4.55 N-12.04 F-16.33
Found: C-56.57 H-4.65 N-11.94 F-16.13

- 199 -

EXAMPLE 792-(S)-(3-Fluoro-5-trifluoromethyl)benzoyloxy)-3-(S)-(4-fluoro)phenyl-4-benzyl morpholine

5 The title compound was prepared in 57% yield from 3-(S)-(4-fluoro)phenyl-4-benzyl-2-morpholinone (from Example 59) using a procedure analogous to Example 67.

Mass Spectrum (CI): m/Z 478 (M+H, 100%)

10 ¹H NMR (CDCl₃, 360 MHz, ppm): δ 2.50 (dt, J= 3.3, 12.0, 1H),
2.96(d, J= 12.0, 1H), 2.98 (d, J= 13.6, 1H), 3.75 (dd, J= 1.7, 11.5, 1H),
3.80 (d, J= 13.6, 1H), 3.75 (dd, J=1.7, 11.5, 1H), 3.80 (d, J= 2.5, 1H),
3.92 (d, J= 13.6, 1H), 4.19 (dt, J= 2.1, 12.0, 1H), 6.20 (d, J= 2.5, 1H),
6.99 (t, J= 8.7, 2H), 7.2-7.37 (m, 5H), 7.51-7.55 (m, 3H), 7.89 (d, J= 8.4, 1H), 8.09 (s, 1H).

EXAMPLE 802-(S)-(1-(3-Fluoro-5-trifluoromethyl)phenyl)ethenyloxy)-3-(S)-(4-fluoro)-phenyl-4-benzyl morpholine

20 The title compound was prepared in 85% yield from 2-(S)-(3-fluoro-5-trifluoromethyl)benzoyloxy)-3-(S)-(4-fluoro)phenyl-4-benzyl morpholine (from Example 79) using a procedure analogous to Example 68.

Mass Spectrum (CI): m/Z 476 (M+H, 100%)

25 ¹H NMR (CDCl₃, 360 MHz, ppm): δ 2.42 (dt, J= 3.6, 12.0 Hz, 1H),
2.90 (d, J= 12.0, 1H), 2.91 (d, J=13.6, 1H), 3.60-3.62 (m, 1H), 3.72 (d, J= 2.6, 1H), 3.92 (d, J= 13.6, 1H), 4.09 (dt, J= 2.4, 12.0, 1H), 4.67 (d, J= 2.9, 1H) 4.76 (d, J= 2.9, 1H), 5.28 (d, J= 2.6, 1H), 7.07 (t, J=8.7, 2H), 7.2-7.37 (m, 7H), 7.53 (s, 1H), 7.57-7.61 (m, 2H).

EXAMPLE 81

- 200 -

2-(S)-(1-(S)-(3-Fluoro-5-trifluoromethyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl morpholine and 2-(S)-(1-(R)-(3-Fluoro-5-trifluoromethyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl morpholine

5 The title compounds were prepared from 2-(S)-(1-(3-fluoro-5-trifluoromethyl)phenyl) ethenyloxy)-3-(S)-(4-fluoro)-phenyl-4-benzyl morpholine (from Example 80) using a procedure analogous to Example 74, but using 10% palladium on charcoal as the catalyst.

For 2-(S)-(1-(S)-(3-Fluoro-5-trifluoromethyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl morpholine:

10 Mass Spectrum (CI): m/z 388 ($M+H$, 100%)

1H NMR ($CDCl_3$, 360 MHz, ppm): δ 1.12 (d, $J=6.5$, 1H), 1.83 (s, 1H), 3.02 (d, $J=10.1$, 1H), 3.16 (dt, $J=3.6, 12.5$, 1H), 3.43 (dd, $J=2.7, 11.4$, 1H), 3.81 (dt, $J=2.9, 11.7$, 1H), 4.09 (d, $J=2.1$, 1H), 4.62 (q, $J=6.5$, 1H), 4.84 (d, $J=2.1$, 1H), 7.05 (t, $J=8.8$, 2H), 7.2 (d, $J=8.8$, 2H), 7.32 (s, 1H), 7.38 (dd, $J=5.5, 8.5$, 2H).

For 2-(S)-(1-(R)-(3-Fluoro-5-trifluoromethyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl morpholine:

Mass Spectrum (CI): m/z 387 (M^+ , 100%)

20 1H NMR ($CDCl_3$, 360 MHz, ppm): δ 1.42 (d, $J=6.6$, 3H), 1.91 (s, 1H), 3.11 (dd, $J=3.2, 12.4$, 1H), 3.22 (dt, $J=3.6, 12.4$, 1H), 3.58-3.62 (m, 1H), 4.01 (d, $J=2.3$, 1H), 4.11 (dt, $J=3.2, 12.0$, 1H), 4.41 (d, $J=2.3$, 1H), 4.80 (q, $J=6.6$, 1H), 6.41 (d, $J=9.2$, 1H), 6.86 (s, 1H), 7.02 (t, $J=8.7$, 2H), 7.08 (d, $J=9.2$, 2H), 7.21-7.26 (m, 2H).

25

EXAMPLE 82

2-(S)-(1-(R)-(3-Fluoro-5-trifluoromethyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1,2,4-triazolo)methyl) morpholine

30 The title compound was prepared from 2-(S)-(1-(R)-(3-fluoro-5-trifluoromethyl) phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl morpholine (from Example 81) using a procedure analogous to Example 70, mp 209-211 °C.

$[\alpha]_D = +65.1$ ($c = 1.0$, methanol)

- 201 -

¹H NMR (CDCl₃, 360 MHz, ppm): δ 1.32 (d, J= 6.4, 1H), 2.38 (t, J= 11.9, 1H), 2.76 (d, J= 13.9, 1H), 2.84 (d, J= 11.5, 1H), 3.32 (s, 1H), 3.40 (d, J=13.9, 1H), 3.49 (s, 1H), 3.61 (d, J= 11.2, 1H), 4.11 (t, J=11.3, 1H), 4.8 (q, J=6.4, 1H), 6.57 (d, J= 9.4, 1H), 6.94 (s, 1H), 7.1 (t, J= 8.7, 2H), 7.39 (d, J= 8.7, 2H), 7.51 (s, 2H), 11.26 (s, 1H), 11.38 (s, 1H).

EXAMPLE 83

2-(S)-(1-(R)-(3,5-Bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl morpholine

Step A: N,N-Diacetyl-4-bromomethyl-2-imidazolone

The title compound was prepared according to the procedure of Dolan and Dushinsky (Journal of the American Chemical Society, 70, 657 (1948)).

Step B: 2-(S)-(1-(R)-(3,5-Bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl-morpholine

A mixture of 1.00 g (2.28 mmol) of 2-(S)-(1-(R)-(3,5-bis(trifluoro-methyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl morpholine (from Example 74), 0.62 g (2.40 mmol) of N,N-diacetyl-4-bromomethyl-2-imidazolone (from Example 83, Step A) and 0.63 g (4.56 mmol) of potassium carbonate in 10 mL of N,N-dimethyl-formamide was stirred at room temperature for 15 minutes. The reaction was diluted with 100 mL of ethyl acetate and washed with water, saturated aqueous sodium chloride solution, dried over magnesium sulfate, and evaporated in vacuo. The resulting oil was dissolved in 10 mL of ethanol; the resulting solution was treated with 1.05 mL of 33% ethanolic methylamine solution and stirred at room temperature for 10 minutes. The reaction mixture was concentrated in vacuo to afford a solid. Recrystallisation from ethyl acetate/methanol afforded 0.63 g of the title compound, mp 192-194 °C.

- 202 -

¹H NMR (d₆-DMSO, 360 MHz, ppm): δ 1.35 (d, J= 6.5, 3H), 2.25 (dt, J=8.7, 1H), 2.60 (d, J= 13.8, 1H), 2.89 (d, J=11.6, 1H), 3.28-3.36 (m, 2H), 3.62 (d, J=10.2, 1H), 4.1 (t, J=10.0, 1H), 4.31 (d, J= 2.7, 1H), 4.92 (q, J= 6.5, 1H), 5.97 (s, 1H), 7.06 (t, J= 8.8, 2H), 7.36 (s, 2H), 7.65-7.85 (m, 2H), 7.84 (s, 1H), 9.58 (s, 1H), 9.8 (s, 1H).

EXAMPLE 84

2-(S)-(1-(R)-(3-Fluoro-5-(trifluoromethyl)phenyl)ethoxy)-3-(S)-(4-fluorophenyl)-4-(4-(2-oxo-1,3-imidazolo)methyl)morpholine

The title compound was prepared from 2-(S)-(1-(R)-(3-fluoro-5-trifluoromethyl) phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl morpholine (from Example 82) using a procedure analogous to Example 83, mp 209-210 °C.

[α]_D = +92.8 (c = 1.0 , methanol).

¹H NMR (d₆-DMSO, 360 MHz, ppm) δ 1.31 (d, J= 6.5, 3H), 2.24 (dt, J= 3.0, 11.9, 1H), 2.6 (d, J= 13.9, 1H), 3.61 (d, J= 11.2, 1H), 4.1 (t, J= 11.0, 1H), 4.29 (d, J= 2.3, 1H), 4.8 (q, J= 6.5, 1H), 6.00 (s, 1H), 6.55 (d, J= 9.3, 1H), 6.94 (s, 1H), 7.11 (t, J= 8.7, 2H), 7.39 (d, J= 8.4, 1H), 7.51 (s, 2H), 9.59 (s, 1H), 9.84 (s, 1H).

EXAMPLE 85

2-(S)-(1-(S)-(3,5-Bis(trifluoromethyl)phenyl)ethoxy)-3-(R)-(4-fluorophenyl)-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)morpholine

The title compound was prepared from (R)-(4-fluoro)phenylglycine using procedures analogous to Example 59, 67, 68, 69 and 70.

[α]_D = -67.7 (c = 0.7, MeOH, 20 °C)

EXAMPLE 86

The following compounds are prepared from 3-(S)-phenyl-4-benzyl-2-morpholinone (from Example 14) or 3-(S)-(4-fluoro)phenyl-

- 203 -

4-benzyl-2-morpholinone (from Example 59) using procedures analogous to Examples 15, 67 - 69 and 74. The hydrogenation of the 1-(substituted-aryl)ethenyloxy intermediates may be done with 10% palladium on carbon (Example 70) or 5% rhodium on alumina catalyst (Example 74) to give rapid reduction of the enol ether. Removal of the 4-benzyl substituent may be done catalytically under extended hydrogenation with 10% palladium on carbon or 5% rhodium on alumina catalyst or (when dehalogenation or cleavage of the ether might occur) in a second step with 1-chloroethyl chloroformate as in Example 44, Step C.

- 1) 2-(R)-(1-(R)-(3-(Chloro)-5-(trifluoromethyl)phenyl)ethoxy)-3-(S)-phenyl-morpholine;
- 2) 2-(R)-(1-(R)-(3,5-(Dimethyl)phenyl)ethoxy)-3-(S)-phenyl-morpholine;
- 3) 2-(R)-(1-(R)-(3-(Fluoro)-5-(methyl)phenyl)ethoxy)-3-(S)-phenyl-morpholine;
- 4) 2-(R)-(1-(R)-(3-(Chloro)-5-(methyl)phenyl)ethoxy)-3-(S)-phenyl-morpholine;
- 5) 2-(R)-(1-(R)-(3-(Bromo)-5-(methyl)phenyl)ethoxy)-3-(S)-phenyl-morpholine;
- 6) 2-(R)-(1-(R)-(3-(Isopropoxy)phenyl)ethoxy)-3-(S)-phenyl-morpholine;
- 7) 2-(R)-(1-(R)-(3-(Isopropoxy)-5-(trifluoromethyl)phenyl)ethoxy)-3-(S)-phenyl-morpholine;
- 8) 2-(R)-(1-(R)-(3-(Chloro)-5-(isopropoxy)phenyl)ethoxy)-3-(S)-phenyl-morpholine;
- 9) 2-(R)-(1-(R)-(3-(Fluoro)-5-(isopropoxy)phenyl)ethoxy)-3-(S)-phenyl-morpholine;
- 10) 2-(R)-(1-(R)-(3-(t-Butyl)phenyl)ethoxy)-3-(S)-phenyl-morpholine;
- 11) 2-(R)-(1-(R)-(3-(t-Butyl)-5-(chloro)phenyl)ethoxy)-3-(S)-phenyl-morpholine;

- 204 -

- 12) 2-(R)-(1-(R)-(3-(t-Butyl)-5-(trifluoromethyl)phenyl)ethoxy)-3-(S)-phenyl-morpholine;
- 13) 2-(R)-(1-(R)-(3,5-(Dimethyl)-4-(fluoro)phenyl)ethoxy)-3-(S)-phenyl-morpholine
- 5 14) 2-(R)-(1-(R)-(3,5-(Dimethyl)-4-(chloro)phenyl)ethoxy)-3-(S)-phenyl-morpholine;
- 15) 2-(R)-(1-(R)-(3,5-Bis(trifluoromethyl)-4-(fluoro)phenyl)ethoxy)-3-(S)-phenyl-morpholine;
- 10 16) 2-(R)-(1-(R)-(3,5-Bis(trifluoromethyl)-4-(chloro)phenyl)ethoxy)-3-(S)-phenyl-morpholine;
- 17) 2-(R)-(1-(R)-(3,5-(Dichloro)phenyl)ethoxy)-3-(S)-phenyl-morpholine;
- 18) 2-(R)-(1-(R)-(3,5-(Difluoro)phenyl)ethoxy)-3-(S)-phenyl-morpholine;
- 15 19) 2-(R)-(1-(R)-(1-(Naphthyl)ethoxy)-3-(S)-phenyl-morpholine;
- 20) 2-(R)-(1-(R)-(1-(4-(Fluoro)naphthyl))ethoxy)-3-(S)-phenyl-morpholine;
- 21) 2-(R)-(1-(R)-(1-(3-(Fluoro)naphthyl))ethoxy)-3-(S)-phenyl-morpholine;
- 20 22) 2-(R)-(1-(R)-(1-(3-(Chloro)naphthyl))ethoxy)-3-(S)-phenyl-morpholine;
- 23) 2-(R)-(1-(R)-(1-(3-(Methyl)naphthyl))ethoxy)-3-(S)-phenyl-morpholine;
- 25 24) 2-(R)-(1-(R)-(1-(3-(Trifluoromethyl)naphthyl))ethoxy)-3-(S)-phenyl-morpholine;
- 25) 2-(S)-(2-Fluoro-5-trifluoromethyl)benzyloxy-3-(S)-phenyl-morpholine;
- 26) 2-(S)-(2-Fluoro-5-trifluoromethyl)benzyloxy-3-(S)-(4-fluoro)phenyl-morpholine;
- 30 27) 2-(R)-(1-(R)-(2-Fluoro-5-trifluoromethyl)phenylethoxy)-3-(S)-phenyl-morpholine;
- 28) 2-(R)-(1-(R)-(2-Fluoro-5-trifluoromethyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-morpholine;

- 205 -

- 29) 2-(S)-(2-Chloro-5-trifluoromethyl)benzyloxy-3-(S)-phenyl-morpholine;
- 30) 2-(S)-(2-Chloro-5-trifluoromethyl)benzyloxy-3-(S)-(4-fluoro)phenyl-morpholine;
- 5 31) 2-(R)-(1-(R)-(2-Chloro-5-trifluoromethyl)phenylethoxy)-3-(S)-phenyl-morpholine;
- 32) 2-(R)-(1-(R)-(2-Chloro-5-trifluoromethyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-morpholine;
- 33) 2-(S)-(3-Methyl)benzyloxy-3-(S)-phenyl-morpholine;
- 10 34) 2-(S)-(3-Methyl)benzyloxy-3-(S)-(4-fluoro)phenyl-morpholine;
- 35) 2-(R)-(1-(R)-(3-Methyl)phenylethoxy)-3-(S)-phenyl-morpholine;
- 36) 2-(R)-(1-(R)-(3-Methyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-morpholine;
- 37) 2-(S)-(3-Bromo)benzyloxy-3-(S)-phenyl-morpholine;
- 15 38) 2-(S)-(3-Bromo)benzyloxy-3-(S)-(4-fluoro)phenyl-morpholine;
- 39) 2-(R)-(1-(R)-(3-Bromo)phenylethoxy)-3-(S)-phenyl-morpholine;
- 40) 2-(R)-(1-(R)-(3-Bromo)phenylethoxy)-3-(S)-(4-fluoro)phenyl-morpholine;
- 41) 2-(S)-(3-Chloro)benzyloxy-3-(S)-phenyl-morpholine;
- 20 42) 2-(S)-(3-Chloro)benzyloxy-3-(S)-(4-fluoro)phenyl-morpholine;
- 43) 2-(R)-(1-(R)-(3-Chloro)phenylethoxy)-3-(S)-phenyl-morpholine;
- 44) 2-(R)-(1-(R)-(3-Chloro)phenylethoxy)-3-(S)-(4-fluoro)phenyl-morpholine;
- 45) 2-(S)-(3-Trifluoromethyl)benzyloxy-3-(S)-phenyl-morpholine;
- 25 46) 2-(S)-(3-Trifluoromethyl)benzyloxy-3-(S)-(4-fluoro)phenyl-morpholine;
- 47) 2-(S)-(3-t-Butyl)benzyloxy-3-(S)-phenyl-morpholine;
- 48) 2-(S)-(3-t-Butyl)benzyloxy-3-(S)-(4-fluoro)phenyl-morpholine;
- 49) 2-(R)-(1-(R)-(3-(thiomethyl)phenyl)ethoxy)-3-(S)-phenyl-morpholine;
- 30 50) 2-(R)-(1-(R)-(3-(thiomethyl)-5-(trifluoromethyl)phenyl)ethoxy)-3-(S)-phenyl-morpholine;
- 51) 2-(R)-(1-(R)-(2,2-(dimethyl)-5-(thiomethyl)-2,3-dihydrobenzofuran-7-yl)ethoxy)-3-(S)-phenyl-morpholine;

- 206 -

- 52) 2-(R)-(1-(R)-(3,5-(dimethoxy)phenyl)ethoxy)-3-(S)-phenyl-morpholine;
- 53) 2-(R)-(1-(R)-(3-(trifluoromethyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-morpholine;
- 5 54) 2-(R)-(1-(R)-(3-(fluoro)-5-(trifluoromethyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-morpholine;
- 55) 2-(R)-(1-(R)-(3-(chloro)-5-(trifluoromethyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-morpholine;
- 10 56) 2-(R)-(1-(R)-(3,5-(dimethyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-morpholine;
- 57) 2-(R)-(1-(R)-(3-(fluoro)-5-(methyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-morpholine;
- 58) 2-(R)-(1-(R)-(3-(chloro)-5-(methyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-morpholine;
- 15 59) 2-(R)-(1-(R)-(3-(bromo)-5-(methyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-morpholine;
- 60) 2-(R)-(1-(R)-(3-(isopropoxy)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-morpholine;
- 20 61) 2-(R)-(1-(R)-(3-(isopropoxy)-5-(trifluoromethyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-morpholine;
- 62) 2-(R)-(1-(R)-(3-(chloro)-5-(isopropoxy)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-morpholine;
- 63) 2-(R)-(1-(R)-(3-(fluoro)-5-(isopropoxy)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-morpholine;
- 25 64) 2-(R)-(1-(R)-(3-(t-butyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-morpholine;
- 65) 2-(R)-(1-(R)-(3-(t-butyl)-5-(chloro)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-morpholine;
- 30 66) 2-(R)-(1-(R)-(3-(t-butyl)-5-(trifluoromethyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-morpholine;
- 67) 2-(R)-(1-(R)-(3,5-(dimethyl)-4-(fluoro)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-morpholine;
- 68) 2-(R)-(1-(R)-(3,5-(dimethyl)-4-(chloro)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-morpholine;

- 207 -

- 69) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)-4-(fluoro)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-morpholine;
- 70) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)-4-(chloro)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-morpholine;
- 5 71) 2-(R)-(1-(R)-(3,5-(dichloro)phenyl)ethoxy)-3-(S)-(4-fluoro)-phenyl-morpholine;
- 72) 2-(R)-(1-(R)-(3,5-(difluoro)phenyl)ethoxy)-3-(S)-(4-fluoro)-phenyl-morpholine;
- 10 73) 2-(R)-(1-(R)-(1-(naphthyl)ethoxy)-3-(S)-(4-fluoro)phenyl-morpholine;
- 74) 2-(R)-(1-(R)-(1-(4-(fluoro)naphthyl))ethoxy)-3-(S)-(4-fluoro)phenyl-morpholine;
- 75) 2-(R)-(1-(R)-(1-(3-(fluoro)naphthyl))ethoxy)-3-(S)-(4-fluoro)phenyl-morpholine;
- 15 76) 2-(R)-(1-(R)-(1-(3-(chloro)naphthyl))ethoxy)-3-(S)-(4-fluoro)phenyl-morpholine;
- 77) 2-(R)-(1-(R)-(1-(3-(methyl)naphthyl))ethoxy)-3-(S)-(4-fluoro)phenyl-morpholine;
- 20 78) 2-(R)-(1-(R)-(1-(3-(trifluoromethyl)naphthyl))ethoxy)-3-(S)-(4-fluoro)phenyl-morpholine;
- 79) 2-(R)-(1-(R)-(3-(thiomethyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-morpholine;
- 80) 2-(R)-(1-(R)-(3-(thiomethyl)-5-(trifluoromethyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-morpholine;
- 25 81) 2-(R)-(1-(R)-(2,2-(dimethyl)-5-(thiomethyl)-2,3-dihydro-benzofuran-7-yl)ethoxy)-3-(S)-(4-fluoro)phenyl-morpholine;
- 82) 2-(R)-(1-(R)-(3,5-(dimethoxy)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-morpholine;
- 30 83) 2-(R)-(1-(R)-(phenyl)ethoxy)-3-(S)-phenyl-morpholine;
- 84) 2-(R)-(1-(R)-(phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-morpholine;
- 85) 2-(R)-(1-(R)-(3-(fluoro)phenyl)ethoxy)-3-(S)-phenyl-morpholine;
- 86) 2-(R)-(1-(R)-(3-(fluoro)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-morpholine;
- 87) 2-(R)-(1-(R)-(4-(fluoro)phenyl)ethoxy)-3-(S)-phenyl-morpholine;

- 208 -

- 88) 2-(R)-(1-(R)-(4-(fluoro)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-morpholine;
- 89) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(3-fluoro)phenyl-morpholine;
- 5 90) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(3,4-difluoro)phenyl-morpholine;
- 91) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(3,4-dichloro)phenyl-morpholine;
- 10 92) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(3,4-dimethyl)phenyl-morpholine;
- 93) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-3,4-methylenedioxyphenyl-morpholine;
- 94) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(2-naphthyl)-morpholine.
- 15

EXAMPLE 87

The following compounds are prepared from the corresponding 2-(S)-(substituted-benzyloxy)-3-(S)-aryl morpholines or 2-(R)-(1-(R)-(substituted-aryl)ethoxy)-3-(S)-aryl morpholines (from Example 86) using procedures analogous to Examples 17, 18, 36, 38, 83 or, in the case of the 4-(5-tetrazolyl)methyl-substituted morpholines, by alkylation of the morpholine (from Example 86) with chloroacetonitrile in the presence of a tertiary amine base in acetonitrile, followed by formation of the final product by reacting the resulting nitrile with either sodium azide or trimethylsilylazide in an appropriate solvent.

20

25

- 1) 2-(R)-(1-(R)-(3-(Chloro)-5-(trifluoromethyl)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl-morpholine;
- 30 2) 2-(R)-(1-(R)-(3,5-(Dimethyl)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl-morpholine;
- 3) 2-(R)-(1-(R)-(3-(Fluoro)-5-(methyl)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1H,4H-1,2,4-triazolo)methyl-morpholine;

- 209 -

- 4) 2-(R)-(1-(R)-(3-(Chloro)-5-(methyl)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl-morpholine;
- 5) 2-(R)-(1-(R)-(3-(Bromo)-5-(methyl)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl-morpholine;
- 5 6) 2-(R)-(1-(R)-(3-(Isopropoxy)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl-morpholine;
- 7) 2-(R)-(1-(R)-(3-(Isopropoxy)-5-(trifluoromethyl)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl-morpholine;
- 10 8) 2-(R)-(1-(R)-(3-(Chloro)-5-(isopropoxy)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl-morpholine;
- 9) 2-(R)-(1-(R)-(3-(Fluoro)-5-(isopropoxy)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl-morpholine;
- 10 10) 2-(R)-(1-(R)-(1-(3-(trifluoromethyl)naphthyl))ethoxy)-3-(S)-phenyl-4-(3-(1,2,4-triazolo)methyl-morpholine;
- 11) 2-(R)-(1-(R)-(3-(t-Butyl)-5-(chloro)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl-morpholine;
- 12) 2-(R)-(1-(R)-(3-(t-Butyl)-5-(trifluoromethyl)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl-morpholine;
- 20 13) 2-(R)-(1-(R)-(3,5-(Dimethyl)-4-(fluoro)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl-morpholine;
- 14) 2-(R)-(1-(R)-(3,5-(Dimethyl)-4-(chloro)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl-morpholine;
- 25 15) 2-(R)-(1-(R)-(3,5-Bis(trifluoromethyl)-4-(fluoro)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl-morpholine;
- 16) 2-(R)-(1-(R)-(3,5-Bis(trifluoromethyl)-4-(chloro)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl-morpholine;
- 30 17) 2-(R)-(1-(R)-(3,5-(Dichloro)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl-morpholine;
- 18) 2-(R)-(1-(R)-(3,5-(Difluoro)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl-morpholine;

- 210 -

- 19) 2-(R)-(1-(R)-(1-(Naphthyl)ethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl-morpholine;
- 20) 2-(R)-(1-(R)-(1-(4-(Fluoro)naphthyl)ethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl-morpholine;
- 5 21) 2-(R)-(1-(R)-(1-(3-(Fluoro)naphthyl)ethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl-morpholine;
- 22) 2-(R)-(1-(R)-(1-(3-(Chloro)naphthyl)ethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl-morpholine;
- 10 23) 2-(R)-(1-(R)-(1-(3-(Methyl)naphthyl)ethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl-morpholine;
- 24) 2-(R)-(1-(R)-(1-(3-(Trifluoromethyl)naphthyl)ethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl-morpholine;
- 25) 2-(S)-(2-Fluoro-5-trifluoromethyl)benzyloxy-3-(S)-phenyl-4-(3-(1,2,4-triazolo)methyl-morpholine;
- 15 26) 2-(S)-(2-Fluoro-5-trifluoromethyl)benzyloxy-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl-morpholine;
- 27) 2-(R)-(1-(R)-(2-Fluoro-5-trifluoromethyl)phenylethoxy)-3-(S)-phenyl-4-(3-(1,2,4-triazolo)methyl-morpholine;
- 20 28) 2-(R)-(1-(R)-(2-Fluoro-5-trifluoromethyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl-morpholine;
- 29) 2-(S)-(2-Fluoro-5-trifluoromethyl)benzyloxy-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl-morpholine;
- 30) 2-(S)-(2-Fluoro-5-trifluoromethyl)benzyloxy-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl-morpholine;
- 25 31) 2-(R)-(1-(R)-(2-Fluoro-5-trifluoromethyl)phenylethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl-morpholine;
- 32) 2-(R)-(1-(R)-(2-Fluoro-5-trifluoromethyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl-morpholine;
- 30 33) 2-(S)-(2-Fluoro-5-trifluoromethyl)benzyloxy-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl-morpholine;
- 34) 2-(S)-(2-Fluoro-5-trifluoromethyl)benzyloxy-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl-morpholine;

- 211 -

- 35) 2-(R)-(1-(R)-(2-Fluoro-5-trifluoromethyl)phenylethoxy)-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl-morpholine;
- 36) 2-(R)-(1-(R)-(2-Fluoro-5-trifluoromethyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl-morpholine;
- 5 37) 2-(S)-(2-Fluoro-5-trifluoromethyl)benzyloxy-3-(S)-phenyl-4-(2-imidazolo)methyl-morpholine;
- 38) 2-(S)-(2-Fluoro-5-trifluoromethyl)benzyloxy-3-(S)-(4-fluoro)phenyl-4-(2-imidazolo)methyl-morpholine;
- 10 39) 2-(R)-(1-(R)-(2-Fluoro-5-trifluoromethyl)phenylethoxy)-3-(S)-phenyl-4-(2-imidazolo)methyl-morpholine;
- 40) 2-(R)-(1-(R)-(2-Fluoro-5-trifluoromethyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(2-imidazolo)methyl-morpholine;
- 41) 2-(S)-(2-Fluoro-5-trifluoromethyl)benzyloxy-3-(S)-phenyl-4-(4-imidazolo)methyl-morpholine;
- 15 42) 2-(S)-(2-Fluoro-5-trifluoromethyl)benzyloxy-3-(S)-(4-fluoro)phenyl-4-(4-imidazolo)methyl-morpholine;
- 43) 2-(R)-(1-(R)-(2-Fluoro-5-trifluoromethyl)phenylethoxy)-3-(S)-phenyl-4-(4-imidazolo)methyl-morpholine;
- 20 44) 2-(R)-(1-(R)-(2-Fluoro-5-trifluoromethyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-imidazolo)methyl-morpholine;
- 45) 2-(S)-(2-Fluoro-5-trifluoromethyl)benzyloxy-3-(S)-phenyl-4-(5-tetrazolo)methyl-morpholine;
- 46) 2-(S)-(2-Fluoro-5-trifluoromethyl)benzyloxy-3-(S)-(4-fluoro)phenyl-4-(5-tetrazolo)methyl-morpholine;
- 25 47) 2-(R)-(1-(R)-(2-Fluoro-5-trifluoromethyl)phenylethoxy)-3-(S)-phenyl-4-(5-tetrazolo)methyl-morpholine;
- 48) 2-(R)-(1-(R)-(2-Fluoro-5-trifluoromethyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(5-tetrazolo)methyl-morpholine;
- 30 49) 2-(S)-(2-Fluoro-5-trifluoromethyl)benzyloxy-3-(S)-phenyl-4-(2-oxo-5H-pyrrol-4-yl)methyl-morpholine;
- 50) 2-(S)-(2-Fluoro-5-trifluoromethyl)benzyloxy-3-(S)-(4-fluoro)phenyl-4-(2-oxo-5H-pyrrol-4-yl)methyl-morpholine;
- 51) 2-(R)-(1-(R)-(2-Fluoro-5-trifluoromethyl)phenylethoxy)-3-(S)-phenyl-4-(2-oxo-5H-pyrrol-4-yl)methyl-morpholine;

- 212 -

- 52) 2-(R)-(1-(R)-(2-Fluoro-5-trifluoromethyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(2-oxo-5H-pyrrol-4-yl)methyl-morpholine;
- 53) 2-(S)-(2-Chloro-5-trifluoromethyl)benzyloxy-3-(S)-phenyl-4-(3-(1,2,4-triazolo)methyl-morpholine;
- 5 54) 2-(S)-(2-Chloro-5-trifluoromethyl)benzyloxy-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl-morpholine;
- 55) 2-(R)-(1-(R)-(2-Chloro-5-trifluoromethyl)phenylethoxy)-3-(S)-phenyl-4-(3-(1,2,4-triazolo)methyl-morpholine;
- 10 56) 2-(R)-(1-(R)-(2-Chloro-5-trifluoromethyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl-morpholine;
- 57) 2-(S)-(2-Chloro-5-trifluoromethyl)benzyloxy-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl-morpholine;
- 58) 2-(S)-(2-Chloro-5-trifluoromethyl)benzyloxy-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl-morpholine;
- 15 59) 2-(R)-(1-(R)-(2-Chloro-5-trifluoromethyl)phenylethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl-morpholine;
- 60) 2-(R)-(1-(R)-(2-Chloro-5-trifluoromethyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl-morpholine;
- 20 61) 2-(S)-(2-Chloro-5-trifluoromethyl)benzyloxy-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl-morpholine;
- 62) 2-(S)-(2-Chloro-5-trifluoromethyl)benzyloxy-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl-morpholine;
- 25 63) 2-(R)-(1-(R)-(2-Chloro-5-trifluoromethyl)phenylethoxy)-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl-morpholine;
- 64) 2-(R)-(1-(R)-(2-Chloro-5-trifluoromethyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl-morpholine;
- 30 65) 2-(S)-(2-Chloro-5-trifluoromethyl)benzyloxy-3-(S)-phenyl-4-(2-imidazolo)methyl-morpholine;
- 66) 2-(S)-(2-Chloro-5-trifluoromethyl)benzyloxy-3-(S)-(4-fluoro)phenyl-4-(2-imidazolo)methyl-morpholine;
- 67) 2-(R)-(1-(R)-(2-Chloro-5-trifluoromethyl)phenylethoxy)-3-(S)-phenyl-4-(2-imidazolo)methyl-morpholine;

- 213 -

- 68) 2-(R)-(1-(R)-(2-Chloro-5-trifluoromethyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(2-imidazolo)methyl-morpholine;
- 69) 2-(S)-(2-Chloro-5-trifluoromethyl)benzyloxy-3-(S)-phenyl-4-(4-imidazolo)methyl-morpholine;
- 5 70) 2-(S)-(2-Chloro-5-trifluoromethyl)benzyloxy-3-(S)-(4-fluoro)phenyl-4-(4-imidazolo)methyl-morpholine;
- 71) 2-(R)-(1-(R)-(2-Chloro-5-trifluoromethyl)phenylethoxy)-3-(S)-phenyl-4-(4-imidazolo)methyl-morpholine;
- 72) 2-(R)-(1-(R)-(2-Chloro-5-trifluoromethyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-imidazolo)methyl-morpholine;
- 10 73) 2-(S)-(2-Chloro-5-trifluoromethyl)benzyloxy-3-(S)-phenyl-4-(5-tetrazolo)methyl-morpholine;
- 74) 2-(S)-(2-Chloro-5-trifluoromethyl)benzyloxy-3-(S)-(4-fluoro)phenyl-4-(5-tetrazolo)methyl-morpholine;
- 15 75) 2-(R)-(1-(R)-(2-Chloro-5-trifluoromethyl)phenylethoxy)-3-(S)-phenyl-4-(5-tetrazolo)methyl-morpholine;
- 76) 2-(R)-(1-(R)-(2-Chloro-5-trifluoromethyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(5-tetrazolo)methyl-morpholine;
- 77) 2-(S)-(2-Chloro-5-trifluoromethyl)benzyloxy-3-(S)-phenyl-4-(2-oxo-5H-pyrrol-4-yl)methyl-morpholine;
- 20 78) 2-(S)-(2-Chloro-5-trifluoromethyl)benzyloxy-3-(S)-(4-fluoro)phenyl-4-(2-oxo-5H-pyrrol-4-yl)methyl-morpholine;
- 79) 2-(R)-(1-(R)-(2-Chloro-5-trifluoromethyl)phenylethoxy)-3-(S)-phenyl-4-(2-oxo-5H-pyrrol-4-yl)methyl-morpholine;
- 25 80) 2-(R)-(1-(R)-(2-Chloro-5-trifluoromethyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(2-oxo-5H-pyrrol-4-yl)methyl-morpholine;
- 81) 2-(S)-(3-Methyl)benzyloxy-3-(S)-phenyl-4-(3-(1,2,4-triazolo)methyl-morpholine;
- 82) 2-(S)-(3-Methyl)benzyloxy-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl-morpholine;
- 30 83) 2-(R)-(1-(R)-(3-Methyl)phenylethoxy)-3-(S)-phenyl-4-(3-(1,2,4-triazolo)methyl-morpholine;
- 84) 2-(R)-(1-(R)-(3-Methyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl-morpholine;

- 214 -

- 85) 2-(S)-(3-Methyl)benzyloxy-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl-morpholine;
- 86) 2-(S)-(3-Methyl)benzyloxy-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl-morpholine;
- 5 87) 2-(R)-(1-(R)-(3-Methyl)phenylethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl-morpholine;
- 88) 2-(R)-(1-(R)-(3-Methyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl-morpholine;
- 10 89) 2-(S)-(3-Methyl)benzyloxy-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl-morpholine;
- 90) 2-(S)-(3-Methyl)benzyloxy-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl-morpholine;
- 91) 2-(R)-(1-(R)-(3-Methyl)phenylethoxy)-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl-morpholine;
- 15 92) 2-(R)-(1-(R)-(3-Methyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl-morpholine;
- 93) 2-(S)-(3-Methyl)benzyloxy-3-(S)-phenyl-4-(2-imidazolo)methyl-morpholine;
- 20 94) 2-(S)-(3-Methyl)benzyloxy-3-(S)-(4-fluoro)phenyl-4-(2-imidazolo)methyl-morpholine;
- 95) 2-(R)-(1-(R)-(3-Methyl)phenylethoxy)-3-(S)-phenyl-4-(2-imidazolo)methyl-morpholine;
- 96) 2-(R)-(1-(R)-(3-Methyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(2-imidazolo)methyl-morpholine;
- 25 97) 2-(S)-(3-Methyl)benzyloxy-3-(S)-phenyl-4-(4-imidazolo)methyl-morpholine;
- 98) 2-(S)-(3-Methyl)benzyloxy-3-(S)-(4-fluoro)phenyl-4-(4-imidazolo)methyl-morpholine;
- 30 99) 2-(R)-(1-(R)-(3-Methyl)phenylethoxy)-3-(S)-phenyl-4-(4-imidazolo)methyl-morpholine;
- 100) 2-(R)-(1-(R)-(3-Methyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-imidazolo)methyl-morpholine;
- 101) 2-(S)-(3-Methyl)benzyloxy-3-(S)-phenyl-4-(5-tetrazolo)methyl-morpholine;

- 215 -

- 102) 2-(S)-(3-Methyl)benzyloxy-3-(S)-(4-fluoro)phenyl-4-(5-tetrazolo)methyl-morpholine;
- 103) 2-(R)-(1-(R)-(3-Methyl)phenylethoxy)-3-(S)-phenyl-4-(5-tetrazolo)methyl-morpholine;
- 5 104) 2-(R)-(1-(R)-(3-Methyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(5-tetrazolo)methyl-morpholine;
- 105) 2-(S)-(3-Methyl)benzyloxy-3-(S)-phenyl-4-(2-oxo-5H-pyrrol-4-yl)methyl-morpholine;
- 10 106) 2-(S)-(3-Methyl)benzyloxy-3-(S)-(4-fluoro)phenyl-4-(2-oxo-5H-pyrrol-4-yl)methyl-morpholine;
- 107) 2-(R)-(1-(R)-(3-Methyl)phenylethoxy)-3-(S)-phenyl-4-(2-oxo-5H-pyrrol-4-yl)methyl-morpholine;
- 108) 2-(R)-(1-(R)-(3-Methyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(2-oxo-5H-pyrrol-4-yl)methyl-morpholine;
- 15 109) 2-(S)-(3-Bromo)benzyloxy-3-(S)-phenyl-4-(3-(1,2,4-triazolo)methyl-morpholine;
- 110) 2-(S)-(3-Bromo)benzyloxy-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl-morpholine;
- 20 111) 2-(R)-(1-(R)-(3-Bromo)phenylethoxy)-3-(S)-phenyl-4-(3-(1,2,4-triazolo)methyl-morpholine;
- 112) 2-(R)-(1-(R)-(3-Bromo)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl-morpholine;
- 113) 2-(S)-(3-Bromo)benzyloxy-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl-morpholine;
- 25 114) 2-(S)-(3-Bromo)benzyloxy-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl-morpholine;
- 115) 2-(R)-(1-(R)-(3-Bromo)phenylethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl-morpholine;
- 30 116) 2-(R)-(1-(R)-(3-Bromo)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl-morpholine;
- 117) 2-(S)-(3-Bromo)benzyloxy-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl-morpholine;
- 118) 2-(S)-(3-Bromo)benzyloxy-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl-morpholine;

- 216 -

- 119) 2-(R)-(1-(R)-(3-Bromo)phenylethoxy)-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl-morpholine;
120) 2-(R)-(1-(R)-(3-Bromo)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl-morpholine;
5 121) 2-(S)-(3-Bromo)benzyloxy-3-(S)-phenyl-4-(2-imidazolo)methyl-morpholine;
122) 2-(S)-(3-Bromo)benzyloxy-3-(S)-(4-fluoro)phenyl-4-(2-imidazolo)methyl-morpholine;
123) 2-(R)-(1-(R)-(3-Bromo)phenylethoxy)-3-(S)-phenyl-4-(2-imidazolo)methyl-morpholine;
10 124) 2-(R)-(1-(R)-(3-Bromo)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(2-imidazolo)methyl-morpholine;
125) 2-(S)-(3-Bromo)benzyloxy-3-(S)-phenyl-4-(4-imidazolo)methyl-morpholine;
15 126) 2-(S)-(3-Bromo)benzyloxy-3-(S)-(4-fluoro)phenyl-4-(4-imidazolo)methyl-morpholine;
127) 2-(R)-(1-(R)-(3-Bromo)phenylethoxy)-3-(S)-phenyl-4-(4-imidazolo)methyl-morpholine;
128) 2-(R)-(1-(R)-(3-Bromo)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-imidazolo)methyl-morpholine;
20 129) 2-(S)-(3-Bromo)benzyloxy-3-(S)-phenyl-4-(5-tetrazolo)methyl-morpholine;
130) 2-(S)-(3-Bromo)benzyloxy-3-(S)-(4-fluoro)phenyl-4-(5-tetrazolo)methyl-morpholine;
25 131) 2-(R)-(1-(R)-(3-Bromo)phenylethoxy)-3-(S)-phenyl-4-(5-tetrazolo)methyl-morpholine;
132) 2-(R)-(1-(R)-(3-Bromo)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(5-tetrazolo)methyl-morpholine;
133) 2-(S)-(3-Bromo)benzyloxy-3-(S)-phenyl-4-(2-oxo-5H-pyrrol-4-yl)methyl-morpholine;
30 134) 2-(S)-(3-Bromo)benzyloxy-3-(S)-(4-fluoro)phenyl-4-(2-oxo-5H-pyrrol-4-yl)methyl-morpholine;
135) 2-(R)-(1-(R)-(3-Bromo)phenylethoxy)-3-(S)-phenyl-4-(2-oxo-5H-pyrrol-4-yl)methyl-morpholine;

- 217 -

- 136) 2-(R)-(1-(R)-(3-Bromo)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(2-oxo-5H-pyrrol-4-yl)methyl-morpholine;
- 137) 2-(S)-(3-Chloro)benzyloxy-3-(S)-phenyl-4-(3-(1,2,4-triazolo)methyl-morpholine;
- 5 138) 2-(S)-(3-Chloro)benzyloxy-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl-morpholine;
- 139) 2-(R)-(1-(R)-(3-Chloro)phenylethoxy)-3-(S)-phenyl-4-(3-(1,2,4-triazolo)methyl-morpholine;
- 10 140) 2-(R)-(1-(R)-(3-Chloro)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl-morpholine;
- 141) 2-(S)-(3-Chloro)benzyloxy-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl-morpholine;
- 142) 2-(S)-(3-Chloro)benzyloxy-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl-morpholine;
- 15 143) 2-(R)-(1-(R)-(3-Chloro)phenylethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl-morpholine;
- 144) 2-(R)-(1-(R)-(3-Chloro)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl-morpholine;
- 20 145) 2-(S)-(3-Chloro)benzyloxy-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl-morpholine;
- 146) 2-(S)-(3-Chloro)benzyloxy-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl-morpholine;
- 147) 2-(R)-(1-(R)-(3-Chloro)phenylethoxy)-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl-morpholine;
- 25 148) 2-(R)-(1-(R)-(3-Chloro)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl-morpholine;
- 149) 2-(S)-(3-Chloro)benzyloxy-3-(S)-phenyl-4-(2-imidazolo)methyl-morpholine;
- 30 150) 2-(S)-(3-Chloro)benzyloxy-3-(S)-(4-fluoro)phenyl-4-(2-imidazolo)methyl-morpholine;
- 151) 2-(R)-(1-(R)-(3-Chloro)phenylethoxy)-3-(S)-phenyl-4-(2-imidazolo)methyl-morpholine;
- 152) 2-(R)-(1-(R)-(3-Chloro)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(2-imidazolo)methyl-morpholine;

- 218 -

- 153) 2-(S)-(3-Chloro)benzyloxy-3-(S)-phenyl-4-(4-imidazolo)methyl-morpholine;
- 154) 2-(S)-(3-Chloro)benzyloxy-3-(S)-(4-fluoro)phenyl-4-(4-imidazolo)methyl-morpholine;
- 5 155) 2-(R)-(1-(R)-(3-Chloro)phenylethoxy)-3-(S)-phenyl-4-(4-imidazolo)methyl-morpholine;
- 156) 2-(R)-(1-(R)-(3-Chloro)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-imidazolo)methyl-morpholine;
- 10 157) 2-(S)-(3-Chloro)benzyloxy-3-(S)-phenyl-4-(5-tetrazolo)methyl-morpholine;
- 158) 2-(S)-(3-Chloro)benzyloxy-3-(S)-(4-fluoro)phenyl-4-(5-tetrazolo)methyl-morpholine;
- 159) 2-(R)-(1-(R)-(3-Chloro)phenylethoxy)-3-(S)-phenyl-4-(5-tetrazolo)methyl-morpholine;
- 15 160) 2-(R)-(1-(R)-(3-Chloro)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(5-tetrazolo)methyl-morpholine;
- 161) 2-(S)-(3-Chloro)benzyloxy-3-(S)-phenyl-4-(2-oxo-5H-pyrrol-4-yl)methyl-morpholine;
- 20 162) 2-(S)-(3-Chloro)benzyloxy-3-(S)-(4-fluoro)phenyl-4-(2-oxo-5H-pyrrol-4-yl)methyl-morpholine;
- 163) 2-(R)-(1-(R)-(3-Chloro)phenylethoxy)-3-(S)-phenyl-4-(2-oxo-5H-pyrrol-4-yl)methyl-morpholine;
- 164) 2-(R)-(1-(R)-(3-Chloro)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(2-oxo-5H-pyrrol-4-yl)methyl-morpholine;
- 25 165) 2-(S)-(3-Trifluoromethyl)benzyloxy-3-(S)-phenyl-4-(3-(1,2,4-triazolo)methyl)morpholine;
- 166) 2-(S)-(3-Trifluoromethyl)benzyloxy-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl-morpholine;
- 30 167) 2-(R)-(1-(R)-(3-Trifluoromethyl)phenylethoxy)-3-(S)-phenyl-4-(3-(1,2,4-triazolo)methyl-morpholine;
- 168) 2-(R)-(1-(R)-(3-Trifluoromethyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl-morpholine;
- 169) 2-(S)-(3-Trifluoromethyl)benzyloxy-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl-morpholine;

- 219 -

- 170) 2-(S)-(3-Trifluoromethyl)benzyloxy-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl-morpholine;
- 171) 2-(R)-(1-(R)-(1-(3-(trifluoromethyl)naphthyl))ethoxy)-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl-morpholine;
- 5 172) 2-(R)-(1-(R)-(3-Trifluoromethyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl-morpholine;
- 173) 2-(S)-(3-Trifluoromethyl)benzyloxy-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl-morpholine;
- 10 174) 2-(S)-(3-Trifluoromethyl)benzyloxy-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl-morpholine;
- 175) 2-(R)-(1-(R)-(3-Trifluoromethyl)phenylethoxy)-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl-morpholine;
- 15 176) 2-(R)-(1-(R)-(3-Trifluoromethyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl-morpholine;
- 177) 2-(S)-(3-Trifluoromethyl)benzyloxy-3-(S)-phenyl-4-(2-imidazolo)methyl-morpholine;
- 178) 2-(S)-(3-Trifluoromethyl)benzyloxy-3-(S)-(4-fluoro)phenyl-4-(2-imidazolo)methyl-morpholine;
- 20 179) 2-(R)-(1-(R)-(3-Trifluoromethyl)phenylethoxy)-3-(S)-phenyl-4-(2-imidazolo)methyl-morpholine;
- 180) 2-(R)-(1-(R)-(3-Trifluoromethyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(2-imidazolo)methyl-morpholine;
- 25 181) 2-(S)-(3-Trifluoromethyl)benzyloxy-3-(S)-phenyl-4-(4-imidazolo)methyl-morpholine;
- 182) 2-(S)-(3-Trifluoromethyl)benzyloxy-3-(S)-(4-fluoro)phenyl-4-(4-imidazolo)methyl-morpholine;
- 183) 2-(R)-(1-(R)-(3-Trifluoromethyl)phenylethoxy)-3-(S)-phenyl-4-(4-imidazolo)methyl-morpholine;
- 30 184) 2-(R)-(1-(R)-(3-Trifluoromethyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-imidazolo)methyl-morpholine;
- 185) 2-(S)-(3-Trifluoromethyl)benzyloxy-3-(S)-phenyl-4-(5-tetrazolo)methyl-morpholine;

- 220 -

- 186) 2-(S)-(3-Trifluoromethyl)benzyloxy-3-(S)-(4-fluoro)phenyl-4-(5-tetrazolo)methyl-morpholine;
- 187) 2-(R)-(1-(R)-(3-Trifluoromethyl)phenylethoxy)-3-(S)-phenyl-4-(5-tetrazolo)methyl-morpholine;
- 5 188) 2-(R)-(1-(R)-(3-Trifluoromethyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(5-tetrazolo)methyl-morpholine;
- 189) 2-(S)-(3-Trifluoromethyl)benzyloxy-3-(S)-phenyl-4-(2-oxo-5H-pyrrol-4-yl)methyl-morpholine;
- 10 190) 2-(S)-(3-Trifluoromethyl)benzyloxy-3-(S)-(4-fluoro)phenyl-4-(2-oxo-5H-pyrrol-4-yl)methyl-morpholine;
- 191) 2-(R)-(1-(R)-(3-Trifluoromethyl)phenylethoxy)-3-(S)-phenyl-4-(2-oxo-5H-pyrrol-4-yl)methyl-morpholine;
- 192) 2-(R)-(1-(R)-(3-Trifluoromethyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(2-oxo-5H-pyrrol-4-yl)methyl-morpholine;
- 15 193) 2-(S)-(3-t-Butyl)benzyloxy-3-(S)-phenyl-4-(3-(1,2,4-triazolo)methyl-morpholine;
- 194) 2-(S)-(3-t-Butyl)benzyloxy-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl-morpholine;
- 20 195) 2-(R)-(1-(R)-(3-t-Butyl)phenylethoxy)-3-(S)-phenyl-4-(3-(1,2,4-triazolo)methyl-morpholine;
- 196) 2-(R)-(1-(R)-(3-t-Butyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl-morpholine;
- 197) 2-(S)-(3-t-Butyl)benzyloxy-3-(S)-phenyl-4-(3-(5-oxo-1,2,4-triazolo)methyl-morpholine;
- 25 198) 2-(S)-(3-t-Butyl)benzyloxy-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1,2,4-triazolo)methyl-morpholine;
- 199) 2-(R)-(1-(R)-(3-t-Butyl)phenylethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1,2,4-triazolo)methyl-morpholine;
- 30 200) 2-(R)-(1-(R)-(3-t-Butyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1,2,4-triazolo)methyl-morpholine;
- 201) 2-(S)-(3-t-Butyl)benzyloxy-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl-morpholine;
- 202) 2-(S)-(3-t-Butyl)benzyloxy-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl-morpholine;

- 221 -

- 203) 2-(R)-(1-(R)-(3-t-Butyl)phenylethoxy)-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl-morpholine;
204) 2-(R)-(1-(R)-(3-t-Butyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl-morpholine;
5 205) 2-(S)-(3-t-Butyl)benzyloxy-3-(S)-phenyl-4-(2-imidazolo)methyl-morpholine;
206) 2-(S)-(3-t-Butyl)benzyloxy-3-(S)-(4-fluoro)phenyl-4-(2-imidazolo)methyl-morpholine;
10 207) 2-(R)-(1-(R)-(3-t-Butyl)phenylethoxy)-3-(S)-phenyl-4-(2-imidazolo)methyl-morpholine;
208) 2-(R)-(1-(R)-(3-t-Butyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(2-imidazolo)methyl-morpholine;
209) 2-(S)-(3-t-Butyl)benzyloxy-3-(S)-phenyl-4-(4-imidazolo)methyl-morpholine;
15 210) 2-(S)-(3-t-Butyl)benzyloxy-3-(S)-(4-fluoro)phenyl-4-(4-imidazolo)methyl-morpholine;
211) 2-(R)-(1-(R)-(3-t-Butyl)phenylethoxy)-3-(S)-phenyl-4-(4-imidazolo)methyl-morpholine;
20 212) 2-(R)-(1-(R)-(3-t-Butyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-imidazolo)methyl-morpholine;
213) 2-(S)-(3-t-Butyl)benzyloxy-3-(S)-phenyl-4-(5-tetrazolo)methyl-morpholine;
214) 2-(S)-(3-t-Butyl)benzyloxy-3-(S)-(4-fluoro)phenyl-4-(5-tetrazolo)methyl-morpholine;
25 215) 2-(R)-(1-(R)-(3-t-Butyl)phenylethoxy)-3-(S)-phenyl-4-(5-tetrazolo)methyl-morpholine;
216) 2-(R)-(1-(R)-(3-t-Butyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(5-tetrazolo)methyl-morpholine;
30 217) 2-(S)-(3-t-Butyl)benzyloxy-3-(S)-phenyl-4-(2-oxo-5H-pyrrol-4-yl)methyl-morpholine;
218) 2-(S)-(3-t-Butyl)benzyloxy-3-(S)-(4-fluoro)phenyl-4-(2-oxo-5H-pyrrol-4-yl)methyl-morpholine;
219) 2-(R)-(1-(R)-(3-t-Butyl)phenylethoxy)-3-(S)-phenyl-4-(2-oxo-5H-pyrrol-4-yl)methyl-morpholine;

- 222 -

- 220) 2-(R)-(1-(R)-(3-t-Butyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(2-oxo-5H-pyrrol-4-yl)methyl-morpholine;
- 221) 2-(R)-(1-(R)-(3-(thiomethylphenyl)ethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl-morpholine;
- 5 222) 2-(R)-(1-(R)-(3-(thiomethylphenyl)ethoxy)-3-(S)-phenyl-4-(3-(1,2,4-triazolo)methyl-morpholine;
- 223) 2-(R)-(1-(R)-(3-(thiomethylphenyl)ethoxy)-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl-morpholine;
- 10 224) 2-(R)-(1-(R)-(3-(thiomethyl-5-(trifluoromethyl)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1,2,4-triazolo)methyl-morpholine;
- 225) 2-(R)-(1-(R)-(3-(thiomethyl-5-(trifluoromethyl)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(1,2,4-triazolo)methyl-morpholine;
- 226) 2-(R)-(1-(R)-(3-(thiomethyl-5-(trifluoromethyl)phenyl)ethoxy)-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl-morpholine;
- 15 227) 2-(R)-(1-(R)-(2,2-(dimethyl)-5-(thiomethyl)-2,3-dihydro-benzofuran-7-yl)ethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl-morpholine;
- 228) 2-(R)-(1-(R)-(2,2-(dimethyl)-5-(thiomethyl)-2,3-dihydro-benzofuran-7-yl)ethoxy)-3-(S)-phenyl-4-(3-(1,2,4-triazolo)-methyl-morpholine;
- 20 229) 2-(R)-(1-(R)-(2,2-(dimethyl)-5-(thiomethyl)-2,3-dihydro-benzofuran-7-yl)ethoxy)-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl-morpholine;
- 230) 2-(R)-(1-(R)-(3,5-(dimethoxy)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl-morpholine;
- 25 231) 2-(R)-(1-(R)-(3,5-(dimethoxy)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(1,2,4-triazolo)methyl-morpholine;
- 232) 2-(R)-(1-(R)-(3,5-(dimethoxy)phenyl)ethoxy)-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl-morpholine;
- 30 233) 2-(R)-(1-(R)-(3-(fluoro)-5-(trifluoromethyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl-morpholine;
- 234) 2-(R)-(1-(R)-(3-(fluoro)-5-(trifluoromethyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl-morpholine;

- 223 -

- 235) 2-(R)-(1-(R)-(3-(fluoro)-5-(trifluoromethyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl-morpholine;
- 236) 2-(R)-(1-(R)-(3-(chloro)-5-(trifluoromethyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl-morpholine;
- 5 237) 2-(R)-(1-(R)-(3-(chloro)-5-(trifluoromethyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl-morpholine;
- 238) 2-(R)-(1-(R)-(3-(chloro)-5-(trifluoromethyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl-morpholine;
- 10 239) 2-(R)-(1-(R)-(3,5-(dimethyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl-morpholine;
- 240) 2-(R)-(1-(R)-(3,5-(dimethyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl-morpholine;
- 15 241) 2-(R)-(1-(R)-(3,5-(dimethyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl-morpholine;
- 242) 2-(R)-(1-(R)-(3-(fluoro)-5-(methyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl-morpholine;
- 20 243) 2-(R)-(1-(R)-(3-(fluoro)-5-(methyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl-morpholine;
- 244) 2-(R)-(1-(R)-(3-(fluoro)-5-(methyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl-morpholine;
- 25 245) 2-(R)-(1-(R)-(3-(chloro)-5-(methyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl-morpholine;
- 246) 2-(R)-(1-(R)-(3-(chloro)-5-(methyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl-morpholine;
- 30 247) 2-(R)-(1-(R)-(3-(chloro)-5-(methyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl-morpholine;
- 248) 2-(R)-(1-(R)-(3-(bromo)-5-(methyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl-morpholine;

- 224 -

- 249) 2-(R)-(1-(R)-(3-(bromo)-5-(methyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl-morpholine;
- 250) 2-(R)-(1-(R)-(3-(bromo)-5-(methyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl-morpholine;
- 5 251) 2-(R)-(1-(R)-(3-(isopropoxy)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl-morpholine;
- 252) 2-(R)-(1-(R)-(3-(isopropoxy)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl-morpholine;
- 10 253) 2-(R)-(1-(R)-(3-(isopropoxy)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl-morpholine;
- 254) 2-(R)-(1-(R)-(3-(isopropoxy)-5-(trifluoromethyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl-morpholine;
- 15 255) 2-(R)-(1-(R)-(3-(isopropoxy)-5-(trifluoromethyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl-morpholine;
- 256) 2-(R)-(1-(R)-(3-(isopropoxy)-5-(trifluoromethyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl-morpholine;
- 20 257) 2-(R)-(1-(R)-(3-(chloro)-5-(isopropoxy)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl-morpholine;
- 258) 2-(R)-(1-(R)-(3-(chloro)-5-(isopropoxy)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl-morpholine;
- 25 259) 2-(R)-(1-(R)-(3-(chloro)-5-(isopropoxy)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl-morpholine;
- 260) 2-(R)-(1-(R)-(3-(fluoro)-5-(isopropoxy)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl-morpholine;
- 30 261) 2-(R)-(1-(R)-(3-(fluoro)-5-(isopropoxy)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl-morpholine;
- 262) 2-(R)-(1-(R)-(3-(fluoro)-5-(isopropoxy)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl-morpholine;

- 225 -

- 263) 2-(R)-(1-(R)-(3-(t-butyl)-5-(chloro)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl-morpholine;
- 5 264) 2-(R)-(1-(R)-(3-(t-butyl)-5-(chloro)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl-morpholine;
- 265) 2-(R)-(1-(R)-(3-(t-butyl)-5-(chloro)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl-morpholine;
- 266) 2-(R)-(1-(R)-(3-(t-butyl)-5-(trifluoromethyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl-morpholine;
- 10 267) 2-(R)-(1-(R)-(3-(t-butyl)-5-(trifluoromethyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl-morpholine;
- 268) 2-(R)-(1-(R)-(3-(t-butyl)-5-(trifluoromethyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl-morpholine;
- 15 269) 2-(R)-(1-(R)-(3,5-(dimethyl)-4-(fluoro)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl-morpholine;
- 270) 2-(R)-(1-(R)-(3,5-(dimethyl)-4-(fluoro)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl-morpholine;
- 20 271) 2-(R)-(1-(R)-(3,5-(dimethyl)-4-(fluoro)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl-morpholine;
- 272) 2-(R)-(1-(R)-(3,5-(dimethyl)-4-(chloro)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl-morpholine;
- 25 273) 2-(R)-(1-(R)-(3,5-(dimethyl)-4-(chloro)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl-morpholine;
- 274) 2-(R)-(1-(R)-(3,5-(dimethyl)-4-(chloro)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl-morpholine;
- 30 275) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)-4-(fluoro)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl-morpholine;
- 276) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)-4-(fluoro)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl-morpholine;

- 226 -

- 277) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)-4-(fluoro)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl-morpholine;
- 5 278) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)-4-(chloro)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl-morpholine;
- 279) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)-4-(chloro)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl-morpholine;
- 10 280) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)-4-(chloro)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl-morpholine;
- 281) 2-(R)-(1-(R)-(3,5-(dichloro)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl-morpholine;
- 15 282) 2-(R)-(1-(R)-(3,5-(dichloro)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl-morpholine;
- 283) 2-(R)-(1-(R)-(3,5-(dichloro)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl-morpholine;
- 20 284) 2-(R)-(1-(R)-(3,5-(difluoro)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl-morpholine;
- 285) 2-(R)-(1-(R)-(3,5-(difluoro)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl-morpholine;
- 25 286) 2-(R)-(1-(R)-(3,5-(difluoro)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl-morpholine;
- 287) 2-(R)-(1-(R)-(1-(naphthyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl-morpholine;
- 288) 2-(R)-(1-(R)-(1-(naphthyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl-morpholine;
- 30 289) 2-(R)-(1-(R)-(1-(naphthyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl-morpholine;
- 290) 2-(R)-(1-(R)-(1-(4-(fluoro)naphthyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl-morpholine;

- 227 -

- 291) 2-(R)-(1-(R)-(1-(4-(fluoro)naphthyl))ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl-morpholine;
- 292) 2-(R)-(1-(R)-(1-(4-(fluoro)naphthyl))ethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl-morpholine;
- 5 293) 2-(R)-(1-(R)-(1-(3-(fluoro)naphthyl))ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl-morpholine;
- 294) 2-(R)-(1-(R)-(1-(3-(fluoro)naphthyl))ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl-morpholine;
- 10 295) 2-(R)-(1-(R)-(1-(3-(fluoro)naphthyl))ethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl-morpholine;
- 296) 2-(R)-(1-(R)-(1-(3-(chloro)naphthyl))ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl-morpholine;
- 15 297) 2-(R)-(1-(R)-(1-(3-(chloro)naphthyl))ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl-morpholine;
- 298) 2-(R)-(1-(R)-(1-(3-(chloro)naphthyl))ethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl-morpholine;
- 20 299) 2-(R)-(1-(R)-(1-(3-(methyl)naphthyl))ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl-morpholine;
- 300) 2-(R)-(1-(R)-(1-(3-(methyl)naphthyl))ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl-morpholine;
- 25 301) 2-(R)-(1-(R)-(1-(3-(methyl)naphthyl))ethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl-morpholine;
- 302) 2-(R)-(1-(R)-(1-(3-(trifluoromethyl)naphthyl))ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl-morpholine;
- 30 303) 2-(R)-(1-(R)-(1-(3-(trifluoromethyl)naphthyl))ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl-morpholine;
- 304) 2-(R)-(1-(R)-(1-(3-(trifluoromethyl)naphthyl))ethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl-morpholine;

- 228 -

- 305) 2-(R)-(1-(R)-(3-(thiomethylphenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl-morpholine;
- 5 306) 2-(R)-(1-(R)-(3-(thiomethylphenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl-morpholine;
- 307) 2-(R)-(1-(R)-(3-(thiomethylphenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl-morpholine;
- 308) 2-(R)-(1-(R)-(3-(thiomethyl-5-(trifluoromethyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl-morpholine;
- 10 309) 2-(R)-(1-(R)-(3-(thiomethyl-5-(trifluoromethyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl-morpholine;
- 310) 2-(R)-(1-(R)-(3-(thiomethyl-5-(trifluoromethyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl-morpholine;
- 15 311) 2-(R)-(1-(R)-(2,2-(dimethyl)-5-(thiomethyl)-2,3-dihydrobenzofuran-7-yl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl-morpholine;
- 20 312) 2-(R)-(1-(R)-(2,2-(dimethyl)-5-(thiomethyl)-2,3-dihydrobenzofuran-7-yl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl-morpholine;
- 313) 2-(R)-(1-(R)-(2,2-(dimethyl)-5-(thiomethyl)-2,3-dihydrobenzofuran-7-yl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl-morpholine;
- 25 314) 2-(R)-(1-(R)-(3,5-(dimethoxy)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl-morpholine;
- 315) 2-(R)-(1-(R)-(3,5-(dimethoxy)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl-morpholine;
- 30 316) 2-(R)-(1-(R)-(3,5-(dimethoxy)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl-morpholine;
- 317) 2-(R)-(1-(R)-(phenyl)ethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl-morpholine;

- 229 -

- 318) 2-(R)-(1-(R)-(phenyl)ethoxy)-3-(S)-phenyl-4-(3-(1,2,4-triazolo)methyl-morpholine;
- 319) 2-(R)-(1-(R)-(phenyl)ethoxy)-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl-morpholine;
- 5 320) 2-(R)-(1-(R)-(phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl-morpholine;
- 321) 2-(R)-(1-(R)-(phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl-morpholine;
- 10 322) 2-(R)-(1-(R)-(phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl-morpholine;
- 323) 2-(R)-(1-(R)-(3-(fluoro)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl-morpholine;
- 324) 2-(R)-(1-(R)-(3-(fluoro)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(1,2,4-triazolo)methyl-morpholine;
- 15 325) 2-(R)-(1-(R)-(3-(fluoro)phenyl)ethoxy)-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl-morpholine;
- 326) 2-(R)-(1-(R)-(3-(fluoro)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl-morpholine;
- 20 327) 2-(R)-(1-(R)-(3-(fluoro)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl-morpholine;
- 328) 2-(R)-(1-(R)-(3-(fluoro)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl-morpholine;
- 329) 2-(R)-(1-(R)-(4-(fluoro)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl-morpholine;
- 25 330) 2-(R)-(1-(R)-(4-(fluoro)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(1,2,4-triazolo)methyl-morpholine;
- 331) 2-(R)-(1-(R)-(4-(fluoro)phenyl)ethoxy)-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl-morpholine;
- 30 332) 2-(R)-(1-(R)-(4-(fluoro)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl-morpholine;
- 332) 2-(R)-(1-(R)-(4-(fluoro)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl-morpholine;
- 333) 2-(R)-(1-(R)-(4-(fluoro)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl-morpholine;

- 230 -

- 334) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(3-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl-morpholine;
- 5 335) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(3-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl-morpholine;
- 336) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(3-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl-morpholine;
- 337) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(3,4-difluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl-morpholine;
- 10 338) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(3,4-difluoro)phenyl-4-(3-(1,2,4-triazolo)methyl-morpholine;
- 339) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(3,4-difluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl-morpholine;
- 15 340) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(3,4-dichloro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl-morpholine;
- 341) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(3,4-dichloro)phenyl-4-(3-(1,2,4-triazolo)methyl-morpholine;
- 20 342) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(3,4-dichloro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl-morpholine;
- 343) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(3,4-dimethyl)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl-morpholine;
- 25 344) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(3,4-dimethyl)phenyl-4-(3-(1,2,4-triazolo)methyl-morpholine;
- 345) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(3,4-dimethyl)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl-morpholine;
- 30 346) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-3,4-methylenedioxyphenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl-morpholine;
- 347) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-3,4-methylenedioxyphenyl-4-(3-(1,2,4-triazolo)methyl-morpholine;

- 348) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-3,4-methylenedioxyphenyl-4-(4-(2-oxo-1,3-imidazolo)methyl-morpholine;
- 5 349) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(2-naphthyl)-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl-morpholine;
- 350) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(2-naphthyl)-4-(3-(1,2,4-triazolo)methyl-morpholine;
- 351) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(2-naphthyl)-4-(4-(2-oxo-1,3-imidazolo)methyl-morpholine;
- 10 352) 2-(R)-(1-(R)-(3-(fluoro)-5-(trifluoromethyl)ethoxy)-3-(S)-phenyl-4-(3-(-1,2,4-triazolo)methyl-morpholine;
- 353) 2-(R)-(1-(R)-(3-(fluoro)-5-(trifluoromethyl)ethoxy)-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl-morpholine;
- 15 354) 2-(R)-(1-(R)-(3-(chloro)-5-(trifluoromethyl)ethoxy)-3-(S)-phenyl-4-(3-(-1,2,4-triazolo)methyl-morpholine;
- 355) 2-(R)-(1-(R)-(3-(chloro)-5-(trifluoromethyl)ethoxy)-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl-morpholine;
- 356) 2-(R)-(1-(R)-(3,5-(dimethyl)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(-1,2,4-triazolo)methyl-morpholine;
- 20 357) 2-(R)-(1-(R)-(3,5-(dimethyl)phenyl)ethoxy)-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl-morpholine;
- 358) 2-(R)-(1-(R)-(3-(fluoro)-5-(methyl)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(-1,2,4-triazolo)methyl-morpholine;
- 25 359) 2-(R)-(1-(R)-(3-(fluoro)-5-(methyl)phenyl)ethoxy)-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl-morpholine;
- 360) 2-(R)-(1-(R)-(3-(chloro)-5-(methyl)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(-1,2,4-triazolo)methyl-morpholine;
- 361) 2-(R)-(1-(R)-(3-(chloro)-5-(methyl)phenyl)ethoxy)-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl-morpholine;
- 30 362) 2-(R)-(1-(R)-(3-(bromo)-5-(methyl)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(-1,2,4-triazolo)methyl-morpholine;
- 363) 2-(R)-(1-(R)-(3-(bromo)-5-(methyl)phenyl)ethoxy)-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl-morpholine;

- 232 -

- 364) 2-(R)-(1-(R)-(3-(isopropoxy)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(-1,2,4-triazolo)methyl-morpholine;
- 365) 2-(R)-(1-(R)-(3-(isopropoxy)phenyl)ethoxy)-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl-morpholine;
- 5 366) 2-(R)-(1-(R)-(3-(isopropoxy)-5-(trifluoromethyl)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(-1,2,4-triazolo)methyl-morpholine;
- 367) 2-(R)-(1-(R)-(3-(isopropoxy)-5-(trifluoromethyl)phenyl)ethoxy)-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl-morpholine;
- 10 368) 2-(R)-(1-(R)-(3-(chloro)-5-(isopropoxy)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(-1,2,4-triazolo)methyl-morpholine;
- 369) 2-(R)-(1-(R)-(3-(chloro)-5-(isopropoxy)phenyl)ethoxy)-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl-morpholine;
- 370) 2-(R)-(1-(R)-(3-(fluoro)-5-(isopropoxy)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(-1,2,4-triazolo)methyl-morpholine;
- 15 371) 2-(R)-(1-(R)-(3-(fluoro)-5-(isopropoxy)phenyl)ethoxy)-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl-morpholine;
- 372) 2-(R)-(1-(R)-(3-(t-butyl)-5-(chloro)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(-1,2,4-triazolo)methyl-morpholine;
- 20 373) 2-(R)-(1-(R)-(3-(t-butyl)-5-(chloro)phenyl)ethoxy)-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl-morpholine;
- 374) 2-(R)-(1-(R)-(3-(t-butyl)-5-(trifluoromethyl)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(-1,2,4-triazolo)methyl-morpholine;
- 375) 2-(R)-(1-(R)-(3-(t-butyl)-5-(trifluoromethyl)phenyl)ethoxy)-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl-morpholine;
- 25 376) 2-(R)-(1-(R)-(3,5-(dimethyl)-4-(fluoro)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(-1,2,4-triazolo)methyl-morpholine;
- 377) 2-(R)-(1-(R)-(3,5-(dimethyl)-4-(fluoro)phenyl)ethoxy)-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl-morpholine;
- 30 378) 2-(R)-(1-(R)-(3,5-(dimethyl)-4-(chloro)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(-1,2,4-triazolo)methyl-morpholine;
- 379) 2-(R)-(1-(R)-(3,5-(dimethyl)-4-(chloro)phenyl)ethoxy)-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl-morpholine;
- 380) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)-4-(fluoro)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(-1,2,4-triazolo)methyl-morpholine;

- 233 -

- 381) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)-4-(fluoro)phenyl)ethoxy)-
3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl-morpholine;
382) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)-4-(chloro)phenyl)ethoxy)-
3-(S)-phenyl-4-(3-(-1,2,4-triazolo)methyl-morpholine;
5 383) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)-4-(chloro)phenyl)ethoxy)-
3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl-morpholine;
384) 2-(R)-(1-(R)-(3,5-(dichloro)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(-
1,2,4-triazolo)methyl-morpholine;
385) 2-(R)-(1-(R)-(3,5-(dichloro)phenyl)ethoxy)-3-(S)-phenyl-4-(4-(2-
10 oxo-1,3-imidazolo)methyl-morpholine;
386) 2-(R)-(1-(R)-(3,5-(difluoro)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(-
1,2,4-triazolo)methyl-morpholine;
387 2-(R)-(1-(R)-(3,5-(difluoro)phenyl)ethoxy)-3-(S)-phenyl-4-(4-(2-
oxo-1,3-imidazolo)methyl-morpholine;
15 388) 2-(R)-(1-(R)-(1-(naphthyl)ethoxy)-3-(S)-phenyl-4-(3-(-1,2,4-
triazolo)methyl-morpholine;
389) 2-(R)-(1-(R)-(1-(naphthyl)ethoxy)-3-(S)-phenyl-4-(4-(2-oxo-1,3-
imidazolo)methyl-morpholine;
390) 2-(R)-(1-(R)-(1-(4-(fluoro)naphthyl))ethoxy)-3-(S)-phenyl-4-(3-
20 (-1,2,4-triazolo)methyl-morpholine;
391) 2-(R)-(1-(R)-(1-(4-(fluoro)naphthyl))ethoxy)-3-(S)-phenyl-4-(4-
(2-oxo-1,3-imidazolo)methyl-morpholine;
392) 2-(R)-(1-(R)-(1-(3-(fluoro)naphthyl))ethoxy)-3-(S)-phenyl-4-(3-
(-1,2,4-triazolo)methyl-morpholine;
25 393) 2-(R)-(1-(R)-(1-(3-(fluoro)naphthyl))ethoxy)-3-(S)-phenyl-4-(4-
(2-oxo-1,3-imidazolo)methyl-morpholine;
394) 2-(R)-(1-(R)-(1-(3-(chloro)naphthyl))ethoxy)-3-(S)-phenyl-4-(3-
(-1,2,4-triazolo)methyl-morpholine;
395) 2-(R)-(1-(R)-(1-(3-(chloro)naphthyl))ethoxy)-3-(S)-phenyl-4-(4-
30 (2-oxo-1,3-imidazolo)methyl-morpholine;
396) 2-(R)-(1-(R)-(1-(3-(methyl)naphthyl))ethoxy)-3-(S)-phenyl-4-(3-
(-1,2,4-triazolo)methyl-morpholine; and
397) 2-(R)-(1-(R)-(1-(3-(methyl)naphthyl))ethoxy)-3-(S)-phenyl-4-(4-
(2-oxo-1,3-imidazolo)methyl-morpholine.

- 234 -

EXAMPLE 885 2-(R)-(2,5-Bis(trifluoromethyl)benzoyloxy)-3-(S)-(4-fluorophenyl)-4-benzyl-morpholine

The title compound was prepared from 3-(S)-(4-fluorophenyl)-4-benzyl-2-morpholinone (from Example 59) using a procedure analogous to Example 67.

Mass Spectrum (CI): m/Z 528 (M+H)

10 ¹H NMR (CDCl₃, 360 MHz, ppm): δ 2.46 (dt, 1H), 2.90 (dd, 2H), 3.76 (dd, J= 11.6, 2.0, 1H), 3.88 (d, J= 13.6, 1H), 4.18 (t, 1H), 6.20 (d, J=2.8, 1H), 7.04 (d, J= 8.4, 2H), 7.24-7.32 (m, 5H), 7.50, (m, 2H), 7.60 (s, 1H), 7.88 (dd, 2H).

EXAMPLE 8915 2-(R)-(1-(2,5-Bis(trifluoromethyl)phenyl)ethenyloxy)-3-(S)-(4-fluorophenyl)-4-benzyl-morpholine

The title compound was prepared from 2-(R)-(2,5-bis(trifluoromethyl) benzoyloxy)-3-(S)-(4-fluorophenyl)-4-benzyl-morpholine (from Example 88) using a procedure analogous to

20 Example 68.
¹H NMR (CDCl₃, 250 MHz, ppm): δ 2.30 (dt, J= 3.5, 11.9, 1H), 2.74 (app d, J= 9.4, 1H), 2.82 (d, J=13.5, 1H), 3.55-3.60 (m, 2H), 3.72 (d, J= 13.5, 1H), 4.10 (dt, J= 2.4, 11.7, 1H), 4.22 (d, J= 2.7, 1H), 4.67 (d, 25 J=2.8, 1H), 5.18 (d, J=2.8, 1H), 6.90 (t, J=8.7, 2H), 7.08 (s, 1H), 7.13-7.23 (m, 5H), 7.36 (dd, J= 5.6, 8.7, 2H), 7.62 (d, J= 8.4, 1H), 7.72 (d, J= 8.4, 1H).

EXAMPLE 9030 2-(R)-(1-(R)-(2,5-Bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(4-fluorophenyl)-morpholine

The title compound was prepared from 2-(R)-(1-(2,5-bis(trifluoromethyl) phenyl)ethenyloxy)-3-(S)-(4-fluorophenyl)-4-

- 235 -

benzyl-morpholine (from Example 89) using a procedure analogous to Example 74.

Mass Spectrum (CI): m/Z 438 (M+H)

5 ¹H NMR Spectrum (HCl salt, d₆-DMSO, 360 MHz, ppm): δ 1.47 (d, J= 8.7, 3H), 3.88 (d, J= 11.8, 1H), 4.20 (dt, J= 3.7, 11.8, 1H), 4.50 (s, 1H), 4.58 (s, 1H), 5.17 (m, 1H), 7.04 (s, 1H), 7.23 (t, J= 8.8, 2H), 7.55 (m, 2H), 7.77 (d, J= 8.1, 1H), 7.88 (d, J= 8.3, 1H), 10.1 (br s, 1H).

10

EXAMPLE 91

2-(R)-(1-(R)-(2,5-Bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(4-fluorophenyl)-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl-morpholine

15 The title compound was prepared from 2-(R)-(1-(R)-(2,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(4-fluorophenyl)-morpholine (from Example 90) using a procedure analogous to Example 70, mp 162-168 °C.

20 ¹H NMR (d₆-DMSO, 360 MHz, ppm) δ 1.37 (d, J= 6.4, 3H), 2.40 (dt, J= 3.3, 11.9, 1H), 2.77 (d, J= 14.0, 1H), 2.86 (d, J= 11.5, 1H), 3.37 (d, J= 14.4, 1H), 3.48 (d, J= 2.7, 1H), 3.64 (d, J= 11.0, 1H), 4.11 (t, J= 9.8, 1H), 4.18 (d, J= 2.8, 1H), 5.16 (q, J= 6.2, 1H), 6.90 (s, 1H), 7.08 (t, J= 8.8, 2H), 7.50 (br t, 1H), 7.74 (d, J= 8.3, 1H), 7.85 (d, J= 8.3, 1H), 11.25 s, 1H), 11.35 (s, 1H).

25

EXAMPLE 92

2-(R)-(1-(R)-(2,5-Bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(4-fluorophenyl)-4-(3-(1,2,4-triazolo)methyl-morpholine

30 The title compound was prepared from 2-(R)-(1-(R)-(2,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(4-fluorophenyl)-morpholine (from Example 90) using a procedure analogous to Example 17, mp 98-100 °C.

Mass Spectrum (CI): m/Z 519 (M+H)

- 236 -

¹H NMR (d₆-DMSO, 360 MHz, ppm): δ 1.36 (d, J= 6.4, 3H), 2.46 (dt, J= 3.26, 11.9, 1H), 2.89 (d, J= 11.0, 1H), 3.16 (d, J= 13.9, 1H), 3.57-3.64 (m, 3H), 4.09 (t, J= 10.5, 1H), 4.18 (d, J= 2.6, 1H), 5.14 (q, J= 6.4, 1H), 6.90 (s, 1H), 7.11 (t, J= 8.7, 2H), 7.48 (m, 2H), 7.72 (d, J= 8.3, 1H), 7.83 (d, J= 8.3, 1H), 8.36 (br s), 13.8 (s, 1H).

EXAMPLE 93

2-(R)-(1-(R)-(2,5-Bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(4-fluorophenyl)-4-(4-(2-oxo-1,3-imidazolo)methyl-morpholine

The title compound was prepared from 2-(R)-(1-(R)-(2,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(4-fluorophenyl)-morpholine (from Example 90) using a procedure analogous to Example 83. A sample was recrystallized from aqueous ethanol, mp 203-205 °C.

¹H NMR (d₆-DMSO, 360 MHz, ppm): δ 1.35 (d, J= 6.4, 3H), 2.25 (dt, J= 3.1, 11.8, 1H), 2.58 (d, J= 13.9, 1H), 2.88 (d, J= 11.6, 1H), 3.24 (d, J= 14.0, 1H), 3.35 (d, J= 2.7, 1H), 3.64 (dd, J= 9.6, 1H), 4.09 (t, J= 9.8, 1H), 4.16 (d, J= 2.7, 1H), 5.14 (q, J= 6.5, 1H), 5.97 (s, 1H), 6.89 (s, 1H), 7.07 (t, J= 8.7, 1H), 7.49 (m, 1H), 7.72 (d, J= 8.1, 1H), 7.83 (d, J= 8.3, 1H), 9.57 (s, 1H), 9.80 (s, 1H).

EXAMPLE 94

Typical Pharmaceutical Compositions Containing a Compound of the Invention

A: Dry Filled Capsules Containing 5 mg of Active Ingredient Per Capsule

<u>Ingredient</u>	<u>Amount per capsule (mg)</u>
Active ingredient	5
Lactose	194
Magnesium stearate	1
Capsule (size No. 1)	200

- 237 -

5 The active ingredient may be reduced to a No. 60 powder and the lactose and magnesium stearate may then be passed through a No. 60 blotting cloth onto the powder. The combined ingredients may then be mixed for about 10 minutes and filled into a No. 1 dry gelatin capsule.

B: Tablet

10 A typical tablet would contain the active ingredient (25 mg), pregelatinized starch USP (82 mg), microcrystalline cellulose (82 mg) and magnesium stearate (1 mg).

C: Suppository

15 Typical suppository formulations for rectal administration contain the active ingredient (0.08-1.0 mg), disodium calcium edetate (0.25-0.5 mg), and polyethylene glycol (775-1600 mg). Other suppository formulations may be made by substituting, for example, butylated hydroxytoluene (0.04-0.08 mg) for the disodium calcium edetate and a hydrogenated vegetable oil (675-1400 mg) such as Suppocire L, Wecobee FS, Wecobee M, Witepsols, and the like, for the
20 polyethylene glycol.

D: Injection

25 A typical injectible formulation contains the active ingredient, sodium phosphate dibasic anhydrous (11.4 mg), benzyl alcohol (0.01 ml) and water for injection (1.0 ml).

30

- 238 -

While the invention has been described and illustrated with reference to certain particular embodiments thereof, those skilled in the art will appreciate that various adaptations, changes, modifications, substitutions, deletions, or additions of procedures and protocols may be
5 made without departing from the spirit and scope of the invention. For example, effective dosages other than the particular dosages as set forth herein above may be applicable as a consequence of variations in the responsiveness of the mammal being treated for any of the indications with the compounds of the invention indicated above. Likewise, the
10 specific pharmacological responses observed may vary according to and depending upon the particular active compounds selected or whether there are present pharmaceutical carriers, as well as the type of formulation and mode of administration employed, and such expected variations or differences in the results are contemplated in accordance
15 with the objects and practices of the present invention. It is intended, therefore, that the invention be defined by the scope of the claims which follow and that such claims be interpreted as broadly as is reasonable.

20

25

30


- 239 -

WHAT IS CLAIMED IS:

1. A compound of the structural formula:

5

10



or a pharmaceutically acceptable salt thereof, wherein:

15 R1 is selected from the group consisting of:

- (1) hydrogen;
- (2) C1-6 alkyl, unsubstituted or substituted with one or more of the substituents selected from:
 - (a) hydroxy,
 - (b) oxo,
 - (c) C1-6 alkoxy,
 - (d) phenyl-C1-3 alkoxy,
 - (e) phenyl,
 - (f) -CN,
 - (g) halo, wherein halo is fluoro, chloro, bromo or iodo,
 - (h) -NR⁹R¹⁰, wherein R⁹ and R¹⁰ are independently selected from:
 - (i) hydrogen,
 - (ii) C1-6 alkyl,
 - (iii) hydroxy-C1-6 alkyl, and
 - (iv) phenyl,
 - (i) -NR⁹COR¹⁰, wherein R⁹ and R¹⁰ are as defined above,
 - (j) -NR⁹CO₂R¹⁰, wherein R⁹ and R¹⁰ are as defined above,

- 240 -

- (k) -CONR⁹R¹⁰, wherein R⁹ and R¹⁰ are as defined above,
- (l) -COR⁹, wherein R⁹ is as defined above,
- (m) -CO₂R⁹, wherein R⁹ is as defined above;
- 5 (n) heterocycle, wherein the heterocycle is selected from the group consisting of:
- (A) benzimidazolyl,
- (B) benzofuranyl,
- (C) benzothiophenyl,
- 10 (D) benzoxazolyl,
- (E) furanyl,
- (F) imidazolyl,
- (G) indolyl,
- (H) isooxazolyl,
- 15 (I) isothiazolyl,
- (J) oxadiazolyl,
- (K) oxazolyl,
- (L) pyrazinyl,
- (M) pyrazolyl,
- 20 (N) pyridyl,
- (O) pyrimidyl,
- (P) pyrrolyl,
- (Q) quinolyl,
- (R) tetrazolyl,
- 25 (S) thiadiazolyl,
- (T) thiazolyl,
- (U) thienyl,
- (V) triazolyl,
- (W) azetidiny,
- 30 (X) 1,4-dioxanyl,
- (Y) hexahydroazepinyl,
- (Z) piperazinyl,
- (AA) piperidiny,
- (AB) pyrrolidinyl,

- 241 -

(AC) tetrahydrofuranyl, and

(AD) tetrahydrothienyl,

and wherein the heterocycle is unsubstituted or substituted with one or more substituent(s) selected from:

(i) C₁₋₆ alkyl, unsubstituted or substituted with halo, -CF₃, -OCH₃, or phenyl,

(ii) C₁₋₆ alkoxy,

(iii) oxo,

(iv) hydroxy,

(v) thioxo,

(vi) -SR⁹, wherein R⁹ is as defined above,

(vii) halo,

(viii) cyano,

(ix) phenyl,

(x) trifluoromethyl,

(xi) -(CH₂)_m-NR⁹R¹⁰, wherein m is 0, 1 or 2, and R⁹ and R¹⁰ are as defined above,

(xii) -NR⁹COR¹⁰, wherein R⁹ and R¹⁰ are as defined above,

(xiii) -CONR⁹R¹⁰, wherein R⁹ and R¹⁰ are as defined above,

(xiv) -CO₂R⁹, wherein R⁹ is as defined above, and

(xv) -(CH₂)_m-OR⁹, wherein m and R⁹ are as defined above;

(3) C₂₋₆ alkenyl, unsubstituted or substituted with one or more of the substituent(s) selected from:

(a) hydroxy,

(b) oxo,

(c) C₁₋₆ alkoxy,

(d) phenyl-C₁₋₃ alkoxy,

(e) phenyl,

- 242 -

- (f) -CN,
(g) halo,
(h) -CONR⁹R¹⁰ wherein R⁹ and R¹⁰ are as defined above,
5 (i) -COR⁹ wherein R⁹ is as defined above,
(j) -CO₂R⁹, wherein R⁹ is as defined above,
(k) heterocycle, wherein the heterocycle is as defined above;
10 (4) C₂₋₆ alkynyl;
(5) phenyl, unsubstituted or substituted with one or more of the substituent(s) selected from:
(a) hydroxy,
(b) C₁₋₆ alkoxy,
(c) C₁₋₆ alkyl,
15 (d) C₂₋₅ alkenyl,
(e) halo,
(f) -CN,
(g) -NO₂,
(h) -CF₃,
20 (i) -(CH₂)_m-NR⁹R¹⁰, wherein m, R⁹ and R¹⁰ are as defined above,
(j) -NR⁹COR¹⁰, wherein R⁹ and R¹⁰ are as defined above,
(k) -NR⁹CO₂R¹⁰, wherein R⁹ and R¹⁰ are as defined above,
25 (l) -CONR⁹R¹⁰, wherein R⁹ and R¹⁰ are as defined above,
(m) -CO₂NR⁹R¹⁰, wherein R⁹ and R¹⁰ are as defined above,
30 (n) -COR⁹, wherein R⁹ is as defined above;
(o) -CO₂R⁹, wherein R⁹ is as defined above;

- 243 -

R² and R³ are independently selected from the group consisting of:

- (1) hydrogen,
- (2) C₁-6 alkyl, unsubstituted or substituted with one or more
of the substituents selected from:
 - (a) hydroxy,
 - (b) oxo,
 - (c) C₁-6 alkoxy,
 - (d) phenyl-C₁-3 alkoxy,
 - (e) phenyl,
 - (f) -CN,
 - (g) halo,
 - (h) -NR⁹R¹⁰, wherein R⁹ and R¹⁰ are as defined above,
 - (i) -NR⁹COR¹⁰, wherein R⁹ and R¹⁰ are as defined
above,
 - (j) -NR⁹CO₂R¹⁰, wherein R⁹ and R¹⁰ are as defined
above,
 - (k) -CONR⁹R¹⁰, wherein R⁹ and R¹⁰ are as defined
above,
 - (l) -COR⁹, wherein R⁹ is as defined above, and
 - (m) -CO₂R⁹, wherein R⁹ is as defined above;
- (3) C₂-6 alkenyl, unsubstituted or substituted with one or more
of the substituent(s) selected from:
 - (a) hydroxy,
 - (b) oxo,
 - (c) C₁-6 alkoxy,
 - (d) phenyl-C₁-3 alkoxy,
 - (e) phenyl,
 - (f) -CN,
 - (g) halo,
 - (h) -CONR⁹R¹⁰ wherein R⁹ and R¹⁰ are as defined
above,
 - (i) -COR⁹ wherein R⁹ is as defined above,
 - (j) -CO₂R⁹, wherein R⁹ is as defined above;

- 244 -

- (4) C₂-6 alkynyl;
- (5) phenyl, unsubstituted or substituted with one or more of the substituent(s) selected from:
- 5 (a) hydroxy,
(b) C₁-6 alkoxy,
(c) C₁-6 alkyl,
(d) C₂-5 alkenyl,
(e) halo,
10 (f) -CN,
(g) -NO₂,
(h) -CF₃,
(i) -(CH₂)_m-NR⁹R¹⁰, wherein m, R⁹ and R¹⁰ are as defined above,
15 (j) -NR⁹COR¹⁰, wherein R⁹ and R¹⁰ are as defined above,
(k) -NR⁹CO₂R¹⁰, wherein R⁹ and R¹⁰ are as defined above,
(l) -CONR⁹R¹⁰, wherein R⁹ and R¹⁰ are as defined above,
20 (m) -CO₂NR⁹R¹⁰, wherein R⁹ and R¹⁰ are as defined above,
(n) -COR⁹, wherein R⁹ is as defined above;
(o) -CO₂R⁹, wherein R⁹ is as defined above;
- 25 and the groups R¹ and R² may be joined together to form a heterocyclic ring selected from the group consisting of:
- 30 (a) pyrrolidinyl,
(b) piperidinyl,
(c) pyrrolyl,
(d) pyridinyl,
(e) imidazolyl,
(f) oxazolyl, and
(g) thiazolyl,
- and wherein the heterocyclic ring is unsubstituted or

- 245 -

substituted with one or more substituent(s) selected from:

- (i) C₁-6alkyl,
- (ii) oxo,
- (iii) C₁-6alkoxy,
- 5 (iv) -NR⁹R¹⁰, wherein R⁹ and R¹⁰ are as defined above,
- (v) halo, and
- (vi) trifluoromethyl;

10 and the groups R² and R³ may be joined together to form a carbocyclic ring selected from the group consisting of:

- (a) cyclopentyl,
- (b) cyclohexyl,
- (c) phenyl,

15 and wherein the carbocyclic ring is unsubstituted or substituted with one or more substituents selected from:

- (i) C₁-6alkyl,
- (ii) C₁-6alkoxy,
- 20 (iii) -NR⁹R¹⁰, wherein R⁹ and R¹⁰ are as defined above,
- (iv) halo, and
- (v) trifluoromethyl;

25 and the groups R² and R³ may be joined together to form a heterocyclic ring selected from the group consisting of:

- (a) pyrrolidinyl,
- (b) piperidinyl,
- (c) pyrrolyl,
- (d) pyridinyl,
- 30 (e) imidazolyl,
- (f) furanyl,
- (g) oxazolyl,
- (h) thienyl, and
- (i) thiazolyl,

- 246 -

and wherein the heterocyclic ring is unsubstituted or substituted with one or more substituent(s) selected from:

- (i) C₁-6alkyl,
- (ii) oxo,
- (iii) C₁-6alkoxy,
- (iv) -NR⁹R¹⁰, wherein R⁹ and R¹⁰ are as defined above,
- (v) halo, and
- (vi) trifluoromethyl;

R⁶, R⁷ and R⁸ are independently selected from the group consisting of:

- (1) hydrogen;
- (2) C₁-6 alkyl, unsubstituted or substituted with one or more of the substituents selected from:
 - (a) hydroxy,
 - (b) oxo,
 - (c) C₁-6 alkoxy,
 - (d) phenyl-C₁-3 alkoxy,
 - (e) phenyl,
 - (f) -CN,
 - (g) halo,
 - (h) -NR⁹R¹⁰, wherein R⁹ and R¹⁰ are as defined above,
 - (i) -NR⁹COR¹⁰, wherein R⁹ and R¹⁰ are as defined above,
 - (j) -NR⁹CO₂R¹⁰, wherein R⁹ and R¹⁰ are as defined above,
 - (k) -CONR⁹R¹⁰, wherein R⁹ and R¹⁰ are as defined above,
 - (l) -COR⁹, wherein R⁹ is as defined above, and
 - (m) -CO₂R⁹, wherein R⁹ is as defined above;
- (3) C₂-6 alkenyl, unsubstituted or substituted with one or more of the substituent(s) selected from:
 - (a) hydroxy,
 - (b) oxo,

- 247 -

- 5
- (c) C1-6 alkoxy,
 - (d) phenyl-C1-3 alkoxy,
 - (e) phenyl,
 - (f) -CN,
 - (g) halo,
 - (h) -CONR⁹R¹⁰ wherein R⁹ and R¹⁰ are as defined above,
 - (i) -COR⁹ wherein R⁹ is as defined above,
 - (j) -CO₂R⁹, wherein R⁹ is as defined above;
- 10
- (4) C2-6 alkynyl;
 - (5) phenyl, unsubstituted or substituted with one or more of the substituent(s) selected from:
- 15
- (a) hydroxy,
 - (b) C1-6 alkoxy,
 - (c) C1-6 alkyl,
 - (d) C2-5 alkenyl,
 - (e) halo,
 - (f) -CN,
 - (g) -NO₂,
 - (h) -CF₃,
 - (i) -(CH₂)_m-NR⁹R¹⁰, wherein m, R⁹ and R¹⁰ are as defined above,
 - (j) -NR⁹COR¹⁰, wherein R⁹ and R¹⁰ are as defined above,
 - (k) -NR⁹CO₂R¹⁰, wherein R⁹ and R¹⁰ are as defined above,
 - (l) -CONR⁹R¹⁰, wherein R⁹ and R¹⁰ are as defined above,
 - (m) -CO₂NR⁹R¹⁰, wherein R⁹ and R¹⁰ are as defined above,
 - (n) -COR⁹, wherein R⁹ is as defined above;
 - (o) -CO₂R⁹, wherein R⁹ is as defined above;
- 20
- 25
- 30
- (6) halo,
 - (7) -CN,

- 248 -

- (8) -CF₃,
(9) -NO₂,
(10) -SR¹⁴, wherein R¹⁴ is hydrogen or C₁₋₅alkyl,
(11) -SOR¹⁴, wherein R¹⁴ is as defined above,
5 (12) -SO₂R¹⁴, wherein R¹⁴ is as defined above,
(13) NR⁹COR¹⁰, wherein R⁹ and R¹⁰ are as defined above,
(14) CONR⁹COR¹⁰, wherein R⁹ and R¹⁰ are as defined above,
(15) NR⁹R¹⁰, wherein R⁹ and R¹⁰ are as defined above,
(16) NR⁹CO₂R¹⁰, wherein R⁹ and R¹⁰ are as defined above,
10 (17) hydroxy,
(18) C₁₋₆alkoxy,
(19) COR⁹, wherein R⁹ is as defined above,
(20) CO₂R⁹, wherein R⁹ is as defined above,
(21) 2-pyridyl,
15 (22) 3-pyridyl,
(23) 4-pyridyl,
(24) 5-tetrazolyl,
(25) 2-oxazolyl, and
20 (26) 2-thiazolyl;

R¹¹, R¹² and R¹³ are independently selected from the definitions of R⁶, R⁷ and R⁸;

X is selected from the group consisting of:

- 25 (1) -O-,
(2) -S-,
(3) -SO-, and
(4) -SO₂-;

30

- 249 -

Y is selected from the group consisting of:

- 5
- (1) a single bond,
 - (2) -O-,
 - (3) -S-,
 - (4) -CO-,
 - (5) -CH₂-,
 - (6) -CHR¹⁵-, and
 - (7) -CR¹⁵R¹⁶-, wherein R¹⁵ and R¹⁶ are independently
10 selected from the group consisting of:
 - (a) C₁₋₆ alkyl, unsubstituted or substituted with
one or more of the substituents selected from:
 - (i) hydroxy,
 - (ii) oxo,
 - (iii) C₁₋₆ alkoxy,
 - (iv) phenyl-C₁₋₃ alkoxy,
 - (v) phenyl,
 - (vi) -CN,
 - (vii) halo,
 - (viii) -NR⁹R¹⁰, wherein R⁹ and R¹⁰ are as defined
15 above,
 - (ix) -NR⁹COR¹⁰, wherein R⁹ and R¹⁰ are as
defined above,
 - (x) -NR⁹CO₂R¹⁰, wherein R⁹ and R¹⁰ are as
defined above,
 - (xi) -CONR⁹R¹⁰, wherein R⁹ and R¹⁰ are as
defined above,
 - (xii) -COR⁹, wherein R⁹ is as defined above, and
 - (xiii) -CO₂R⁹, wherein R⁹ is as defined above;
 - (b) phenyl, unsubstituted or substituted with one or more
20 of the substituent(s) selected from:
 - (i) hydroxy,
 - (ii) C₁₋₆ alkoxy,
 - (iii) C₁₋₆ alkyl,
- 25
- 30

- 250 -

- 5 (iv) C₂₋₅ alkenyl,
(v) halo,
(vi) -CN,
(vii) -NO₂,
(viii) -CF₃,
(ix) -(CH₂)_m-NR⁹R¹⁰, wherein m, R⁹ and R¹⁰
are as defined above,
10 (x) -NR⁹COR¹⁰, wherein R⁹ and R¹⁰ are as
defined above,
(xi) -NR⁹CO₂R¹⁰, wherein R⁹ and R¹⁰ are as
defined above,
(xii) -CONR⁹R¹⁰, wherein R⁹ and R¹⁰ are as
defined above,
15 (xiii) -CO₂NR⁹R¹⁰, wherein R⁹ and R¹⁰ are as
defined above,
(xiv) -COR⁹, wherein R⁹ is as defined above, and
(xv) -CO₂R⁹, wherein R⁹ is as defined above; and
20 Z is C₁₋₆ alkyl.
- 25
- 30

- 251 -

2. The compound of Claim 1 wherein:

R¹ is C₁-6 alkyl, substituted with one or more of the substituents selected from:

heterocycle, wherein the heterocycle is selected from the group consisting of:

- (A) benzimidazolyl,
- (B) imidazolyl,
- (C) isooxazolyl,
- (D) isothiazolyl,
- (E) oxadiazolyl,
- (F) pyrazinyl,
- (G) pyrazolyl,
- (H) pyridyl,
- (I) pyrrolyl,
- (J) tetrazolyl,
- (K) thiadiazolyl,
- (L) triazolyl, and
- (M) piperidiny,

and wherein the heterocycle is unsubstituted or substituted with one or more substituent(s) selected from:

- (i) C₁-6 alkyl, unsubstituted or substituted with halo, -CF₃, -OCH₃, or phenyl,
- (ii) C₁-6 alkoxy,
- (iii) oxo,
- (iv) thioxo,
- (v) cyano,
- (vi) -SCH₃,
- (vii) phenyl,
- (viii) hydroxy,
- (ix) trifluoromethyl, and
- (x) -(CH₂)_m-NR⁹R¹⁰, wherein m is 0, 1 or 2, and wherein R⁹ and R¹⁰ are independently selected from:

- 252 -

- 5
- (I) hydrogen,
 - (II) C₁₋₆ alkyl,
 - (III) hydroxy-C₁₋₆ alkyl, and
 - (IV) phenyl,
 - (xi) -NR⁹COR¹⁰, wherein R⁹ and R¹⁰ are as defined above, and
 - (xii) -CONR⁹R¹⁰, wherein R⁹ and R¹⁰ are as defined above.

10

3. The compound of Claim 1 wherein:

R² and R³ are independently selected from the group consisting of:

- 15
- (1) hydrogen,
 - (2) C₁₋₆ alkyl,
 - (3) C₂₋₆ alkenyl, and
 - (4) phenyl;

R⁶, R⁷ and R⁸ are independently selected from the group consisting of:

- 20
- (1) hydrogen,
 - (2) C₁₋₆ alkyl,
 - (3) fluoro,
 - (4) chloro,
 - (5) bromo,
 - 25 (6) iodo, and
 - (7) -CF₃;

R¹¹, R¹² and R¹³ are independently selected from the group consisting of:

- 30
- (1) hydrogen,
 - (2) C₁₋₆ alkyl,
 - (3) fluoro,
 - (4) chloro,
 - (5) bromo,



- 253 -

(6) iodo, and



(7) -CF₃;

X is -O-;



5 Y is -O-; and

Z is C₁₋₄ alkyl.10 4. The compound of Claim 1 wherein Z is C₁₋₄ alkyl.5. The compound of Claim 1 wherein Z is -CH₃.15 6. The compound of Claim 1 wherein R¹ is selected from the group consisting of:



20



25



30



- 254 -

5

















10

15


20

25

30



- 255 -




7. The compound of Claim 1 wherein R¹ is selected from the group consisting of:
- 25 (1,2,4-triazolo)methyl; and
(5-oxo-1H,4H-1,2,4-triazolo)methyl.

8. The compound of Claim 1 wherein R¹ is selected from the group consisting of:
- 30 (1,3-imidazolo)methyl; and
(2-oxo-1,3-imidazolo)methyl.


- 256 -

9. The compound of Claim 1 of the structural formula:



or a pharmaceutically acceptable salt thereof, wherein R¹, R², R³, R⁶, R⁷, R⁸, R¹¹, R¹², R¹³, Y and Z are as defined in Claim 1.

10. The compound of Claim 1 of the structural
15 formula II:




or a pharmaceutically acceptable salt thereof, wherein R¹, R², R³, R⁶, R⁷, R⁸, R¹¹, R¹², R¹³ and Z are as defined in Claim 1.

30


- 257 -

11. The compound of Claim 1 of the structural formula III:



or a pharmaceutically acceptable salt thereof, wherein R¹, R², R³, R⁶, R⁷, R⁸, R¹¹, R¹², R¹³ and Z are as defined in Claim 1.


12. The compound of Claim 1 of the structural formula:



or a pharmaceutically acceptable salt thereof, wherein R¹, R², R³, R⁶, R⁷, R⁸, R¹¹, R¹², R¹³, Y and Z are as defined in Claim 1.


- 258 -

13. The compound of Claim 1 of the structural formula:



or a pharmaceutically acceptable salt thereof, wherein R¹, R², R³, R⁶, R⁷, R⁸, R¹¹, R¹², R¹³, Y and Z are as defined in Claim 1.

14. The compound of Claim 1 of the structural formula:



25 or a pharmaceutically acceptable salt thereof, wherein R¹, R², R³, R⁶, R⁷, R⁸, R¹¹, R¹², R¹³, Y and Z are as defined in Claim 1.

30

- 259 -

15. A compound which is selected from the group consisting of:

- 5 87) 2-(R)-(1-(S)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-phenyl-morpholine;
- 88) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-phenyl-morpholine;
- 10 89) 2-(R)-(1-(S)-(3,5-Bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl-morpholine;
- 90) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl-morpholine;
- 15 93) 2-(R)-(1-(S)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-morpholine;
- 20 94) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-morpholine;
- 95) 2-(R)-(1-(S)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl-morpholine;
- 25 96) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl-morpholine;
- 30 98) 2-(R)-(1-(R)-(1-(3-(methyl)naphthyl))ethoxy)-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl-morpholine;
- 99) 2-(R)-(1-(R)-(3-(fluoro)-5-(trifluoromethyl)phenyl)ethoxy)-3-(S)-phenyl-morpholine;

- 260 -

- 100) 2-(R)-(1-(R)-(3-(fluoro)-5-(trifluoromethyl)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
5
- 101) 2-(R)-(1-(R)-(3-(chloro)-5-(trifluoromethyl)phenyl)ethoxy)-3-(S)-phenyl-morpholine;
- 102) 2-(R)-(1-(R)-(3-(chloro)-5-(trifluoromethyl)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
10
- 103) 2-(R)-(1-(R)-(3,5-(dimethyl)phenyl)ethoxy)-3-(S)-phenyl-morpholine;
15
- 104) 2-(R)-(1-(R)-(3,5-(dimethyl)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 105) 2-(R)-(1-(R)-(3-(fluoro)-5-(methyl)phenyl)ethoxy)-3-(S)-phenyl-morpholine;
20
- 106) 2-(R)-(1-(R)-(3-(fluoro)-5-(methyl)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 107) 2-(R)-(1-(R)-(3-(chloro)-5-(methyl)phenyl)ethoxy)-3-(S)-phenyl-morpholine;
25
- 108) 2-(R)-(1-(R)-(3-(chloro)-5-(methyl)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
30
- 109) 2-(R)-(1-(R)-(3-(bromo)-5-(methyl)phenyl)ethoxy)-3-(S)-phenyl-morpholine;

- 261 -

- 110) 2-(R)-(1-(R)-(3-(bromo)-5-(methyl)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 5 111) 2-(R)-(1-(R)-(3-(isopropoxy)phenyl)ethoxy)-3-(S)-phenyl-morpholine;
- 112) 2-(R)-(1-(R)-(3-(isopropoxy)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 10 113) 2-(R)-(1-(R)-(3-(isopropoxy)-5-trifluoromethyl)phenyl)ethoxy)-3-(S)-phenyl-morpholine;
- 114) 2-(R)-(1-(R)-(3-(isopropoxy)-5-trifluoromethyl)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 15 115) 2-(R)-(1-(R)-(3-(chloro)-5-(isopropoxy)phenyl)ethoxy)-3-(S)-phenyl-morpholine;
- 20 116) 2-(R)-(1-(R)-(3-(chloro)-5-(isopropoxy)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 117) 2-(R)-(1-(R)-(3-(fluoro)-5-(isopropoxy)phenyl)ethoxy)-3-(S)-phenyl-morpholine;
- 25 118) 2-(R)-(1-(R)-(3-(fluoro)-5-(isopropoxy)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 30 121) 2-(R)-(1-(R)-(3-(t-butyl)-5-(chloro)phenyl)ethoxy)-3-(S)-phenyl-morpholine;
- 122) 2-(R)-(1-(R)-(3-(t-butyl)-5-(chloro)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;

- 262 -

- 123) 2-(R)-(1-(R)-(3-(t-butyl)-5-(trifluoromethyl)phenyl)ethoxy)-3-(S)-phenyl-morpholine;
- 5 124) 2-(R)-(1-(R)-(3-(t-butyl)-5-(trifluoromethyl)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 10 125) 2-(R)-(1-(R)-(3,5-(dimethyl)-4-(fluoro)phenyl)ethoxy)-3-(S)-phenyl-morpholine;
- 126) 2-(R)-(1-(R)-(3,5-(dimethyl)-4-(fluoro)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 15 127) 2-(R)-(1-(R)-(3,5-(dimethyl)-4-(chloro)phenyl)ethoxy)-3-(S)-phenyl-morpholine;
- 128) 2-(R)-(1-(R)-(3,5-(dimethyl)-4-(chloro)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 20 129) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)-4-(fluoro)phenyl)ethoxy)-3-(S)-phenyl-morpholine;
- 25 130) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)-4-(fluoro)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 131) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)-4-chloro)phenyl)ethoxy)-3-(S)-phenyl-morpholine;
- 30 132) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)-4-chloro)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;

- 263 -

- 133) 2-(R)-(1-(R)-(3,5-(dichloro)phenyl)ethoxy)-3-(S)-phenyl-morpholine;
- 5 134) 2-(R)-(1-(R)-(3,5-(dichloro)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 135) 2-(R)-(1-(R)-(3,5-(difluoro)phenyl)ethoxy)-3-(S)-phenyl-morpholine;
- 10 136) 2-(R)-(1-(R)-(3,5-(difluoro)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 15 153) 2-(R)-(1-(R)-(2-fluoro-5-trifluoromethyl)phenylethoxy)-3-(S)-phenyl-morpholine;
- 154) 2-(R)-(1-(R)-(2-fluoro-5-trifluoromethyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-morpholine;
- 20 157) 2-(R)-(1-(R)-(2-fluoro-5-trifluoromethyl)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 158) 2-(R)-(1-(R)-(2-fluoro-5-trifluoromethyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 25 161) 2-(R)-(1-(R)-(2-fluoro-5-trifluoromethyl)phenylethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 162) 2-(R)-(1-(R)-(2-fluoro-5-trifluoromethyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 30 165) 2-(R)-(1-(R)-(2-fluoro-5-trifluoromethyl)phenylethoxy)-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;

- 264 -

- 166) 2-(R)-(1-(R)-(2-fluoro-5-trifluoromethyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 5 169) 2-(R)-(1-(R)-(2-fluoro-5-trifluoromethyl)phenylethoxy)-3-(S)-phenyl-4-(2-imidazolo)methyl-morpholine;
- 170) 2-(R)-(1-(R)-(2-fluoro-5-trifluoromethyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(2-imidazolo)methyl-morpholine;
- 10 173) 2-(R)-(1-(R)-(2-fluoro-5-trifluoromethyl)phenylethoxy)-3-(S)-phenyl-4-(4-imidazolo)methyl-morpholine;
- 174) 2-(R)-(1-(R)-(2-fluoro-5-trifluoromethyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-imidazolo)methyl-morpholine;
- 15 177) 2-(R)-(1-(R)-(2-fluoro-5-trifluoromethyl)phenylethoxy)-3-(S)-phenyl-4-(5-tetrazolo)methyl-morpholine;
- 20 178) 2-(R)-(1-(R)-(2-fluoro-5-trifluoromethyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(5-tetrazolo)methyl-morpholine;
- 181) 2-(R)-(1-(R)-(2-fluoro-5-trifluoromethyl)phenylethoxy)-3-(S)-phenyl-4-(2-oxo-5H-pyrrol-4-yl)methyl-morpholine;
- 25 182) 2-(R)-(1-(R)-(2-fluoro-5-trifluoromethyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(2-oxo-5H-pyrrol-4-yl)methyl-morpholine;
- 185) 2-(R)-(1-(R)-(2-chloro-5-trifluoromethyl)phenylethoxy)-3-(S)-phenyl-morpholine;
- 30 186) 2-(R)-(1-(R)-(2-chloro-5-trifluoromethyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-morpholine;

- 265 -

- 189) 2-(R)-(1-(R)-(2-chloro-5-trifluoromethyl)phenylethoxy)-3-(S)-phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 5 190) 2-(R)-(1-(R)-(2-chloro-5-trifluoromethyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 193) 2-(R)-(1-(R)-(2-chloro-5-trifluoromethyl)phenylethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 10 194) 2-(R)-(1-(R)-(2-chloro-5-trifluoromethyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 15 197) 2-(R)-(1-(R)-(2-chloro-5-trifluoromethyl)phenylethoxy)-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 198) 2-(R)-(1-(R)-(2-chloro-5-trifluoromethyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 20 201) 2-(R)-(1-(R)-(2-chloro-5-trifluoromethyl)phenylethoxy)-3-(S)-phenyl-4-(2-imidazolo)methyl)-morpholine;
- 202) 2-(R)-(1-(R)-(2-chloro-5-trifluoromethyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(2-imidazolo)methyl)-morpholine;
- 25 205) 2-(R)-(1-(R)-(2-chloro-5-trifluoromethyl)phenylethoxy)-3-(S)-phenyl-4-(4-imidazolo)methyl-morpholine;
- 206) 2-(R)-(1-(R)-(2-chloro-5-trifluoromethyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-imidazolo)methyl-morpholine;
- 30 209) 2-(R)-(1-(R)-(2-chloro-5-trifluoromethyl)phenylethoxy)-3-(S)-phenyl-4-(5-tetrazolo)methyl-morpholine;

- 266 -

- 210) 2-(R)-(1-(R)-(2-chloro-5-trifluoromethyl)phenylethoxy)-3-(S)-
(4-fluoro)phenyl-4-(5-tetrazolo)methyl-morpholine;
- 5 213) 2-(R)-(1-(R)-(2-chloro-5-trifluoromethyl)phenylethoxy)-3-(S)-
phenyl-4-(2-oxo-5H-pyrrol-4-yl)methyl-morpholine;
- 214) 2-(R)-(1-(R)-(2-Chloro-5-trifluoromethyl)phenylethoxy)-3-(S)-
(4-fluoro)phenyl-4-(2-oxo-5H-pyrrol-4-yl)methyl-morpholine;
- 10 217) 2-(R)-(1-(R)-(3-methyl)phenylethoxy)-3-(S)-phenyl-morpholine;
- 218) 2-(R)-(1-(R)-(3-methyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-
morpholine;
- 15 221) 2-(R)-(1-(R)-(3-methyl)phenylethoxy)-3-(S)-phenyl-4-(3-(1,2,4-
triazolo)methyl)-morpholine;
- 222) 2-(R)-(1-(R)-(3-methyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-
20 (3-(1,2,4-triazolo)methyl)-morpholine;
- 225) 2-(R)-(1-(R)-(3-methyl)phenylethoxy)-3-(S)-phenyl-4-(3-(5-oxo-
1,2,4-triazolo)methyl)-morpholine;
- 226) 2-(R)-(1-(R)-(3-methyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-
25 (3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 229) 2-(R)-(1-(R)-(3-methyl)phenylethoxy)-3-(S)-phenyl-4-(4-(2-oxo-
1,3-imidazolo)methyl)-morpholine;
- 30 230) 2-(R)-(1-(R)-(3-methyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-
(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 233) 2-(R)-(1-(R)-(3-methyl)phenylethoxy)-3-(S)-phenyl-4-(2-
imidazolo)methyl-morpholine;

- 267 -

- 234) 2-(R)-(1-(R)-(3-methyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(2-imidazolo)methyl-morpholine;
- 5 237) 2-(R)-(1-(R)-(3-methyl)phenylethoxy)-3-(S)-phenyl-4-(4-imidazolo)methyl-morpholine;
- 238) 2-(R)-(1-(R)-(3-methyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-imidazolo)methyl-morpholine;
- 10 241) 2-(R)-(1-(R)-(3-methyl)phenylethoxy)-3-(S)-phenyl-4-(5-tetrazolo)methyl-morpholine;
- 15 242) 2-(R)-(1-(R)-(3-methyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(5-tetrazolo)methyl-morpholine;
- 245) 2-(R)-(1-(R)-(3-methyl)phenylethoxy)-3-(S)-phenyl-4-(2-oxo-5H-pyrrol-4-yl)methyl-morpholine;
- 20 246) 2-(R)-(1-(R)-(3-methyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(2-oxo-5H-pyrrol-4-yl)methyl-morpholine;
- 249) 2-(R)-(1-(R)-(3-bromo)phenylethoxy)-3-(S)-phenyl-morpholine;
- 25 250) 2-(R)-(1-(R)-(3-bromo)phenylethoxy)-3-(S)-(4-fluoro)phenyl-morpholine;
- 253) 2-(R)-(1-(R)-(3-bromo)phenylethoxy)-3-(S)-phenyl-4-(3-(1,2,4-triazolo)methyl-morpholine;
- 30 254) 2-(R)-(1-(R)-(3-bromo)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl-morpholine;

- 268 -

- 257) 2-(R)-(1-(R)-(3-bromo)phenylethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 5 258) 2-(R)-(1-(R)-(3-bromo)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 261) 2-(R)-(1-(R)-(3-bromo)phenylethoxy)-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 10 262) 2-(R)-(1-(R)-(3-bromo)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 265) 2-(R)-(1-(R)-(3-bromo)phenylethoxy)-3-(S)-phenyl-4-(2-imidazolo)methyl-morpholine;
- 15 266) 2-(R)-(1-(R)-(3-bromo)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(2-imidazolo)methyl-morpholine;
- 20 269) 2-(R)-(1-(R)-(3-bromo)phenylethoxy)-3-(S)-phenyl-4-(4-imidazolo)methyl-morpholine;
- 270) 2-(R)-(1-(R)-(3-bromo)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-imidazolo)methyl-morpholine;
- 25 273) 2-(R)-(1-(R)-(3-bromo)phenylethoxy)-3-(S)-phenyl-4-(5-tetrazolo)methyl-morpholine;
- 274) 2-(R)-(1-(R)-(3-bromo)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(5-tetrazolo)methyl-morpholine;
- 30 277) 2-(R)-(1-(R)-(3-bromo)phenylethoxy)-3-(S)-phenyl-4-(2-oxo-5H-pyrrol-4-yl)methyl-morpholine;

- 269 -

- 278) 2-(R)-(1-(R)-(3-bromo)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(2-oxo-5H-pyrrol-4-yl)methyl-morpholine;
- 5 281) 2-(R)-(1-(R)-(3-chloro)phenylethoxy)-3-(S)-phenyl-morpholine;
- 282) 2-(R)-(1-(R)-(3-chloro)phenylethoxy)-3-(S)-(4-fluoro)phenyl-morpholine;
- 10 285) 2-(R)-(1-(R)-(3-chloro)phenylethoxy)-3-(S)-phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 286) 2-(R)-(1-(R)-(3-chloro)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 15 289) 2-(R)-(1-(R)-(3-chloro)phenylethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 290) 2-(R)-(1-(R)-(3-chloro)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 20 293) 2-(R)-(1-(R)-(3-chloro)phenylethoxy)-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 294) 2-(R)-(1-(R)-(3-chloro)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 25 297) 2-(R)-(1-(R)-(3-chloro)phenylethoxy)-3-(S)-phenyl-4-(2-imidazolo)methyl-morpholine;
- 30 298) 2-(R)-(1-(R)-(3-chloro)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(2-imidazolo)methyl-morpholine;
- 301) 2-(R)-(1-(R)-(3-chloro)phenylethoxy)-3-(S)-phenyl-4-(4-imidazolo)methyl-morpholine;

- 270 -

- 302) 2-(R)-(1-(R)-(3-chloro)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-imidazolo)methyl-morpholine;
- 5 305) 2-(R)-(1-(R)-(3-chloro)phenylethoxy)-3-(S)-phenyl-4-(5-tetrazolo)methyl-morpholine;
- 306) 2-(R)-(1-(R)-(3-chloro)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(5-tetrazolo)methyl-morpholine;
- 10 309) 2-(R)-(1-(R)-(3-chloro)phenylethoxy)-3-(S)-phenyl-4-(2-oxo-5H-pyrrol-4-yl)methyl-morpholine;
- 310) 2-(R)-(1-(R)-(3-chloro)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(2-oxo-5H-pyrrol-4-yl)methyl-morpholine;
- 15 313) 2-(R)-(1-(R)-(3-trifluoromethyl)phenylethoxy)-3-(S)-phenyl-morpholine;
- 20 314) 2-(R)-(1-(R)-(3-trifluoromethyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-morpholine;
- 317) 2-(R)-(1-(R)-(3-trifluoromethyl)phenylethoxy)-3-(S)-phenyl-4-(3-1H,4H-1,2,4-triazolo)methyl-morpholine;
- 25 318) 2-(R)-(1-(R)-(3-trifluoromethyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl-morpholine;
- 30 321) 2-(R)-(1-(R)-(3-trifluoromethyl)phenylethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 322) 2-(R)-(1-(R)-(3-trifluoromethyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;

- 271 -

- 325) 2-(R)-(1-(R)-(3-trifluoromethyl)phenylethoxy)-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 5 326) 2-(R)-(1-(R)-(3-trifluoromethyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 329) 2-(R)-(1-(R)-(3-trifluoromethyl)phenylethoxy)-3-(S)-phenyl-4-(2-imidazolo)methyl-morpholine;
- 10 330) 2-(R)-(1-(R)-(3-trifluoromethyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(2-imidazolo)methyl-morpholine;
- 333) 2-(R)-(1-(R)-(3-trifluoromethyl)phenylethoxy)-3-(S)-phenyl-4-(4-imidazolo)methyl-morpholine;
- 15 334) 2-(R)-(1-(R)-(3-trifluoromethyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-imidazolo)methyl-morpholine;
- 20 337) 2-(R)-(1-(R)-(3-trifluoromethyl)phenylethoxy)-3-(S)-phenyl-4-(5-tetrazolo)methyl-morpholine;
- 338) 2-(R)-(1-(R)-(3-trifluoromethyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(5-tetrazolo)methyl-morpholine;
- 25 341) 2-(R)-(1-(R)-(3-trifluoromethyl)phenylethoxy)-3-(S)-phenyl-4-(2-oxo-5H-pyrrol-4-yl)methyl-morpholine;
- 342) 2-(R)-(1-(R)-(3-trifluoromethyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(2-oxo-5H-pyrrol-4-yl)methyl-morpholine;
- 30 345) 2-(R)-(1-(R)-(3-t-butyl)phenylethoxy)-3-(S)-phenyl-morpholine;

- 272 -

- 346) 2-(R)-(1-(R)-(3-t-butyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-morpholine;
- 5 349) 2-(R)-(1-(R)-(3-t-butyl)phenylethoxy)-3-(S)-phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 350) 2-(R)-(1-(R)-(3-t-butyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 10 353) 2-(R)-(1-(R)-(3-t-butyl)phenylethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 354) 2-(R)-(1-(R)-(3-t-butyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 15 357) 2-(R)-(1-(R)-(3-t-butyl)phenylethoxy)-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 358) 2-(R)-(1-(R)-(3-t-butyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 20 361) 2-(R)-(1-(R)-(3-t-butyl)phenylethoxy)-3-(S)-phenyl-4-(2-imidazolo)methyl-morpholine;
- 25 362) 2-(R)-(1-(R)-(3-t-butyl)phenylethoxy)-3-(S)-(4-fluoro)phenyl-4-(2-imidazolo)methyl-morpholine;
- 365) 2-(R)-(1-(R)-(3-t-butyl)phenylethoxy)-3-(S)-phenyl-4-(4-imidazolo)methyl-morpholine;
- 30 366) 2-(R)-(1-(R)-(3-(t-butyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-imidazolo)methyl-morpholine;

- 273 -

- 369) 2-(R)-(1-(R)-(3-(t-butyl)phenyl)ethoxy)-3-(S)-phenyl-4-(5-tetrazolo)methyl-morpholine;
- 5 370) 2-(R)-(1-(R)-(3-(t-butyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(5-tetrazolo)methyl-morpholine;
- 373) 2-(R)-(1-(R)-(3-(t-butyl)phenyl)ethoxy)-3-(S)-phenyl-4-(2-oxo-5H-pyrrol-4-yl)methyl-morpholine;
- 10 374) 2-(R)-(1-(R)-(3-(t-butyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(2-oxo-5H-pyrrol-4-yl)methyl-morpholine;
- 378) 2-(R)-(1-(R)-(2,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(4-fluorophenyl)-morpholine;
- 15 379) 2-(R)-(1-(R)-(2,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(4-fluorophenyl)-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 20 380) 2-(R)-(1-(R)-(2,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(4-fluorophenyl)-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 381) 2-(R)-(1-(R)-(2,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(4-fluorophenyl)-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 25 382) 2-(R)-(1-(R)-(3-(thiomethyl)phenyl)ethoxy)-3-(S)-phenyl-morpholine;
- 383) 2-(R)-(1-(R)-(3-(thiomethyl)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 30 384) 2-(R)-(1-(R)-(3-(thiomethyl)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;

- 274 -

- 385) 2-(R)-(1-(R)-(3-(thiomethylphenyl)ethoxy)-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 5 386) 2-(R)-(1-(R)-(3-(thiomethyl)-5-(trifluoromethyl)phenyl)ethoxy)-3-(S)-phenyl-morpholine;
- 387) 2-(R)-(1-(R)-(3-(thiomethyl-5-(trifluoromethyl)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 10 388) 2-(R)-(1-(R)-(3-(thiomethyl-5-(trifluoromethyl)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 15 389) 2-(R)-(1-(R)-(3-(thiomethyl-5-(trifluoromethyl)phenyl)ethoxy)-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 394) 2-(R)-(1-(R)-(3,5-(dimethoxy)phenyl)ethoxy)-3-(S)-phenyl-morpholine;
- 20 395) 2-(R)-(1-(R)-(3,5-(dimethoxy)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 396) 2-(R)-(1-(R)-(3,5-(dimethoxy)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 25 397) 2-(R)-(1-(R)-(3,5-(dimethoxy)phenyl)ethoxy)-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 30 398) 2-(R)-(1-(R)-(3-(fluoro)-5-(trifluoromethyl)phenyl)-ethoxy)-3-(S)-(4-fluorophenyl)-morpholine;
- 399) 2-(R)-(1-(R)-(3-(fluorophenyl)-5-(trifluoromethyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;

- 275 -

- 400) 2-(R)-(1-(R)-(3-(fluorophenyl)-5-(trifluoromethyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 5 401) 2-(R)-(1-(R)-(3-(fluorophenyl)-5-(trifluoromethyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 402) 2-(R)-(1-(R)-(3-(chloro)-5-(trifluoromethyl)phenyl)-ethoxy)-3-(S)-(4-fluoro)phenyl-morpholine;
- 10 403) 2-(R)-(1-(R)-(3-(chlorophenyl)-5-(trifluoromethyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 15 404) 2-(R)-(1-(R)-(3-(chlorophenyl)-5-(trifluoromethyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 405) 2-(R)-(1-(R)-(3-(chlorophenyl)-5-(trifluoromethyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 20 406) 2-(R)-(1-(R)-(3,5-(dimethyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-morpholine;
- 407) 2-(R)-(1-(R)-(3,5-(dimethyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 25 408) 2-(R)-(1-(R)-(3,5-(dimethyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 30 409) 2-(R)-(1-(R)-(3,5-(dimethyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;

- 276 -

- 410) 2-(R)-(1-(R)-(3-(fluoro)-5-(methyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-morpholine;
- 5 411) 2-(R)-(1-(R)-(3-(fluoro)-5-(methyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 10 412) 2-(R)-(1-(R)-(3-(fluoro)-5-(methyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 413) 2-(R)-(1-(R)-(3-(fluoro)-5-(methyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 15 414) 2-(R)-(1-(R)-(3-(chloro)-5-(methyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-morpholine;
- 415) 2-(R)-(1-(R)-(3-(chloro)-5-(methyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 20 416) 2-(R)-(1-(R)-(3-(chloro)-5-(methyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 25 417) 2-(R)-(1-(R)-(3-(chloro)-5-(methyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 418) 2-(R)-(1-(R)-(3-(bromo)-5-(methyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-morpholine;
- 30 419) 2-(R)-(1-(R)-(3-(bromo)-5-(methyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;

- 277 -

- 420) 2-(R)-(1-(R)-(3-(bromo)-5-(methyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 5 421) 2-(R)-(1-(R)-(3-(bromo)-5-(methyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 422) 2-(R)-(1-(R)-(3-(isopropoxy)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-morpholine;
- 10 423) 2-(R)-(1-(R)-(3-(isopropoxy)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 15 424) 2-(R)-(1-(R)-(3-(isopropoxy)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 425) 2-(R)-(1-(R)-(3-(isopropoxy)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 20 426) 2-(R)-(1-(R)-(3-(isopropoxy)-5-(trifluoromethyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-morpholine;
- 427) 2-(R)-(1-(R)-(3-(isopropoxy)-5-(trifluoromethyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 25 428) 2-(R)-(1-(R)-(3-(isopropoxy)-5-(trifluoromethyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 30 429) 2-(R)-(1-(R)-(3-(isopropoxy)-5-(trifluoromethyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;

- 278 -

- 430) 2-(R)-(1-(R)-(3-(chloro)-5-(isopropoxy)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-morpholine;
- 5 431) 2-(R)-(1-(R)-(3-(chloro)-5-(isopropoxy)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 10 432) 2-(R)-(1-(R)-(3-(chloro)-5-(isopropoxy)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 433) 2-(R)-(1-(R)-(3-(chloro)-5-(isopropoxy)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 15 434) 2-(R)-(1-(R)-(3-(fluoro)-5-(isopropoxy)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-morpholine;
- 435) 2-(R)-(1-(R)-(3-(fluoro)-5-(isopropoxy)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 20 436) 2-(R)-(1-(R)-(3-(fluoro)-5-(isopropoxy)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 25 437) 2-(R)-(1-(R)-(3-(fluoro)-5-(isopropoxy)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 438) 2-(R)-(1-(R)-(3-(t-butyl)-5-(chloro)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-morpholine;
- 30 439) 2-(R)-(1-(R)-(3-(t-butyl)-5-(chloro)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;

- 279 -

- 440) 2-(R)-(1-(R)-(3-(t-butyl)-5-(chloro)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 5 441) 2-(R)-(1-(R)-(3-(t-butyl)-5-(chloro)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 442) 2-(R)-(1-(R)-(3-(t-butyl)-5-(trifluoromethyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-morpholine;
- 10 443) 2-(R)-(1-(R)-(3-(t-butyl)-5-(trifluoromethyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 15 444) 2-(R)-(1-(R)-(3-(t-butyl)-5-(trifluoromethyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 445) 2-(R)-(1-(R)-(3-(t-butyl)-5-(trifluoromethyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 20 446) 2-(R)-(1-(R)-(3,5-(dimethyl)-4-(fluoro)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-morpholine;
- 25 447) 2-(R)-(1-(R)-(3,5-(dimethyl)-4-(fluoro)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 448) 2-(R)-(1-(R)-(3,5-(dimethyl)-4-(fluoro)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 30 449) 2-(R)-(1-(R)-(3,5-(dimethyl)-4-(fluoro)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;

- 280 -

- 450) 2-(R)-(1-(R)-(3,5-(dimethyl)-4-(chloro)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-morpholine;
- 5 451) 2-(R)-(1-(R)-(3,5-(dimethyl)-4-(chloro)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 10 452) 2-(R)-(1-(R)-(3,5-(dimethyl)-4-(chloro)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 453) 2-(R)-(1-(R)-(3,5-(dimethyl)-4-(chloro)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 15 454) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)-4-(fluoro)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-morpholine;
- 455) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)-4-(fluoro)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 20 456) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)-4-(fluoro)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 25 457) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)-4-(fluoro)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 30 458) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)-4-(chloro)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-morpholine;
- 459) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)-4-(chloro)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;

- 281 -

- 460) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)-4-(chloro)phenyl)ethoxy)-
3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 5 461) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)-4-(chloro)phenyl)ethoxy)-
3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-
morpholine;
- 462) 2-(R)-(1-(R)-(3,5-(dichloro)phenyl)ethoxy)-3-(S)-(4-
10 fluoro)phenyl-morpholine;
- 463) 2-(R)-(1-(R)-(3,5-(dichloro)phenyl)ethoxy)-3-(S)-(4-
fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-
morpholine;
- 15 464) 2-(R)-(1-(R)-(3,5-(dichloro)phenyl)ethoxy)-3-(S)-(4-
fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 465) 2-(R)-(1-(R)-(3,5-(dichloro)phenyl)ethoxy)-3-(S)-(4-
20 fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 466) 2-(R)-(1-(R)-(3,5-(difluoro)phenyl)ethoxy)-3-(S)-(4-
fluoro)phenyl-morpholine;
- 25 467) 2-(R)-(1-(R)-(3,5-(difluoro)phenyl)ethoxy)-3-(S)-(4-
fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-
morpholine;
- 468) 2-(R)-(1-(R)-(3,5-(difluoro)phenyl)ethoxy)-3-(S)-(4-
30 fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 469) 2-(R)-(1-(R)-(3,5-(difluoro)phenyl)ethoxy)-3-(S)-(4-
fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;

- 282 -

- 494) 2-(R)-(1-(R)-(3-(thiomethyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-morpholine;
- 5 495) 2-(R)-(1-(R)-(3-(thiomethyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 10 496) 2-(R)-(1-(R)-(3-(thiomethyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 497) 2-(R)-(1-(R)-(3-(thiomethyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 15 498) 2-(R)-(1-(R)-(3-(thiomethyl)-5-(trifluoromethyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-morpholine;
- 499) 2-(R)-(1-(R)-(3-(thiomethyl-5-(trifluoromethyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 20 500) 2-(R)-(1-(R)-(3-(thiomethyl-5-(trifluoromethyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 25 501) 2-(R)-(1-(R)-(3-(thiomethyl-5-(trifluoromethyl)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 30 506) 2-(R)-(1-(R)-(3,5-(dimethoxy)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-morpholine;
- 507) 2-(R)-(1-(R)-(3,5-(dimethoxy)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;

- 283 -

- 508) 2-(R)-(1-(R)-(3,5-(dimethoxy)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 5 509) 2-(R)-(1-(R)-(3,5-(dimethoxy)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 510) 2-(R)-(1-(R)-(phenyl)ethoxy)-3-(S)-phenyl-morpholine;
- 10 511) 2-(R)-(1-(R)-(phenyl)ethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 512) 2-(R)-(1-(R)-(phenyl)ethoxy)-3-(S)-phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 15 513) 2-(R)-(1-(R)-(phenyl)ethoxy)-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 514) 2-(R)-(1-(R)-(phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl morpholine;
- 20 515) 2-(R)-(1-(R)-(phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 516) 2-(R)-(1-(R)-(phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 25 517) 2-(R)-(1-(R)-(phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 518) 2-(R)-(1-(R)-(3-(fluoro)phenyl)ethoxy)-3-(S)-phenyl-morpholine;
- 30 519) 2-(R)-(1-(R)-(3-(fluoro)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;

- 284 -

- 520) 2-(R)-(1-(R)-(3-(fluoro)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 5 521) 2-(R)-(1-(R)-(3-(fluoro)phenyl)ethoxy)-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 522) 2-(R)-(1-(R)-(3-(fluoro)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-morpholine;
- 10 523) 2-(R)-(1-(R)-(3-(fluoro)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 524) 2-(R)-(1-(R)-(3-(fluoro)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 15 525) 2-(R)-(1-(R)-(3-(fluoro)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 526) 2-(R)-(1-(R)-(4-(fluoro)phenyl)ethoxy)-3-(S)-phenyl-morpholine;
- 20 527) 2-(R)-(1-(R)-(4-(fluoro)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 528) 2-(R)-(1-(R)-(4-(fluoro)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 25 529) 2-(R)-(1-(R)-(4-(fluoro)phenyl)ethoxy)-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 530) 2-(R)-(1-(R)-(4-(fluoro)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-morpholine;
- 30 531) 2-(R)-(1-(R)-(4-(fluoro)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;

- 285 -

- 532) 2-(R)-(1-(R)-(4-(fluoro)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 5 533) 2-(R)-(1-(R)-(4-(fluoro)phenyl)ethoxy)-3-(S)-(4-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 534) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(3-fluoro)phenyl-morpholine;
- 10 535) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(3-fluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 15 536) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(3-fluoro)phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 537) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(3-fluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 20 538) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(3,4-difluoro)phenyl-morpholine;
- 539) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(3,4-difluoro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 25 540) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(3,4-difluoro)phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 30 541) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(3,4-difluoro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;

- 286 -

- 542) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(3,4-dichloro)phenyl-morpholine;
- 5 543) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(3,4-dichloro)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 10 544) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(3,4-dichloro)phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 545) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(3,4-dichloro)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 15 546) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(3,4-dimethyl)phenyl-morpholine;
- 547) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(3,4-dimethyl)phenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 20 548) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(3,4-dimethyl)phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 25 549) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(3,4-dimethyl)phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 550) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-3,4-methylenedioxyphenyl-morpholine;
- 30 551) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-3,4-methylenedioxyphenyl-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;

- 287 -

- 552) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-3,4-methylenedioxyphenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 5 553) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-3,4-methylenedioxyphenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 10 554) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(2-naphthyl)-morpholine;
- 555) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(2-naphthyl)-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 15 556) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(2-naphthyl)-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 557) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(2-naphthyl)-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 20 558) 2-(R)-(1-(R)-(3-(fluorophenyl)-5-(trifluoromethyl)ethoxy)-3-(S)-phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 559) 2-(R)-(1-(R)-(3-(fluorophenyl)-5-(trifluoromethyl)ethoxy)-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 25 560) 2-(R)-(1-(R)-(3-(chlorophenyl)-5-(trifluoromethyl)ethoxy)-3-(S)-phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 561) 2-(R)-(1-(R)-(3-(chlorophenyl)-5-(trifluoromethyl)ethoxy)-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 30 562) 2-(R)-(1-(R)-(3,5-(dimethyl)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;

- 288 -

- 563) 2-(R)-(1-(R)-(3,5-(dimethyl)phenyl)ethoxy)-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 5 564) 2-(R)-(1-(R)-(3-(fluoro)-5-(methyl)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 565) 2-(R)-(1-(R)-(3-(fluoro)-5-(methyl)phenyl)ethoxy)-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 10 566) 2-(R)-(1-(R)-(3-(chloro)-5-(methyl)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 567) 2-(R)-(1-(R)-(3-(chloro)-5-(methyl)phenyl)ethoxy)-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 15 568) 2-(R)-(1-(R)-(3-(bromo)-5-(methyl)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 569) 2-(R)-(1-(R)-(3-(bromo)-5-(methyl)phenyl)ethoxy)-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 20 570) 2-(R)-(1-(R)-(3-(isopropoxy)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 25 571) 2-(R)-(1-(R)-(3-(isopropoxy)phenyl)ethoxy)-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 572) 2-(R)-(1-(R)-(3-(isopropoxy)-5-(trifluoromethyl)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 30 573) 2-(R)-(1-(R)-(3-(isopropoxy)-5-(trifluoromethyl)phenyl)ethoxy)-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;

- 289 -

- 574) 2-(R)-(1-(R)-(3-(chloro)-5-(isopropoxy)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 5 575) 2-(R)-(1-(R)-(3-(chloro)-5-(isopropoxy)phenyl)ethoxy)-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 576) 2-(R)-(1-(R)-(3-(fluoro)-5-(isopropoxy)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 10 577) 2-(R)-(1-(R)-(3-(fluoro)-5-(isopropoxy)phenyl)ethoxy)-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 578) 2-(R)-(1-(R)-(3-(t-butyl)-5-(chloro)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 15 579) 2-(R)-(1-(R)-(3-(t-butyl)-5-(chloro)phenyl)ethoxy)-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 580) 2-(R)-(1-(R)-(3-(t-butyl)-5-(trifluoromethyl)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 20 581) 2-(R)-(1-(R)-(3-(t-butyl)-5-(trifluoromethyl)phenyl)ethoxy)-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 582) 2-(R)-(1-(R)-(3,5-(dimethyl)-4-(fluoro)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 25 583) 2-(R)-(1-(R)-(3,5-(dimethyl)-4-(fluoro)phenyl)ethoxy)-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 30 584) 2-(R)-(1-(R)-(3,5-(dimethyl)-4-(chloro)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(-1H,4H-1,2,4-triazolo)methyl)-morpholine;

- 290 -

- 585) 2-(R)-(1-(R)-(3,5-(dimethyl)-4-(chloro)phenyl)ethoxy)-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 5 586) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)-4-(fluoro)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 587) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)-4-(fluoro)phenyl)ethoxy)-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 10 588) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)-4-(chloro)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 589) 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)-4-(chloro)phenyl)ethoxy)-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 15 590) 2-(R)-(1-(R)-(3,5-(dichloro)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(-1H,4H-1,2,4-triazolo)methyl)-morpholine;
- 591) 2-(R)-(1-(R)-(3,5-(dichloro)phenyl)ethoxy)-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;
- 20 592) 2-(R)-(1-(R)-(3,5-(difluoro)phenyl)ethoxy)-3-(S)-phenyl-4-(3-(1,2,4-triazolo)methyl)-morpholine;
- 25 593) 2-(R)-(1-(R)-(3,5-(difluoro)phenyl)ethoxy)-3-(S)-phenyl-4-(4-(2-oxo-1,3-imidazolo)methyl)-morpholine;

or a pharmaceutically acceptable salt thereof.

- 291 -

16. A pharmaceutical composition comprising a pharmaceutically acceptable carrier and an effective amount of the compound of Claim 1.

5 17. A method for antagonizing the effect of substance P at its receptor site or for the blockade of neurokinin-1 receptors in a mammal which comprises the administration to the mammal of the compound of Claim 1 in an amount that is effective for antagonizing the effect of substance P at its receptor site in the mammal.

10 18. A method for antagonizing the effect of neurokinin A at its receptor site or for the blockade of neurokinin-2 receptors in a mammal which comprises the administration to the mammal of the compound of Claim 1 in an amount that is effective for antagonizing the effect of neurokinin A at its receptor site in the mammal.

15 19. A method of treating or preventing pain or nociception attributable to or associated with migraine in a mammal in need thereof which comprises the administration to the mammal of an effective amount of the compound of Claim 1.

20 20. A method of treating or preventing a condition selected from the group consisting of: diabetic neuropathy; peripheral neuropathy; AIDS related neuropathy; chemotherapy-induced neuropathy; and neuralgia, in a mammal in need thereof which comprises the administration to the mammal of an effective amount of the compound of Claim 1.

25 21. A method for the treatment or prevention of asthma in a mammal in need thereof which comprises the administration to the mammal of an effective amount of the compound of Claim 1, either alone or in combination with a neurokinin-2 receptor antagonist or with a β_2 -adrenergic receptor agonist.

30

- 292 -

22. A method for the treatment of cystic fibrosis in a mammal in need thereof which comprises the administration to the mammal of an effective amount of the compound of Claim 1.

5 23. A method for the treatment or prevention of emesis in a mammal in need thereof which comprises the administration to the mammal of an effective amount of the compound of Claim 1.

10

15


20

25

30

- 293 -

24. A process for the preparation of a compound of structural formula IV:



or a pharmaceutically acceptable salt thereof, wherein:

15 R¹ is selected from the group consisting of:

- (1) hydrogen;
 - (2) C₁-6 alkyl, unsubstituted or substituted with one or more of the substituents selected from:
 - (a) hydroxy,
 - (b) oxo,
 - (c) C₁-6 alkoxy,
 - (d) phenyl-C₁-3 alkoxy,
 - (e) phenyl,
 - (f) -CN,
 - (g) halo, wherein halo is fluoro, chloro, bromo or iodo,
 - (h) -NR⁹R¹⁰, wherein R⁹ and R¹⁰ are independently selected from:
 - (i) hydrogen,
 - (ii) C₁-6 alkyl,
 - (iii) hydroxy-C₁-6 alkyl, and
 - (iv) phenyl,
 - (i) -NR⁹COR¹⁰, wherein R⁹ and R¹⁰ are as defined above,
 - (j) -NR⁹CO₂R¹⁰, wherein R⁹ and R¹⁰ are as defined above,
- 20
- 25
- 30

- 294 -

- (k) -CONR⁹R¹⁰, wherein R⁹ and R¹⁰ are as defined above,
- (l) -COR⁹, wherein R⁹ is as defined above,
- (m) -CO₂R⁹, wherein R⁹ is as defined above;
- 5 (n) heterocycle, wherein the heterocycle is selected from the group consisting of:
- (A) benzimidazolyl,
- (B) benzofuranyl,
- (C) benzothiophenyl,
- 10 (D) benzoxazolyl,
- (E) furanyl,
- (F) imidazolyl,
- (G) indolyl,
- (H) isooxazolyl,
- 15 (I) isothiazolyl,
- (J) oxadiazolyl,
- (K) oxazolyl,
- (L) pyrazinyl,
- (M) pyrazolyl,
- 20 (N) pyridyl,
- (O) pyrimidyl,
- (P) pyrrolyl,
- (Q) quinolyl,
- (R) tetrazolyl,
- 25 (S) thiadiazolyl,
- (T) thiazolyl,
- (U) thienyl,
- (V) triazolyl,
- (W) azetidinyll,
- 30 (X) 1,4-dioxanyl,
- (Y) hexahydroazepinyl,
- (Z) piperazinyl,
- (AA) piperidinyl,
- (AB) pyrrolidinyl,

- 295 -

(AC) tetrahydrofuranyl, and

(AD) tetrahydrothienyl,

and wherein the heterocycle is unsubstituted or substituted with one or more substituent(s) selected from:

(i) C₁₋₆ alkyl, unsubstituted or substituted with halo, -CF₃, -OCH₃, or phenyl,

(ii) C₁₋₆ alkoxy,

(iii) oxo,

(iv) hydroxy,

(v) thioxo,

(vi) -SR⁹, wherein R⁹ is as defined above,

(vii) halo,

(viii) cyano,

(ix) phenyl,

(x) trifluoromethyl,

(xi) -(CH₂)_m-NR⁹R¹⁰, wherein m is 0, 1 or 2, and R⁹ and R¹⁰ are as defined above,

(xii) -NR⁹COR¹⁰, wherein R⁹ and R¹⁰ are as defined above,

(xiii) -CONR⁹R¹⁰, wherein R⁹ and R¹⁰ are as defined above,

(xiv) -CO₂R⁹, wherein R⁹ is as defined above, and

(xv) -(CH₂)_m-OR⁹, wherein m and R⁹ are as defined above;

(3) C₂₋₆ alkenyl, unsubstituted or substituted with one or more of the substituent(s) selected from:

(a) hydroxy,

(b) oxo,

(c) C₁₋₆ alkoxy,

(d) phenyl-C₁₋₃ alkoxy,

(e) phenyl,

- 296 -

- (f) -CN,
(g) halo,
(h) -CONR⁹R¹⁰ wherein R⁹ and R¹⁰ are as defined above,
5 (i) -COR⁹ wherein R⁹ is as defined above,
(j) -CO₂R⁹, wherein R⁹ is as defined above,
(k) heterocycle, wherein the heterocycle is as defined above;
(4) C₂₋₆ alkynyl;
10 (5) phenyl, unsubstituted or substituted with one or more of the substituent(s) selected from:
(a) hydroxy,
(b) C₁₋₆ alkoxy,
(c) C₁₋₆ alkyl,
15 (d) C₂₋₅ alkenyl,
(e) halo,
(f) -CN,
(g) -NO₂,
(h) -CF₃,
20 (i) -(CH₂)_m-NR⁹R¹⁰, wherein m, R⁹ and R¹⁰ are as defined above,
(j) -NR⁹COR¹⁰, wherein R⁹ and R¹⁰ are as defined above,
(k) -NR⁹CO₂R¹⁰, wherein R⁹ and R¹⁰ are as defined above,
25 (l) -CONR⁹R¹⁰, wherein R⁹ and R¹⁰ are as defined above,
(m) -CO₂NR⁹R¹⁰, wherein R⁹ and R¹⁰ are as defined above,
30 (n) -COR⁹, wherein R⁹ is as defined above;
(o) -CO₂R⁹, wherein R⁹ is as defined above;

- 297 -

R² and R³ are independently selected from the group consisting of:

- (1) hydrogen,
- (2) C₁₋₆ alkyl, unsubstituted or substituted with one or more
of the substituents selected from:
 - (a) hydroxy,
 - (b) oxo,
 - (c) C₁₋₆ alkoxy,
 - (d) phenyl-C₁₋₃ alkoxy,
 - (e) phenyl,
 - (f) -CN,
 - (g) halo,
 - (h) -NR⁹R¹⁰, wherein R⁹ and R¹⁰ are as defined above,
 - (i) -NR⁹COR¹⁰, wherein R⁹ and R¹⁰ are as defined
above,
 - (j) -NR⁹CO₂R¹⁰, wherein R⁹ and R¹⁰ are as defined
above,
 - (k) -CONR⁹R¹⁰, wherein R⁹ and R¹⁰ are as defined
above,
 - (l) -COR⁹, wherein R⁹ is as defined above, and
 - (m) -CO₂R⁹, wherein R⁹ is as defined above;
- (3) C₂₋₆ alkenyl, unsubstituted or substituted with one or more
of the substituent(s) selected from:
 - (a) hydroxy,
 - (b) oxo,
 - (c) C₁₋₆ alkoxy,
 - (d) phenyl-C₁₋₃ alkoxy,
 - (e) phenyl,
 - (f) -CN,
 - (g) halo,
 - (h) -CONR⁹R¹⁰ wherein R⁹ and R¹⁰ are as defined
above,
 - (i) -COR⁹ wherein R⁹ is as defined above,
 - (j) -CO₂R⁹, wherein R⁹ is as defined above;

- 298 -

- (4) C2-6 alkynyl;
- (5) phenyl, unsubstituted or substituted with one or more of the substituent(s) selected from:
- (a) hydroxy,
 - (b) C1-6 alkoxy,
 - (c) C1-6 alkyl,
 - (d) C2-5 alkenyl,
 - (e) halo,
 - (f) -CN,
 - (g) -NO₂,
 - (h) -CF₃,
 - (i) -(CH₂)_m-NR⁹R¹⁰, wherein m, R⁹ and R¹⁰ are as defined above,
 - (j) -NR⁹COR¹⁰, wherein R⁹ and R¹⁰ are as defined above,
 - (k) -NR⁹CO₂R¹⁰, wherein R⁹ and R¹⁰ are as defined above,
 - (l) -CONR⁹R¹⁰, wherein R⁹ and R¹⁰ are as defined above,
 - (m) -CO₂NR⁹R¹⁰, wherein R⁹ and R¹⁰ are as defined above,
 - (n) -COR⁹, wherein R⁹ is as defined above;
 - (o) -CO₂R⁹, wherein R⁹ is as defined above;

and the groups R¹ and R² may be joined together to form a heterocyclic ring selected from the group consisting of:

- (a) pyrrolidinyl,
- (b) piperidinyl,
- (c) pyrrolyl,
- (d) pyridinyl,
- (e) imidazolyl,
- (f) oxazolyl, and
- (g) thiazolyl,

and wherein the heterocyclic ring is unsubstituted or

- 299 -

substituted with one or more substituent(s) selected from:

- (i) C₁₋₆alkyl,
- (ii) oxo,
- (iii) C₁₋₆alkoxy,
- 5 (iv) -NR⁹R¹⁰, wherein R⁹ and R¹⁰ are as defined above,
- (v) halo, and
- (vi) trifluoromethyl;

10 and the groups R² and R³ may be joined together to form a carbocyclic ring selected from the group consisting of:

- (a) cyclopentyl,
- (b) cyclohexyl,
- (c) phenyl,

15 and wherein the carbocyclic ring is unsubstituted or substituted with one or more substituents selected from:

- (i) C₁₋₆alkyl,
- (ii) C₁₋₆alkoxy,
- 20 (iii) -NR⁹R¹⁰, wherein R⁹ and R¹⁰ are as defined above,
- (iv) halo, and
- (v) trifluoromethyl;

25 and the groups R² and R³ may be joined together to form a heterocyclic ring selected from the group consisting of:

- (a) pyrrolidinyl,
- (b) piperidinyl,
- (c) pyrrolyl,
- (d) pyridinyl,
- 30 (e) imidazolyl,
- (f) furanyl,
- (g) oxazolyl,
- (h) thienyl, and
- (i) thiazolyl,

- 300 -

and wherein the heterocyclic ring is unsubstituted or substituted with one or more substituent(s) selected from:

- (i) C₁₋₆alkyl,
- (ii) oxo,
- (iii) C₁₋₆alkoxy,
- (iv) -NR⁹R¹⁰, wherein R⁹ and R¹⁰ are as defined above,
- (v) halo, and
- (vi) trifluoromethyl;

X is selected from the group consisting of:

- (1) -O-,
- (2) -S-,
- (3) -SO-, and
- (4) -SO₂-;

R⁶, R⁷ and R⁸ are independently selected from the group consisting of:

- (1) hydrogen;
- (2) C₁₋₆ alkyl, unsubstituted or substituted with one or more of the substituents selected from:
 - (a) hydroxy,
 - (b) oxo,
 - (c) C₁₋₆ alkoxy,
 - (d) phenyl-C₁₋₃ alkoxy,
 - (e) phenyl,
 - (f) -CN,
 - (g) halo,
 - (h) -NR⁹R¹⁰, wherein R⁹ and R¹⁰ are as defined above,
 - (i) -NR⁹COR¹⁰, wherein R⁹ and R¹⁰ are as defined above,
 - (j) -NR⁹CO₂R¹⁰, wherein R⁹ and R¹⁰ are as defined above,
 - (k) -CONR⁹R¹⁰, wherein R⁹ and R¹⁰ are as defined above,

- 301 -

- (l) -COR⁹, wherein R⁹ is as defined above, and
(m) -CO₂R⁹, wherein R⁹ is as defined above;
- (3) C₂-6 alkenyl, unsubstituted or substituted with one or more
of the substituent(s) selected from:
- 5 (a) hydroxy,
(b) oxo,
(c) C₁-6 alkoxy,
(d) phenyl-C₁-3 alkoxy,
(e) phenyl,
10 (f) -CN,
(g) halo,
(h) -CONR⁹R¹⁰ wherein R⁹ and R¹⁰ are as defined
above,
(i) -COR⁹ wherein R⁹ is as defined above,
15 (j) -CO₂R⁹, wherein R⁹ is as defined above;
- (4) C₂-6 alkynyl;
- (5) phenyl, unsubstituted or substituted with one or more of the
substituent(s) selected from:
- 20 (a) hydroxy,
(b) C₁-6 alkoxy,
(c) C₁-6 alkyl,
(d) C₂-5 alkenyl,
(e) halo,
(f) -CN,
25 (g) -NO₂,
(h) -CF₃,
(i) -(CH₂)_m-NR⁹R¹⁰, wherein m, R⁹ and R¹⁰ are as
defined above,
(j) -NR⁹COR¹⁰, wherein R⁹ and R¹⁰ are as defined
above,
30 (k) -NR⁹CO₂R¹⁰, wherein R⁹ and R¹⁰ are as defined
above,
(l) -CONR⁹R¹⁰, wherein R⁹ and R¹⁰ are as defined
above,

- 302 -

- (m) $-\text{CO}_2\text{NR}^9\text{R}^{10}$, wherein R^9 and R^{10} are as defined above;
- (n) $-\text{COR}^9$, wherein R^9 is as defined above;
- (o) $-\text{CO}_2\text{R}^9$, wherein R^9 is as defined above;
- 5 (6) halo,
- (7) $-\text{CN}$,
- (8) $-\text{CF}_3$,
- (9) $-\text{NO}_2$,
- 10 (10) $-\text{SR}^{14}$, wherein R^{14} is hydrogen or C_{1-5} alkyl,
- (11) $-\text{SOR}^{14}$, wherein R^{14} is as defined above,
- (12) $-\text{SO}_2\text{R}^{14}$, wherein R^{14} is as defined above,
- (13) $\text{NR}^9\text{COR}^{10}$, wherein R^9 and R^{10} are as defined above,
- (14) $\text{CONR}^9\text{COR}^{10}$, wherein R^9 and R^{10} are as defined above,
- (15) NR^9R^{10} , wherein R^9 and R^{10} are as defined above,
- 15 (16) $\text{NR}^9\text{CO}_2\text{R}^{10}$, wherein R^9 and R^{10} are as defined above,
- (17) hydroxy,
- (18) C_{1-6} alkoxy,
- (19) COR^9 , wherein R^9 is as defined above,
- 20 (20) CO_2R^9 , wherein R^9 is as defined above,
- (21) 2-pyridyl,
- (22) 3-pyridyl,
- (23) 4-pyridyl,
- (24) 5-tetrazolyl,
- 25 (25) 2-oxazolyl, and
- (26) 2-thiazolyl;

R^{11} , R^{12} and R^{13} are independently selected from the definitions of R^6 , R^7 and R^8 ;

30

- 303 -

Y is selected from the group consisting of:

- 5
- (1) a single bond,
 - (2) -O-,
 - (3) -S-,
 - (4) -CO-,
 - (5) -CH₂-,
 - (6) -CHR¹⁵-, and
 - (7) -CR¹⁵R¹⁶-, wherein R¹⁵ and R¹⁶ are independently
10 selected from the group consisting of:
 - (a) C₁₋₆ alkyl, unsubstituted or substituted with
one or more of the substituents selected from:
 - (i) hydroxy,
 - (ii) oxo,
 - (iii) C₁₋₆ alkoxy,
 - 15 (iv) phenyl-C₁₋₃ alkoxy,
 - (v) phenyl,
 - (vi) -CN,
 - (vii) halo,
 - (viii) -NR⁹R¹⁰, wherein R⁹ and R¹⁰ are as defined
20 above,
 - (ix) -NR⁹COR¹⁰, wherein R⁹ and R¹⁰ are as
defined above,
 - (x) -NR⁹CO₂R¹⁰, wherein R⁹ and R¹⁰ are as
defined above,
 - 25 (xi) -CONR⁹R¹⁰, wherein R⁹ and R¹⁰ are as
defined above,
 - (xii) -COR⁹, wherein R⁹ is as defined above, and
 - (xiii) -CO₂R⁹, wherein R⁹ is as defined above;
 - (b) phenyl, unsubstituted or substituted with one or more
30 of the substituent(s) selected from:
 - (i) hydroxy,
 - (ii) C₁₋₆ alkoxy,
 - (iii) C₁₋₆ alkyl,

- 304 -

- 5
- (iv) C₂₋₅ alkenyl,
(v) halo,
(vi) -CN,
(vii) -NO₂,
(viii) -CF₃,
(ix) -(CH₂)_m-NR⁹R¹⁰, wherein m, R⁹ and R¹⁰
are as defined above,
(x) -NR⁹COR¹⁰, wherein R⁹ and R¹⁰ are as
10 defined above,
(xi) -NR⁹CO₂R¹⁰, wherein R⁹ and R¹⁰ are as
defined above,
(xii) -CONR⁹R¹⁰, wherein R⁹ and R¹⁰ are as
defined above,
(xiii) -CO₂NR⁹R¹⁰, wherein R⁹ and R¹⁰ are as
15 defined above,
(xiv) -COR⁹, wherein R⁹ is as defined above, and
(xv) -CO₂R⁹, wherein R⁹ is as defined above;


20 Z is C₁₋₆ alkyl;

25

30

- 305 -

which comprises contacting a compound of formula V:



wherein R¹, R², R³, R⁶, R⁷, R⁸, R¹¹, R¹² and R¹³ are as defined above;

15 with an inorganic or an organic acid selected from the group consisting of:

toluenesulfonic acid, methanesulfonic acid, sulfuric acid,
hydrochloric acid and mixtures thereof,

in an aprotic solvent selected from the group consisting of:

20 toluene, benzene, dimethylformamide, tetrahydrofuran,
diethylether, dimethoxyethane, ethyl acetate, and mixtures thereof,


at a temperature from 0°C to solvent reflux temperature for a sufficient time to produce a compound of structural formula IV.

25

30

- 306 -

25. A process for the preparation of a compound of structural formula XI:



or a pharmaceutically acceptable salt thereof, wherein:

R¹ is selected from the group consisting of:

- 15
- (1) hydrogen;
 - (2) C₁-6 alkyl, unsubstituted or substituted with one or more of the substituents selected from:
 - (a) hydroxy,
 - (b) oxo,
 - (c) C₁-6 alkoxy,
 - (d) phenyl-C₁-3 alkoxy,
 - (e) phenyl,
 - (f) -CN,
 - (g) halo,
 - (h) -NR⁹R¹⁰, wherein R⁹ and R¹⁰ are independently selected from:
 - (i) hydrogen,
 - (ii) C₁-6 alkyl,
 - (iii) hydroxy-C₁-6 alkyl, and
 - (iv) phenyl,
- 20
- (i) -NR⁹COR¹⁰, wherein R⁹ and R¹⁰ are as defined above,
 - (j) -NR⁹CO₂R¹⁰, wherein R⁹ and R¹⁰ are as defined above,
- 25
- 30

- 307 -

- (k) -CONR⁹R¹⁰, wherein R⁹ and R¹⁰ are as defined above,
- (l) -COR⁹, wherein R⁹ is as defined above,
- (m) -CO₂R⁹, wherein R⁹ is as defined above;
- (n) heterocycle, wherein the heterocycle is selected from the group consisting of:
- (A) benzimidazolyl,
 - (B) benzofuranyl,
 - (C) benzothiophenyl,
 - (D) benzoxazolyl,
 - (E) furanyl,
 - (F) imidazolyl,
 - (G) indolyl,
 - (H) isooxazolyl,
 - (I) isothiazolyl,
 - (J) oxadiazolyl,
 - (K) oxazolyl,
 - (L) pyrazinyl,
 - (M) pyrazolyl,
 - (N) pyridyl,
 - (O) pyrimidyl,
 - (P) pyrrolyl,
 - (Q) quinolyl,
 - (R) tetrazolyl,
 - (S) thiadiazolyl,
 - (T) thiazolyl,
 - (U) thienyl,
 - (V) triazolyl,
 - (W) azetidiny,
 - (X) 1,4-dioxanyl,
 - (Y) hexahydroazepinyl,
 - (Z) piperazinyl,
 - (AA) piperidinyl,
 - (AB) pyrrolidinyl,

- 308 -

(AC) tetrahydrofuranyl, and

(AD) tetrahydrothienyl,

and wherein the heterocycle is unsubstituted or substituted with one or more substituent(s) selected from:

5

(i) C₁₋₆ alkyl, unsubstituted or substituted with halo, -CF₃, -OCH₃, or phenyl,

(ii) C₁₋₆ alkoxy,

10

(iii) oxo,

(iv) hydroxy,

(v) thioxo,

(vi) -SR⁹, wherein R⁹ is as defined above,

(vii) halo,

15

(viii) cyano,

(ix) phenyl,

(x) trifluoromethyl,

(xi) -(CH₂)_m-NR⁹R¹⁰, wherein m is 0, 1 or 2, and R⁹ and R¹⁰ are as defined above,

20

(xii) -NR⁹COR¹⁰, wherein R⁹ and R¹⁰ are as defined above,

(xiii) -CONR⁹R¹⁰, wherein R⁹ and R¹⁰ are as defined above,

(xiv) -CO₂R⁹, wherein R⁹ is as defined above, and

25

(xv) -(CH₂)_m-OR⁹, wherein m and R⁹ are as defined above;

(3) C₂₋₆ alkenyl, unsubstituted or substituted with one or more of the substituent(s) selected from:

30

(a) hydroxy,

(b) oxo,

(c) C₁₋₆ alkoxy,

(d) phenyl-C₁₋₃ alkoxy,

(e) phenyl,

(f) -CN,

(g) halo,

- 309 -

- 5
- (h) $-\text{CONR}^9\text{R}^{10}$ wherein R^9 and R^{10} are as defined above,
(i) $-\text{COR}^9$ wherein R^9 is as defined above,
(j) $-\text{CO}_2\text{R}^9$, wherein R^9 is as defined above,
(k) heterocycle, wherein the heterocycle is as defined above;
- (4) C2-6 alkynyl;
- 10 (5) phenyl, unsubstituted or substituted with one or more of the substituent(s) selected from:
- (a) hydroxy,
(b) C1-6 alkoxy,
(c) C1-6 alkyl,
(d) C2-5 alkenyl,
(e) halo,
15 (f) $-\text{CN}$,
(g) $-\text{NO}_2$,
(h) $-\text{CF}_3$,
(i) $-(\text{CH}_2)_m-\text{NR}^9\text{R}^{10}$, wherein m , R^9 and R^{10} are as defined above,
20 (j) $-\text{NR}^9\text{COR}^{10}$, wherein R^9 and R^{10} are as defined above,
(k) $-\text{NR}^9\text{CO}_2\text{R}^{10}$, wherein R^9 and R^{10} are as defined above,
(l) $-\text{CONR}^9\text{R}^{10}$, wherein R^9 and R^{10} are as defined above,
25 (m) $-\text{CO}_2\text{NR}^9\text{R}^{10}$, wherein R^9 and R^{10} are as defined above,
(n) $-\text{COR}^9$, wherein R^9 is as defined above;
30 (o) $-\text{CO}_2\text{R}^9$, wherein R^9 is as defined above;

R^2 and R^3 are independently selected from the group consisting of:

- (1) hydrogen,
(2) C1-6 alkyl, unsubstituted or substituted with one or more of the substituents selected from:

- 310 -

- 5 (a) hydroxy,
(b) oxo,
(c) C₁-6 alkoxy,
(d) phenyl-C₁-3 alkoxy,
(e) phenyl,
(f) -CN,
(g) halo,
(h) -NR⁹R¹⁰, wherein R⁹ and R¹⁰ are as defined above,
10 (i) -NR⁹COR¹⁰, wherein R⁹ and R¹⁰ are as defined above,
(j) -NR⁹CO₂R¹⁰, wherein R⁹ and R¹⁰ are as defined above,
(k) -CONR⁹R¹⁰, wherein R⁹ and R¹⁰ are as defined above,
15 (l) -COR⁹, wherein R⁹ is as defined above, and
(m) -CO₂R⁹, wherein R⁹ is as defined above;
(3) C₂-6 alkenyl, unsubstituted or substituted with one or more
of the substituent(s) selected from:
20 (a) hydroxy,
(b) oxo,
(c) C₁-6 alkoxy,
(d) phenyl-C₁-3 alkoxy,
(e) phenyl,
(f) -CN,
25 (g) halo,
(h) -CONR⁹R¹⁰ wherein R⁹ and R¹⁰ are as defined above,
(i) -COR⁹ wherein R⁹ is as defined above,
(j) -CO₂R⁹, wherein R⁹ is as defined above;
30 (4) C₂-6 alkynyl;
(5) phenyl, unsubstituted or substituted with one or more of the
substituent(s) selected from:
(a) hydroxy,
(b) C₁-6 alkoxy,

- 311 -

- 5 (c) C₁-6 alkyl,
(d) C₂-5 alkenyl,
(e) halo,
(f) -CN,
(g) -NO₂,
(h) -CF₃,
(i) -(CH₂)_m-NR⁹R¹⁰, wherein m, R⁹ and R¹⁰ are as defined above,
10 (j) -NR⁹COR¹⁰, wherein R⁹ and R¹⁰ are as defined above,
(k) -NR⁹CO₂R¹⁰, wherein R⁹ and R¹⁰ are as defined above,
(l) -CONR⁹R¹⁰, wherein R⁹ and R¹⁰ are as defined above,
15 (m) -CO₂NR⁹R¹⁰, wherein R⁹ and R¹⁰ are as defined above,
(n) -COR⁹, wherein R⁹ is as defined above;
(o) -CO₂R⁹, wherein R⁹ is as defined above;
20 and the groups R¹ and R² may be joined together to form a heterocyclic ring selected from the group consisting of:
(a) pyrrolidinyl,
(b) piperidinyl,
(c) pyrrolyl,
(d) pyridinyl,
25 (e) imidazolyl,
(f) oxazolyl, and
(g) thiazolyl,
and wherein the heterocyclic ring is unsubstituted or substituted with one or more substituent(s) selected
30 from:
(i) C₁-6alkyl,
(ii) oxo,
(iii) C₁-6alkoxy,
(iv) -NR⁹R¹⁰, wherein R⁹ and R¹⁰ are as defined above,

- 312 -

- (v) halo, and
- (vi) trifluoromethyl;

and the groups R² and R³ may be joined together to form a carbocyclic ring selected from the group consisting of:

5

- (a) cyclopentyl,
- (b) cyclohexyl,
- (c) phenyl,

and wherein the carbocyclic ring is unsubstituted or substituted with one or more substituents selected from:

10

- (i) C₁-6alkyl,
- (ii) C₁-6alkoxy,
- (iii) -NR⁹R¹⁰, wherein R⁹ and R¹⁰ are as defined above,
- (iv) halo, and
- (v) trifluoromethyl;

15

and the groups R² and R³ may be joined together to form a heterocyclic ring selected from the group consisting of:

20

- (a) pyrrolidinyl,
- (b) piperidinyl,
- (c) pyrrolyl,
- (d) pyridinyl,
- (e) imidazolyl,
- (f) furanyl,
- (g) oxazolyl,
- (h) thienyl, and
- (i) thiazolyl,

25

and wherein the heterocyclic ring is unsubstituted or substituted with one or more substituent(s) selected from:

30

- (i) C₁-6alkyl,
- (ii) oxo,
- (iii) C₁-6alkoxy,
- (iv) -NR⁹R¹⁰, wherein R⁹ and R¹⁰ are as defined above,

- 313 -

- (v) halo, and
- (vi) trifluoromethyl;

R⁶, R⁷ and R⁸ are independently selected from the group consisting of:

- 5 (1) hydrogen;
- (2) C₁₋₆ alkyl, unsubstituted or substituted with one or more of the substituents selected from:
 - (a) hydroxy,
 - 10 (b) oxo,
 - (c) C₁₋₆ alkoxy,
 - (d) phenyl-C₁₋₃ alkoxy,
 - (e) phenyl,
 - (f) -CN,
 - (g) halo,
 - 15 (h) -NR⁹R¹⁰, wherein R⁹ and R¹⁰ are as defined above,
 - (i) -NR⁹COR¹⁰, wherein R⁹ and R¹⁰ are as defined above,
 - (j) -NR⁹CO₂R¹⁰, wherein R⁹ and R¹⁰ are as defined above,
 - 20 (k) -CONR⁹R¹⁰, wherein R⁹ and R¹⁰ are as defined above,
 - (l) -COR⁹, wherein R⁹ is as defined above, and
 - (m) -CO₂R⁹, wherein R⁹ is as defined above;
- 25 (3) C₂₋₆ alkenyl, unsubstituted or substituted with one or more of the substituent(s) selected from:
 - (a) hydroxy,
 - (b) oxo,
 - 30 (c) C₁₋₆ alkoxy,
 - (d) phenyl-C₁₋₃ alkoxy,
 - (e) phenyl,
 - (f) -CN,
 - (g) halo,

- 314 -


- (h) $-\text{CONR}^9\text{R}^{10}$ wherein R^9 and R^{10} are as defined above,
- (i) $-\text{COR}^9$ wherein R^9 is as defined above,
- (j) $-\text{CO}_2\text{R}^9$, wherein R^9 is as defined above;
- 5 (4) C₂₋₆ alkynyl;
- (5) phenyl, unsubstituted or substituted with one or more of the substituent(s) selected from:
- (a) hydroxy,
- (b) C₁₋₆ alkoxy,
- 10 (c) C₁₋₆ alkyl,
- (d) C₂₋₅ alkenyl,
- (e) halo,
- (f) $-\text{CN}$,
- (g) $-\text{NO}_2$,
- 15 (h) $-\text{CF}_3$,
- (i) $-(\text{CH}_2)_m-\text{NR}^9\text{R}^{10}$, wherein m , R^9 and R^{10} are as defined above,
- (j) $-\text{NR}^9\text{COR}^{10}$, wherein R^9 and R^{10} are as defined above,
- 20 (k) $-\text{NR}^9\text{CO}_2\text{R}^{10}$, wherein R^9 and R^{10} are as defined above,
- (l) $-\text{CONR}^9\text{R}^{10}$, wherein R^9 and R^{10} are as defined above,
- (m) $-\text{CO}_2\text{NR}^9\text{R}^{10}$, wherein R^9 and R^{10} are as defined above,
- 25 (n) $-\text{COR}^9$, wherein R^9 is as defined above;
- (o) $-\text{CO}_2\text{R}^9$, wherein R^9 is as defined above;
- (6) halo,
- (7) $-\text{CN}$,
- 30 (8) $-\text{CF}_3$,
- (9) $-\text{NO}_2$,
- (10) $-\text{SR}^{14}$, wherein R^{14} is hydrogen or C₁₋₆alkyl,
- (11) $-\text{SOR}^{14}$, wherein R^{14} is as defined above,
- (12) $-\text{SO}_2\text{R}^{14}$, wherein R^{14} is as defined above,

- 315 -

- (13) $\text{NR}^9\text{COR}^{10}$, wherein R^9 and R^{10} are as defined above,
 (14) $\text{CONR}^9\text{COR}^{10}$, wherein R^9 and R^{10} are as defined above,
 (15) NR^9R^{10} , wherein R^9 and R^{10} are as defined above,
 (16) $\text{NR}^9\text{CO}_2\text{R}^{10}$, wherein R^9 and R^{10} are as defined above,
 5 (17) hydroxy,
 (18) C_{1-6} alkoxy,
 (19) COR^9 , wherein R^9 is as defined above,
 (20) CO_2R^9 , wherein R^9 is as defined above,
 10 (21) 2-pyridyl,
 (22) 3-pyridyl,
 (23) 4-pyridyl,
 (24) 5-tetrazolyl,
 (25) 2-oxazolyl, and
 15 (26) 2-thiazolyl;

R^{11} , R^{12} and R^{13} are independently selected from the definitions of
 R^6 , R^7 and R^8 ;
 Z is C_{1-6} alkyl;


20 which comprises contacting a compound of formula IX:



wherein R^1 , R^2 , R^3 , R^{11} , R^{12} and R^{13} are defined as above;
 30 with a hydride reducing agent selected from a group consisting of:
 diisobutylaluminum hydride: lithium tri(sec-butyl)borohydride: and
 lithium aluminum hydride; in an organic solvent at low temperature;
 followed by acylation of the resultant alcohol/alkoxide with a substituted
 benzoyl halide, substituted benzoic anhydride, substituted benzoic mixed
 anhydride or substituted activated benzoate ester (e.g. p-nitrophenyl

- 316 -

ester or N-hydroxysuccinimide ester), in which the phenyl ring of these acylating agents is substituted with R⁶, R⁷, and R⁸ as defined above, in an organic solvent at low temperature for a sufficient time to produce a compound of structural formula X:



15 and subsequently contacting the compound of formula X with a titanium
ylide (generated from reagents selected from: μ -chloro- μ -methylene-
(bis(cyclopentadienyl)titanium)dimethylaluminum; or dimethyl
titanocene; or the reagent prepared by the reduction of a 1,1-
dibromoalkane with zinc and titanium tetrachloride in the presence of
20 N,N,N',N'-tetramethylethylenediamine) to afford an enol ether which is
then hydrogenated in the presence of a catalyst selected from
palladium on carbon, platinum on carbon, or rhodium on carbon to
afford the compound of formula XI.

25

30

INTERNATIONAL SEARCH REPORT

Int. Application No
PCT/US 94/14497

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 C07D265/32 C07D279/12 C07D413/06 C07D413/04 C07D413/14

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 6 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP,A,0 528 495 (MERCK SHARP & DOHME LTD.) 24 February 1993 see claims	1-25
P,A	EP,A,0 577 394 (MERCK & CO., INC.) 5 January 1994 see claims & WO,A,94 00440 cited in the application	1-25
P,A	WO,A,94 19323 (MERCK SHARP & DOHME LTD.) 1 September 1994 see claims	1-25.

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

9 March 1995

Date of mailing of the international search report

23.03.95

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+31-70) 340-3016

Authorized officer

Chouly, J

INTERNATIONAL SEARCH REPORT

information on patent family members

Inter national Application No
PCT/US 94/14497

Patent document cited in search report	Publication date	Patent family member(s)		Publication date
EP-A-0528495	24-02-93	AU-A-	2413892	16-03-93
		CA-A-	2112397	04-03-93
		EP-A-	0600952	15-06-94
		WO-A-	9304040	04-03-93
		JP-T-	6510034	10-11-94

EP-A-0577394	05-01-94	AU-B-	4156893	06-01-94
		AU-B-	4656193	24-01-94
		CA-A-	2099233	30-12-93
		CN-A-	1087902	15-06-94
		FI-A-	946133	28-12-94
		JP-A-	6172178	21-06-94
		SI-A-	9300346	31-12-93
		WO-A-	9400440	06-01-94

WO-A-9400440	06-01-94	AU-B-	4156893	06-01-94
		AU-B-	4656193	24-01-94
		CA-A-	2099233	30-12-93
		CN-A-	1087902	15-06-94
		EP-A-	0577394	05-01-94
		FI-A-	946133	28-12-94
		JP-A-	6172178	21-06-94
		SI-A-	9300346	31-12-93

WO-A-9419323	01-09-94	AU-B-	6109394	14-09-94
